**ACTIVE INGREDIENT:** 

Captan N-[(trichloromethyl)thio] -4-cyclohexene-1, 2-dicarboximide

**INERT INGREDIENTS:** 

EPA REG. NO. 2749-295 NET WEIGHT

## CAUTION: KEEP OUT OF THE REACH OF CHILDREN. OF THE REACH OF CHILDREN. OF THE REACH OF CHILDREN. OF THE REACH OF CHILDREN.

MAY BE HARMFUL IF SWALLOWED. MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN. AVOID INHALATION OF DUST. AVOID CONTACT WITH SKIN, EYES OR CLOTHING.

IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES - GET MEDICAL ATTENTION.

**KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS.** DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORA-TING OR CRUSHING AND BURYING IN A SAFE PLACE.

THIS PRODUCT IS TOXIC TO FISH, BIRDS, AND OTHER WILDLIFE. KEEP OUT OF LAKES, STREAMS, OR PONDS. DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR, DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM AREAS TREATED. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIP-MENT, OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

## DIRECTIONS FOR USE

Captan 25 Seed Protectant is especially formulated for treatment of seed for the control of seed and soilborne diseases, including damping-off, seed decay organisms, and seedling blights, and is prepared for use directly in the planter or drill box.

The amount of seed protectant, as shown in the recommendations, should be poured over the seed in the planter or drill box and stirred with the seed for uniform distribution throughout the hopper. The amounts recommended will provide adequate coverage of the seed and any excess, that does not adhere to the seed coat, will flow with the seed into the furrow, giving additional protection against soil-borne diseases. May be used on previously treated seed.

Do not use treated seed for feed, food or oil purposes.

## RECOMMENDATIONS

SOYBEANS: Use Captan 25 Seed Protectant at the rate of 4 oz. per bu., or 7 oz. per 100 lbs. of seed as a planter box treatment.

-PEANUTS (Shalled): For the control of Asperigillus Niger and other seed and soil-borne diseases, use 2 to 4 oz. Captan 25 Seed Protectant per bushel, or 4 to 6 oz. per 100 lbs. of seed as a planter box treatment. WHEAT: Use Captan 25 Seed Protectant at the rate of 3 oz. per bu., or 4½ oz. per 100 lbs. of seed as a drill box treatment. This treatment is not recommended for the control of bunt (smut).

Manufactured for: J CHEMICAL CO., INC. AGRICULTURAL CHEMICALS DIV. ACETO 126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368

