

CAPTAN 25
SEED PROTECTANT
FOR SOYBEANS, PEANUTS AND WHEAT
15% NON-OIL LUBRICANT ADDED

ACTIVE INGREDIENT:
Captan N- (trichloromethyl)thio/ -4-cyclohexene-1,
2-dicarboximide
INERT INGREDIENTS:

2749-295
25%
ACCEPTED
TOTAL 100%
JAN 29 1973
EPA REG. NO. 2749-295
NET WEIGHT

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.

MAY BE HARMFUL IF SWALLOWED. MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN. AVOID INHALATION OF DUST. AVOID CONTACT WITH SKIN, EYES OR CLOTHING.

IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES - GET MEDICAL ATTENTION.

KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

THIS PRODUCT IS TOXIC TO FISH, BIRDS, AND OTHER WILDLIFE. KEEP OUT OF LAKES, STREAMS, OR PONDS. DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM AREAS TREATED. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DIRECTIONS FOR USE

Captan 25 Seed Protectant is especially formulated for treatment of seed for the control of seed and soil-borne diseases, including damping-off, seed decay organisms, and seedling blights, and is prepared for use directly in the planter or drill box.

The amount of seed protectant, as shown in the recommendations, should be poured over the seed in the planter or drill box and stirred with the seed for uniform distribution throughout the hopper. The amounts recommended will provide adequate coverage of the seed and any excess, that does not adhere to the seed coat, will flow with the seed into the furrow, giving additional protection against soil-borne diseases. May be used on previously treated seed.

Do not use treated seed for feed, food or oil purposes.

RECOMMENDATIONS

SOYBEANS: Use Captan 25 Seed Protectant at the rate of 4 oz. per bu., or 7 oz. per 100 lbs. of seed as a planter box treatment.

CAPTAN 25 SEED PROTECTANT FOR SOYBEANS, PEANUTS AND WHEAT 15% NON-OIL LUBRICANT ADDED

ACTIVE INGREDIENT:

Captan N, (trichloromethyl)thio, 4 cyclohexene-1,
2-dicarboximide

INERT INGREDIENTS:

	25%
	75%
TOTAL	100%

EPA REG. NO. 2749-295

NET WEIGHT

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

MAY BE HARMFUL IF SWALLOWED. MAY CAUSE IRRITATION OF EYES, NOSE,
THROAT AND SKIN. AVOID INHALATION. AVOID CONTACT WITH SKIN,
EYES OR CLOTHING.

IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES - GET
MEDICAL ATTENTION.

KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORA-
TING OR CRUSHING AND BURYING IN A SAFE PLACE.

THIS PRODUCT IS TOXIC TO FISH, BIRDS, AND OTHER WILDLIFE. KEEP OUT OF
LAKES, STREAMS, OR PONDS. DO NOT APPLY WHERE RUNOFF IS LIKELY TO
OCCUR. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM
AREAS TREATED. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIP-
MENT, OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON
THIS LABEL.

DIRECTIONS FOR USE

Captan 25 Seed Protectant is especially formulated for treatment of seed for the control of seed and soil-
borne diseases, including damping-off, seed decay organisms, and seedling blights, and is prepared for use
directly in the planter or drill box.

The amount of seed protectant, as shown in the recommendations, should be poured over the seed in the
planter or drill box and stirred with the seed for uniform distribution throughout the hopper. The amounts
recommended will provide adequate coverage of the seed and any excess, that does not adhere to the seed
coat, will flow with the seed into the furrow, giving additional protection against soil-borne diseases. May
be used on previously treated seed.

Do not use treated seed for feed, food or oil purposes.

RECOMMENDATIONS

SOYBEANS: Use Captan 25 Seed Protectant at the rate of 4 oz. per bu., or 7 oz. per 100 lbs. of seed as a
planter box treatment.

PEANUTS (Shell): For the control of *Aspergillus Niger* and other seed and soil-borne diseases, use 2 to 4
oz. Captan 25 Seed Protectant per bushel, or 4 to 6 oz. per 100 lbs. of seed as a planter box treatment.

WHEAT: Use Captan 25 Seed Protectant at the rate of 3 oz. per bu., or 4 1/2 oz. per 100 lbs. of seed as a drill
box treatment. This treatment is not recommended for the control of bunt (smut).



Manufactured for
ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICALS DIV.

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368