



# ATRAZINE 80W INDUSTRIAL WEED KILLER

FOR NON-SELECTIVE WEED CONTROL ON INDUSTRIAL SITES AND  
NON-CROP AREAS  
DO NOT USE ON AGRICULTURAL LANDS.

ACTIVE INGREDIENT: *April 25, 1974*  
Atrazine (3-Chloro-4-ethylamino-6-isopropyl-amino-s-triazine) 80%

INERT INGREDIENTS: *2749-371* 20%  
TOTAL 100%

EPA REG. NO. 2749-371  
NET WEIGHT

## CAUTION: KEEP OUT OF REACH OF CHILDREN.

HARMFUL IF SWALLOWED. AVOID CONTACT WITH EYES, PRO-  
LONGED CONTACT WITH SKIN, INHALATION OF DUST AND CON-  
TAMINATION OF FOOD AND FEED.

DO NOT CONTAMINATE DOMESTIC OR IRRIGATION WATER SUP-  
PLIES OR LAKES, STREAMS OR PONDS.

DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR  
DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED  
ON THIS LABEL.

DO NOT REUSE EMPTY CONTAINER. DESTROY WHEN EMPTY.

### GENERAL INFORMATION:

ATRAZINE 80W IS A NON-SELECTIVE HERBICIDE FOR USE ON  
NON-CROP AREAS. ATRAZINE 80W CAN BE APPLIED BEFORE OR  
AFTER PLANT GROWTH BEGINS. ALTHOUGH ATRAZINE 80W HAS  
SOME FOLIAGE ACTIVITY, MOST OF ITS ACTIVITY IS THROUGH  
ROOT ABSORPTION. THEREFORE, RAINFALL IS REQUIRED TO  
MOVE IT INTO THE ROOT ZONE. VERY DRY SOIL CONDITIONS  
AND LACK OF SUFFICIENT RAINFALL MAY RESULT IN POOR WEED

CONTROL. IF POST-EMERGENCE APPLICATIONS ARE MADE, WEEDS  
SHOULD BE YOUNG AND GROWING ACTIVELY.

USE IN AREAS WHERE LONG-TERM CONTROL OF BROADLEAF  
GRASS VEGETATION IS DESIRED, SUCH AS 1) RAILROAD,  
WAY AND UTILITY RIGHTS-OF-WAY; 2) PETROLEUM TANK F  
3) INDUSTRIAL SITES; 4) LUMBER YARDS; 5) AROUND  
HOUSES AND CHEMICAL PLANTS, AND 6) ALONG INDUS  
FENCE LINES. WHEN APPLIED TO THE SOIL, THIS PRO  
USUALLY INHIBITS PLANT GROWTH FOR ONE YEAR OR MO  
SHOULD NOT BE USED ON LAND TO BE CROPPED OR NEA  
JACENT DESIRABLE TREES, SHRUBS, OR PLANTS OR INJURY  
OCCUR.

ATRAZINE 80W IS NON-CORROSIVE TO EQUIPMENT. STORE  
ATRAZINE 80W IN A DRY PLACE.

### DIRECTIONS FOR USE:

Use sufficient water to assure thorough ground and plant coverage. Use at least  
of water for each pound of ATRAZINE 80W; more water may be practical when  
growth is heavy. Mechanical, bypass, or jet agitation is necessary to keep ATRAZINE  
80W in suspension during application.

To control annual broadleaf and grass weeds (such as ~~henry's grass~~, ~~chestnut~~  
~~grass~~, ~~lambquarters~~, ~~foxtail~~, ~~ragweed~~, ~~puncturevine~~) and biennials (such as  
~~mullein~~): Broadcast 6-12½ lbs. of ATRAZINE 80W per acre.

To control most annual and many perennial broadleaf and grass weeds (such as  
~~burdock~~, ~~Canada thistle~~, ~~dogfennel~~, ~~orchard grass~~, ~~plantain~~, ~~quackgrass~~, ~~pu~~  
~~redtop~~ and ~~smooth brome~~): Broadcast 12½-25 lbs. of ATRAZINE 80W per acre.

To control hard-to-kill biennial and perennial weeds (such as ~~bull thistle~~ and ~~sow~~  
~~thistle~~): Broadcast 25-50 lbs. of ATRAZINE 80W per acre.

For longer residual control in regions of high rainfall and a long growing season  
Broadcast 25-50 lbs. of ATRAZINE 80W per acre.

For small areas, 4½ ozs. per 1000 sq. ft. is equivalent to 12½ lbs. per acre.

The use of this product being beyond the control of the manufacturer, no guar-  
antee, expressed or implied, is made as to the effects of such or the results to be obtained  
if not used in accordance with directions or established safe practice. The buyer  
assumes all responsibility, including injury or damage, resulting from its misuse  
or in combination with other materials.



Manufactured for:  
ACETO CHEMICAL CO., INC.  
AGRICULTURAL CHEMICALS DIVISION

126 02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11355

2749-371

T



# ATRAZINE 80W INDUSTRIAL WEED KILLER

FOR NON-SELECTIVE WEED CONTROL ON INDUSTRIAL SITES AND  
~~IN-CROP AREAS~~  
DO NOT USE ON AGRICULTURAL LANDS.

ACTIVE INGREDIENT: *April 25, 1974*

Atrazine (3-Chloro-4-ethylamino-6-isopropyl-amino-s-triazine)	80%
INERT INGREDIENTS: <i>2749-371</i>	20%
TOTAL	100%

EPA REG. NO. 2749-371  
NET WEIGHT \_\_\_\_\_

## CAUTION: KEEP OUT OF REACH OF CHILDREN.

HAZARDOUS IF SWALLOWED. AVOID CONTACT WITH EYES, PROLONGED CONTACT WITH SKIN, INHALATION OF DUST AND CONTAMINATION OF FOOD AND FEED.

- ) NOT CONTAMINATE DOMESTIC OR IRRIGATION WATER SUPPLIES OR LAKES, STREAMS OR PONDS.
- ) NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.
- ) NOT REUSE EMPTY CONTAINER. DESTROY WHEN EMPTY.

GENERAL INFORMATION:  
ATRAZINE 80W IS A NON-SELECTIVE HERBICIDE FOR USE ON IN-CROP AREAS. ATRAZINE 80W CAN BE APPLIED BEFORE OR AFTER PLANT GROWTH BEGINS. ALTHOUGH ATRAZINE 80W HAS SOME FOLIAGE ACTIVITY, MOST OF ITS ACTIVITY IS THROUGH ROOT ABSORPTION. THEREFORE, RAINFALL IS REQUIRED TO MOVE IT INTO THE ROOT ZONE. VERY DRY SOIL CONDITIONS AND LACK OF SUFFICIENT RAINFALL MAY RESULT IN POOR WEED

CONTROL. IF POST-EMERGENCE APPLICATIONS ARE MADE, WEEDS SHOULD BE YOUNG AND GROWING ACTIVELY.

USE IN AREAS WHERE LONG-TERM CONTROL OF BROADLEAF AND GRASS VEGETATION IS DESIRED, SUCH AS 1) RAILROAD, HIGHWAY AND UTILITY RIGHTS-OF-WAY; 2) PETROLEUM TANK FARMS; 3) INDUSTRIAL SITES; 4) LUMBER YARDS; 5) AROUND WAREHOUSES AND CHEMICAL PLANTS, AND 6) ALONG INDUSTRIAL FENCE LINES. WHEN APPLIED TO THE SOIL, THIS PRODUCT USUALLY INHIBITS PLANT GROWTH FOR ONE YEAR OR MORE. IT SHOULD NOT BE USED ON LAND TO BE CROPPED OR NEAR ADJACENT DESIRABLE TREES, SHRUBS, OR PLANTS OR INJURY MAY OCCUR.

ATRAZINE 80W IS NON-CORROSIVE TO EQUIPMENT. STORE ATRAZINE 80W IN A DRY PLACE.

DIRECTIONS FOR USE:  
Use sufficient water to assure thorough ground and plant coverage. Use at least 1 gallon of water for each pound of ATRAZINE 80W; more water may be practical where weed growth is heavy. Mechanical, bypass, or jet agitation is necessary to keep ATRAZINE 80W in suspension during application.

To control annual broadleaf and grass weeds (such as ~~henry's grass, chestgrass, crabgrass, lambsquarters, foxtail, ragweed, puncturevine~~ and biennials (such as ~~turkey mullein~~): Broadcast 6-12½ lbs. of ATRAZINE 80W per acre.

To control most annual and many perennial broadleaf and grass weeds (such as ~~bluegrass, burdock, Canada thistle, dogfennel, orchard grass, plantain, quackgrass, purpletop, redtop and smooth brome~~): Broadcast 12½-25 lbs. of ATRAZINE 80W per acre.

To control hard-to-kill biennial and perennial weeds (such as ~~bull thistle and sow thistle~~): Broadcast 25-50 lbs. of ATRAZINE 80W per acre.

For longer residual control in regions of high rainfall and a long growing season, apply 25-50 lbs. of ATRAZINE 80W per acre.

For small areas, 4½ ozs. per 1000 sq. ft. is equivalent to 12½ lbs. per acre.

The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.



Manufactured by  
**ACETO CHEMICAL CO., INC**  
AGRICULTURAL CHEMICALS DIVISION  
126 02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368

2749-371



# ATRAZINE 80W INDUSTRIAL WEED KILLER

FOR NON-SELECTIVE WEED CONTROL ON INDUSTRIAL SITES AND  
NON-CROP AREAS

DO NOT USE ON AGRICULTURAL LANDS.

ACTIVE INGREDIENT:

Atrazine (3-Chloro-4-ethylamino-6-isopropyl-amino-s-triazine) 80%

INERT INGREDIENTS:

20%  
TOTAL 100%

EPA REG. NO. 2749-371  
NET WEIGHT

**CAUTION:** KEEP OUT OF REACH OF CHILDREN.

HARMFUL IF SWALLOWED. AVOID CONTACT WITH EYES, PROLONGED CONTACT WITH SKIN, INHALATION OF DUST AND CONTAMINATION OF FOOD AND FEED.

DO NOT CONTAMINATE DOMESTIC OR IRRIGATION WATER SUPPLIES OR LAKES, STREAMS OR PONDS.

DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DO NOT REUSE EMPTY CONTAINER. DESTROY WHEN EMPTY.  
GENERAL INFORMATION:

ATRAZINE 80W IS A NON-SELECTIVE HERBICIDE FOR USE ON NON-CROP AREAS. ATRAZINE 80W CAN BE APPLIED BEFORE OR AFTER PLANT GROWTH BEGINS. ALTHOUGH ATRAZINE 80W HAS SOME FOLIAGE ACTIVITY, MOST OF ITS ACTIVITY IS THROUGH ROOT ABSORPTION. THEREFORE, RAINFALL IS REQUIRED TO MOVE IT INTO THE ROOT ZONE. VERY DRY SOIL CONDITIONS AND LACK OF SUFFICIENT RAINFALL MAY RESULT IN POOR WEED