



ZIRAM 76 WP FUNG

Active Ingredient
Ziram
Inert Ingredients

Ziram 76 WP Fungicide
16.25%
76.00%
24.00%
100.00%

AC
JU
the
and
regis
1973

TO MIX - Fill the spray tank about 2/3 full of water and with the agitator running slowly sift in the required amount of this material. Finish filling the tank with water and let mixing continue in the tank until thoroughly mixed before applying, and then with the agitator running apply at once.

Almonds - Monilia Brown Rot Blossom blight and Shot Hole (Coryneum Blight) use 2 lbs per 100 gallons of water at pre-blossom, blossom and petal fall stages. Scab use 2 lbs in 100 gallons of water. Apply at petal fall. Repeat 5 weeks later. Use Precaution. Do not exceed 8.0 lbs per acre after petal fall. Do not exceed 16.44 lbs per acre up to petal fall.

Apples - Scab (West Coast only) - use 1.12 to 2 lbs in 100 gallons of water at preblossom, calyx and cover stages. Perennial Blight and Bullseye Rot (Fusicladium) - use 1.12 to 2 lbs in 100 gallons of water at pre-harvest prior to fall rains. Repeat if necessary, especially, if excessive rainfall occurs. Use Precaution. Apples may be injured by Zim compounds under certain conditions. Do not exceed 15.8 lbs per acre per application. Do not spray within 7 days of harvest.

Apricots - Shot Hole - Coryneum Blight - apply 1.12 to 2 lbs per 100 gallons of water at pre-blossom, blossom and petal fall stages and in early cover sprays. Use Precaution. Do not apply more than 12 lbs per acre at any one time.

Apricots - Monilia Brown Rot Blossom blight - apply 1.12 to 2 lbs per 100 gallons water at pre-blossom, blossom and petal fall stages. Check for leaf spots. Apply 1.12 to 2 lbs per 100 gallons water. Apply 1.12 to 2 lbs per 100 gallons water about two weeks after shuck fall. Do not apply more than 20 lbs per acre.

If in eyes flush with plants of water for 15 minutes. Get medical attention.

Hazards to Humans and Domestic Animals

Avoid contact with walls, eyes, or inhaled. May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin or clothing. Avoid breathing dust or spray mist. Wash thoroughly after handling.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water courses. Use equipment to dispose of wastes. Do not apply when weather conditions favor drift on the area treated. Do not apply when drift is likely to occur.

Keep away from fire, sparks and hot surfaces. Do not use in areas where fire is likely to occur. Do not use in areas where fire is likely to occur.

Do not use in areas where fire is likely to occur. Do not use in areas where fire is likely to occur. Do not use in areas where fire is likely to occur.

Pears - Scab (West Coast Only) - use 1.12 to 2 lbs in 100 gallons of water in pre-harvest sprays. Use Precaution. Do not exceed 14.3 lbs per acre. If excessive rain fall occurs after harvest.

Peaches - Monilia Brown Rot Leaf Curl and Shot-Hole (Coryneum Blight) - use 1.12 to 2 lbs per 100 gallons of water just before fall rain or in November while trees are yet dormant and dormant. Do not exceed 14.3 lbs per acre. Do not exceed 10 lbs/acre after residues by wiping or brushing.

Cucumbers, Melons - Scab and Shot-Hole - use 1.12 to 2 lbs per 100 gallons of water per acre. Application before disease is expected. Use Precaution.

Beans (Snap and Lima) - Anthracnose - use 1.12 to 2 lbs per 100 gallons per acre when plants are large. Make applications starting when true leaves appear. Do not exceed 5.0 lbs per acre. Do not apply 4 days of harvest.

Tomatoes - Anthracnose - Early - use 1.12 to 2 lbs per 100 gallons of water starting 2 to 3 weeks after fruiting. Repeat at 7 to 10 day intervals as needed. Do not exceed 5.0 lbs per acre.

EPA REG. NO. 274-433

EPA EST. NO.

best in field

ZIRAM 76 WP FUNGICIDE

Active Ingredient
 *Ziram
 Inert Ingredients

*Zinc dimethyl dithiocarbamate
 Zinc expressed as metallic

16.25%

76.00%
 24.00%
 100.00%

ACCEPTED
 JUL 2 1980
 274 UC4

about 2/3 full of water and
 lift in the required amount
 a tank with water and let
 it thoroughly mixed before
 running apply at once

Blossom blight and Shot
 per 100 gallons of water at
 all stages. Scab - use 2 lbs
 per 100 gallons of water at
 petal fall. Repeat 5 weeks
 later. Use 8.0 lbs per acre after
 petal fall up to petal fall

use 1 1/2 to 2 lbs in
 100 gallons of water at
 petal fall and cover stages.
 Rot (Fusarium) (Pacific
 States) - use 2 lbs per
 100 gallons of water
 Repeat if necessary, as
 necessary. Apply
 under certain conditions
 or application. Do not

If in eyes, flush with plenty of water for 15 minutes. Get medical attention

Hazards to Humans and Domestic Animals

May be harmful if swallowed or inhaled. May cause irritation of
 eyes, nose, throat and skin. Avoid contact with eyes, skin or clothing.
 Avoid breathing dust or spray mist. Wash thoroughly after handling.

This product is toxic to fish. Keep out of streams or ponds. Do not contaminate
 water by cleaning of equipment, or disposal of wastes. Do not apply when weather con-
 ditions favor drift from the areas treated. Do not apply when runoff is likely to occur.

Keep away from fire, sparks and hot surfaces

If in violation of Federal law, use this product in a manner inconsistent with its
 labeling

Do not contaminate water, food or feed by storage or disposal. Open dumping is pro-
 hibited. Pesticide spray mixture or residue may not be used or chemically reprocessed
 should be disposed of in a landfill approved for pesticides or buried in a safe place away
 from water supplies. Dispose of containers in an incinerator or landfill approved for
 pesticides containers or burn in a safe place. Consult Federal, State or local authorities
 for disposal of containers.

Pears Scab (West Coast Only) - Use 1 to 1-1/2 pounds in
 100 gallons of water in pre-blossom and early cover sprays
 Storage Rots (Oregon) - Use 1 to 1-1/2 lbs. in 100 gallons
 of water as pre-harvest sprays. Use prior to Fall rains and re-
 peat if excessive rain fall occurs. Do not apply within 7 days
 of harvest

Peaches Monilinia Brown Rot Apply 1 1/2 to 2 lbs. per
 100 gallons of water at full bloom, petal fall and cover stages.
 Leaf Curl and Shot Hole (Coryneum Blight) - Apply 1 1/2
 to 2 lbs per 100 gallons of water in dormant and popcorn
 stages. For West Coast use on peaches for leaf curl, apply
 just before fall rain or in November and repeat in 3 months
 while trees are yet dormant and again at popcorn stage. Use
 Precautions Do not exceed 14.8 lbs/acre prior to petal fall.
 Do not exceed 10 lbs/acre after petal fall. Remove excess
 residues by wiping or brushing

Cucumbers, Melons Scab and Anthracnose apply 2 lbs
 per 100 gallons of water per acre at 7 day intervals. Begin
 application before disease is expected to appear or at first
 sign of infection. Use Precaution. Do not exceed 5.0 lbs per
 acre

Beans (Snap and Lima) Anthracnose - apply 2.0 lbs. per
 100 gallons per acre when plants are small and at 3.0 lbs per
 100 gallons to large plants. Make applications at 7 day inter-
 vals starting when true leaves appear. Use Precaution. Do not
 exceed 5.0 lbs. per acre. Do not apply to snap beans within
 4 days of harvest

Tomatoes Anthracnose Early Blight Use 2.0 lbs per
 100 gallons of water starting 2 to 4 weeks after transplanting.
 Repeat at 7 to 10 day intervals as necessary. Use Precaution.
 Do not exceed 5.0 lbs per acre

NET WEIGHT 10 LBS

FRAISE NO

BEST DOCUMENT AVAILABLE