



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
2749-519

Date of Issuance:  
JUN 28 1996

NOTICE OF PESTICIDE:  
 Registration  
 Reregistration  
  
(under FIFRA, as amended)

Term of Issuance:  
Conditional

Name of Pesticide Product:  
CIPC 2 EC

Name and Address of Registrant (include ZIP Code):  
Aceto Agriculture Chemicals Corp.  
One Hollow Lane  
Lake Success, NY 11042

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes listed below before you release the product for shipment:

a. Add the phrase, "EPA Reg. No. 2749-519".

b. In the Environmental Hazards section reinstate "Do not contaminate water by disposal of equipment washwaters".

Signature of Approving Official:

Date:

JUN 28 1996



CIPC 2 EC

POTATO SPROUT INHIBITOR

ONE GALLON OF THIS PRODUCT WILL TREAT 2,000 CWT OF POTATOES

ACTIVE INGREDIENT:

\* Chlorpropham 25%

INERT INGREDIENTS:

75%

TOTAL: 100%

\*Isopropyl N-(3-chlorophenyl) carbamate

Contains 2 lbs. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS BELOW

EPA REG. NO. 2749-519 EPA EST NO. 37507-MT-1

NET CONTENTS \_\_\_\_\_ GALLONS

Manufactured For:

Aceto Agricultural Chemicals Corp.  
1 Hollow Lane  
Lake Success, New York 11042

ACCEPTED  
with COMMENTS  
In EPA Letter-Dated

JUN 28 1996

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

2749-519

PRECAUTIONARY STATEMENTS  
Hazards to Humans and Domestic Animals

CAUTION: Causes eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. This product may cause allergic skin reaction.

Remove contaminated clothing and wash clothing before reuse.

Statement of Practical Treatment

If Swallowed: If conscious, drink large quantities of water, then induce vomiting by administering an emetic such as Ipecac syrup. If an emetic is not available, vomiting should be induced by placing a finger far back in the throat. Administer additional water. Take immediately to a hospital or physician. If unconscious or in convulsions, take immediately to a hospital. DO NOT attempt to induce vomiting or give anything by mouth to an unconscious or convulsing person. If in eyes: Flush with plenty of water for at least 15 minutes. Get medical attention. If on skin: Wash with plenty of soap and water. Get medical attention.

Physical or Chemical Hazards

Flammable. Keep away from heat and open flame.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT CONTAMINATE WATER BY DISPOSAL OF EQUIPMENT WASHWATERS.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label.


STORAGE AND DISPOSAL

PROHIBITIONS: Keep container closed. Do not contaminate water, food or feed by storage or disposal. This product inhibits germination of seed potatoes.

manufacturer and/or seller, including, but not limited to, timing and method of application, variety and condition of stored crop, and mixtures or treatments with other chemicals. Wash equipment thoroughly with soap and water before using with seed potatoes.

**EQUIPMENT:** This product is usually applied as a 1% emulsion spray on the drying table, or similar station, after the potatoes have been washed. Automatic proportioners or injection systems that apply an equivalent rate of active ingredient may also be used. Equipment should be set to spray evenly across the rolls moving the potatoes. Rolling or tumbling action improves coverage compared to flat conveyor belts. Thorough coverage of the potato by this emulsion will provide the best sprout inhibition.

**APPLICATION:** For dilute spray, prepare a 1% emulsion by adding 1 gallon of this product to 24 gallons of water. Agitate enough to form a uniform emulsion. For best results, agitate or recirculate emulsion while in use. Apply at a rate of 1 quart of 1% emulsion per 20 bags of potatoes (100 pounds per bag). For proportioner/injector applications, use sufficient water for thorough coverage while maintaining the application rate of 1 gallon product per 2000 cwt. potatoes.



**DISCLAIMER**

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**STORAGE:** Avoid storage below freezing temperatures. Prolonged storage may cause crystal formation. Warming contents and agitation will restore material to usable condition, but do not heat above 120°F. Vent caps when using indirect heat such as warm water bath, and do not use direct heat or flame.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEM-TREC (800)424-9300.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Metal: Triple rinse (or equivalent). then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

### APPLICATION

This product is used for treating potatoes to inhibit sprouting after storage and washing. This product will prevent wound periderm formation of potatoes: therefore, it should be used only after bruises and cuts have healed (normally a minimum of 2 weeks).

#### DO NOT APPLY IN THE FIELD

**IMPORTANT NOTE:** Potatoes may deteriorate or otherwise be damaged by bacteria, fungi, disease, insects, old age effects such as internal sprouting, environmental factors (such as humidity, temperature and airflow), and other causes which are unrelated to the application of this product, and which may occur whether or not this product is applied to the potatoes.

**DO NOT USE ON SEED POTATOES.** This product will inhibit the sprouting of seed potatoes. The inhibition of sprouting at recommended rates is usually effective for up to a year. The time period of effective sprout inhibition may vary depending on many conditions beyond the control of the