

PM 22

2749-520

6/25/96

19 185



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
2749-520

Date of Issuance:
JUN 25 1996

NOTICE OF PESTICIDE:
 X Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:
Conditional

Name of Pesticide Product:
CIPC 98A

Name and Address of Registrant (include ZIP Code):
Aceto Agriculture Chemicals Corp.
One Hollow Lane
Lake Success, NY 11042

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Add the phrase "EPA Registration No. 2749-520" to your label before you release the product for shipment.
3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

Signature of Approving Official:

/s/

Date:

JUN 25 1996

page 2
EPA Reg. No. 2749-520

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (7505C)


Enclosure

385

CIPC 98A

RECD EPA/OPP/DPD

'96 JUN 18 10:19



AEROSOL GRADE - POTATO SPROUT INHIBITOR

Active ingredient: Chlorpropham* (Isopropyl N-(3-chlorophenyl) carbamate)	By Weight
Inert ingredients:	98.0
Total	100.0%

* Contains 9.66 pounds of active ingredient per gallon.

CAUTION

KEEP OUT OF REACH OF CHILDREN
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapors or dust. Avoid contact with skin, eyes, and clothing. In case of contact immediately flush with plenty of water. Get medical attention if symptoms persist.

Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

IN CASE OF CONTACT: Flush eyes and skin with plenty of water for several minutes. Get medical attention if eye or skin irritation occurs.

IF SWALLOWED: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger placed on the back of the victim's tongue. Give fluids until vomiting is clear. If victim is unconscious, do not induce vomiting or give anything by mouth.

IF INHALED: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

If in eyes: Flush with plenty of water for 15 minutes. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

EMERGENCY INFORMATION

For spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters.

Net Contents: 2.5 gallons.

EPA REG. NO. 2749- EPA EST. NO.

Manufactured For: **ACETO AGRICULTURAL CHEMICALS CORPORATION**
One Hollow Lane, Lake Success, NY 11042-1215

ACCEPTED
with COMMENTS
In EPA Letter Dated

6/25/96

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
2749-520

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER
INCONSISTENT WITH ITS LABELING

NOTICE


CIPC 98A is used as an aerosol for treating potatoes for sprout inhibition during storage.
If entry into the storage area is necessary during or immediately following application before the fog has settled, protective clothing and respirators must be worn.

- Do not apply in the field.
- Do not use on seed potatoes. This inhibition of sprouting at recommended rates is usually effective up to a year regardless of removal from storage.
- Do not allow vapors to come in contact with, or get near to, storage areas used for seed potatoes. Let six months elapse before using treated storage area for seed potatoes. Air system components (including ducts) and building must be thoroughly cleaned before area is used for storage of seed potatoes.

CIPC 98A will prevent periderm formation of potatoes, therefore, it should be used only after bruises and cuts have healed (normally minimum of two weeks.)

FORCED AIR DISTRIBUTION METHOD

Assemble unit as shown. Insert aerosol generator intake hose in CIPC 98A container.
Set air ducts for recirculation.
Place exhaust end of aerosol generator at center of plenum (air mixing chamber) pointing it in direction of air flow. This will assure the best possible distribution of CIPC 98A throughout the duct system.



TREATMENT OF STORAGE OR OTHER AREAS THAT DO NOT HAVE RECIRCULATION AIR SYSTEMS

PRIOR to placing the potatoes in the area to be treated, make the following preparations:
On the floor of the area, install an air duct approximately 12 inches by 12 inches running the length or width of the potato pile leaving a false wall space at both ends for air circulation. The ducts should be spaced 10 to 12 feet apart and can be perforated metal pipe, slotted wood construction or if the potatoes are in bags, by bridging a 12 inch space between two rows of bags in the bottom layer with bags placed crosswise the space.
At the end of each duct in the false wall space where the fog is to be introduced, place a squirrel cage fan positioned to force air through the duct. The exit end of the duct must be blocked to force the air up through the piled potatoes.

When the area is filled and ready to treat, the following steps should be taken:
Close off any ventilating systems.
Start the squirrel cage fans.
Introduce the fog as near as possible to the bottom of the false wall space containing the fans.
Operate the fans until the fog has settled.
Reactive the ventilation systems 4 hours following application or when the fog has settled.

NOTE: When treating small areas such as trailer trucks or railroad cars, it is recommended that low volume aerosol operators such as a "Swing Fog" be used.

APPLICATION

Application of **CHLORPROPHAM** can be made anytime after the curing period and before sprouting of potatoes occurs.

1. Apply at rate of 1 pound active ingredient per 1,000 bushels (600 cwt.). One gallon treats 10,000 bushels (6,000 cwt.). Treat according to volume of storage (see table below for conversion).
2. For low volume application, use **FORCED AIR** recirculation through the pile at rates up to 5.0 CFM per ton of potatoes. For other applications, use the lowest **FORCED AIR** recirculation available for the applicator. Check for uniform air distribution throughout the potato pile.
3. Keep storage closed during application. Continue to recirculate air in storage for at least 4 hours or until the fog has settled. Return storage to normal ventilation as soon as possible following application.
1 cwt. = 1.67 bushels = 2.5 cubic feet; 1 bushel = 60 pounds = 1.6 cubic feet

RECOMMENDED CHLORPROPHAM RATE

To calculate the rate needed use the following formula:

$$\% \text{ of Standard Application Rate} = \frac{(2 \times T) + [(5 \times M) - 5]}{100}$$

Where: Standard Application Rate =

1 pound active ingredient/600 cwt.

T = Storage Temperature

M = Number of Months Storage Time

Example Calculation: (Potatoes stored at 45°F for six (6) months)

$$\text{Required Rate} = \frac{(2 \times 45) + [(5 \times 6) - 5]}{100} = 115\%$$

/1 - Rates for 55°F and 60°F are for processing potatoes only. Chart assumes treatment soon after suberization time.

TIME MONTHS	STORAGE TEMPERATURE				
	40°F	45°F	50°F	55°F/1	60°F/1
1	80%	90%	100%	110%	120%
2	85%	95%	105%	115%	125%
3	90%	100%	110%	120%	130%
4	95%	105%	115%	125%	135%
5	100%	110%	120%	130%	140%
6	105%	115%	125%	135%	145%
7	110%	120%	130%	140%	150%
8	115%	125%	135%	145%	155%
9	120%	130%	140%	150%	160%
10	125%	135%	145%	155%	165%

EXTENDED STORAGE

If the potatoes are held in storage longer than originally anticipated, the potatoes may be retreated. Using the above chart, the retreatment dosage is equal to the total amount required for the extended storage time less the amount used for the original storage time.

For example, potatoes stored at 45°F were treated based on a two (2) month storage period. A decision was made to extend storage to five (5) months (three months in addition to original 2 months). The total extended storage would require a total of 110% of the standard rate, the original treatment would use 95% of the standard rate. In this case, retreatment would require 15% (110-95%) of the standard treatment.

STORAGE AND DISPOSAL

STORAGE

Keep container closed. Do not contaminate water, food, or feed by storage or disposal. This product inhibits germination of seed potatoes.

PESTICIDE DISPOSAL

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Do not reuse as a container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

CONDITIONS OF SALE

ACETO warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to manufacturer, and buyer assumes the risk for any such use.