



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

April 22, 2025

SENT BY EMAIL

John F. Wright
jwrightch@comcast.net
ACETO LIFE SCIENCES, L.L.C., D/B/A ACTYLIS
4 Tri Harbor Court
Port Washington NY, 11050

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - Update company names and correct typographical errors to the label.
Product Name: HALOSULFURON 75WDG HERBICIDE
Admin Number: 2749-528
EPA Receipt Date: 02/12/2025
Action Case Number: 00643934

Dear John F. Wright:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Jamey Shuler via email at shuler.jamey@epa.gov.

Sincerely,

Kable Bo Davis

Kable Bo Davis, Senior Advisor
FHB, RD
Office of Pesticide Programs

HALOSULFURON-METHYL	GROUP	2	HERBICIDE
---------------------	-------	---	-----------

[Information in brackets [] is optional]

HALOSULFURON 75WDG HERBICIDE

[Alternate brand names include Profine 75™ Herbicide, Halomax 75™ Herbicide]

HALOSULFURON 75WDG HERBICIDE is an herbicide for selective pre-emergent and post-emergent control of listed weeds including both broadleaf weeds and nutsedge in:

- alfalfa;
- artichoke;
- beans (including dry);
- bushberry Crop Subgroup 13-07B;
- caneberry (blackberry, loganberry, raspberry, (black and red), wild raspberry; cultivars varieties and/or hybrids of these) Crop Subgroup 13-07A;
- corn (field corn, field corn grown for seed, sweet corn and popcorn);
- cotton;
- fallow ground;
- fruits (cantaloupes, honeydews, Crenshaw melons, watermelons);
- lowbush blueberries Crop Subgroup 13-07B
- millet, proso;
- okra;
- pasture, rangeland & CRP forage grasses/hay Crop Group 17;
- pome fruit (apple; arazole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivar, varieties, and/or hybrids of these) Crop Group 11-10;
- grain sorghum (milo);
- rhubarb;
- rice;
- soybeans, seed;
- succulent shelled pea and bean subgroup (Any succulent shelled cultivar of bean (Phaaseolus) including lima bean, green; broad bean; succulent; (vigna including blackeyed pea, cowpea, southern pea, (Pisum) including green pea, and pigeon pea) Crop Subgroup 6B;
- small fruit vine climbing subgroup except fuzzy kiwi (amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these.) Crop Subgroup 13-07F;
- sugarcane;
- tree nuts (Almonds, Beechnuts, Brazil nuts, Butternuts, Cashews, Chestnuts, Chinquapins, Filberts, Hickory nuts, Macadamia nuts, Pecans, Pistachios, Walnuts {Black and English});
- tuberous and corm vegetables subgroup (arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; potato; sweet potato; tanier; turmeric; yam bean; yam, true Crop Subgroup 1C
- turfgrasses (established lawns, ornamental turfgrass, landscaped areas, commercial and residential turfgrass), and other non-crop sites (including airports, cemeteries, fallow areas, golf courses, landscaped areas, public recreation areas, residential property, roadsides, school grounds, sod or turf seed farms, sports fields, landscaped areas with established woody ornamentals, fairgrounds, race tracks, tennis courts, campgrounds and rights-of-way);
- vegetables (asparagus, cucurbit vegetables group, cucumbers, dry beans, fruiting vegetables group, peppers {chili and bell}, pumpkin, succulent snap beans, tomatoes and winter squash).

Read the entire label before using this product.

Use only according to label instructions.

[See [inside booklet] [back panel] for [First Aid], Precautionary Statements] [and] [Directions for use]]

[Peel here for complete directions for use and additional precautionary Statements]

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

NOTIFICATION

2749-528

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

04/22/2025

ACTIVE INGREDIENT:	% BY WT.
Halosulfuron-methyl.....	75%
OTHER INGREDIENTS:.....	25%
TOTAL	100%

EPA Reg. No. 2749-528	Net Contents: 10, 20 ounces
EPA Est. No. 065387-AR-001	0.03 ounces (0.9 grams) [water-soluble bag]
	1.3 ounces
	55 lbs. (25kg)

Manufactured for:
Aceto Life Sciences, L.L.C. [d/b/a Actylis](#)
4 Tri Harbor Court
Port Washington, NY 11050

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUSES MODERATE EYE IRRITATION. HARMFUL IF SWALLOWED. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid	
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call poison control center or physician for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call poison control center or physician immediately for treatment advice. Remove visible particles from mouth. Have person rinse mouth thoroughly with water, spit out rinse water. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or physician, or going for treatment. FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887.	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- long-sleeved shirt and long pants, and
- shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS:

When handlers use closed systems, or enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should:

- Remove PPE immediately after handling this product.
- Wash the outside of the gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FOR CHEMICAL SPILL, LEAK, FIRE, EXPOSURE OR MEDICAL EMERGENCY INVOLVING THIS PRODUCT
CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887.

ENVIRONMENTAL HAZARDS

This product is toxic to non-target vascular plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Ground Water Advisory: Halosulfuron-methyl is known to leach through soil into ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of halosulfuron-methyl from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Non-Target Organism Advisor This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Aceto Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forest, nurseries and green houses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during this restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves, made of nitrile rubber, neoprene rubber or polyethylene.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

Product Information

HALOSULFURON 75WDG HERBICIDE is a sulfonylurea herbicide that works by inhibition of acetolactate synthase (ALS). Many factors such as application rate, weed species, weed pressure, conditions of weeds including size and climatic factors impact the degree of weed control. Applications made to actively growing weeds at the early stages of development as described below will optimize performance. In post-emergent weed applications, early treatment is best to control the weeds vying (competing) with the crop. For residual control from early post-emergent treatments (in corn) a second application may be needed to control later germination of weeds.

HALOSULFURON 75WDG HERBICIDE is quick to act on targeted weeds by stunting growth allowing the crop to overtake the development of the targeted weeds. Once the development of the targeted weeds is stunted, the leaves and growing point begin to discolor and die. Complete control typically occurs within 7 to 14 days depending on the weed size, species and growing conditions. Depending on the stage and development of the targeted weeds, control generally takes place in 7 to 14 days.

Windblown Soil Particles

Halosulfuron 75WDG Herbicide has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying Halosulfuron 75WDG Herbicide if prevailing local conditions may be expected to result in off-site movement.

Weed Resistance Management

For resistance management, Halosulfuron 75WDG Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Halosulfuron 75WDG Herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

See specific crop Use Directions for maximum single application rate, annual maximum number of applications and amount of active ingredients.

To delay herbicide resistance, take one or more of the following steps:

Rotate the use of Halosulfuron 75WDG Herbicide or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as 002749-00528.20250211.Notification

the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.

Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

Users should scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

Users should report lack of performance to the registrant or their representative.

Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

Mixing Instructions

HALOSULURON 75WDG HERICIDE is a water dispersible granule designed to be diluted with water at the rates listed in the specific crop use directions. Fill the spray tank with approximately ½ of the desired volume with water or carrier. With the agitation operating, add the specified amount of the formulation as listed in the targeted crop use directions. Complete the filling process while maintaining agitation. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the carrier source. Add nonionic surfactant and other spray additives as the last ingredients in the tank. Allow time to fully disperse.

Since this product forms a suspension in water, it is important to maintain good agitation during mixing and spraying. If the spray suspension is allowed to settle for a short period of time, be sure to agitate the spray suspension for a minimum 10 minutes. Apply spray solutions within 24 hours after mixing.

Spray Additives

Spray additives such as nonionic surfactant (NIS), or Crop Oil Concentrate (COC) and liquid nitrogen fertilizer (e.g. 28-0-0) are used with HALOSULFURON 75 WDG HERBICIDE to improve performance. The typical nonionic surfactant contains a minimum of 80% NIS and is accepted by the EPA for use on food crops. The use rate is 0.25 to 0.5% NIS concentrate (1 to 2 quarts per 100 gallons of spray mixture). An alternative for the nonionic surfactant is a Crop Oil Concentrate. The typical Crop Oil Concentrate is a phytobland oil (petroleum) or crop origin (vegetable) based product that containing a minimum 14% surfactant to allow it to be miscible with water. The use rate for the Crop Oil Concentrate is 1% vol/vol (1 gallon per 100 gallons of spray mixture). NIS or COC is the only spray additives required to improve efficacy. Do not use both NIS and COC in the spray mixture. Use liquid nitrogen for those tank mix partners which required a liquid nitrogen additive to improve performance. Consult the tank mixture partner's labels for specific additive requirements and interactions. In place of the liquid nitrogen fertilizer, a high quality, spray grade ammonium sulfate (e.g. 21-0-0) is used at a use rate of 2 to 4 pounds per acre. Use either NIS or COC in the spray mixture.

For specific details, consult the use directions in crop section of the label.

Use Rate Equivalency

Since HALOSULFRUON 75 WDG HERBICIDE contains 75% active ingredient per lb. of product, the following table expresses the use rate equivalency of oz. of this product in term of lbs. halosulfuron-methyl on a per acre basis.

oz. of Product per acre	lbs. halosulfuron- methyl per acre
$\frac{1}{2}$	0.0235
$\frac{2}{3}$	0.031
$\frac{3}{4}$	0.035
1	0.047
$1\frac{1}{3}$	0.062
$1\frac{1}{2}$	0.070
2	0.094
$2\frac{2}{3}$	0.125
$5\frac{1}{3}$	0.250

Application Methods

Apply this product by ground or with aerial equipment to produce uniform coverage on growing weeds or soil to achieve consistent weed control.

Uniform, thorough spray coverage is important to achieve consistent weed control. Calibrate application equipment according to manufacturer's specifications. Use nozzle type arrangements that provide optimum spray distribution and maximum coverage while avoid contact to sensitive crop foliage.

Thoroughly clean application equipment immediately after use and prior to spraying a crop other than corn or grain sorghum. See Spray Equipment Cleanout section of this label for complete details.

Ground Applications

When HALOSULFURON 75WDG HERBICIDE is applied by ground equipment, use in a minimum of 10 gallons of water per acre for a broadcast application. In dense weed populations and thick canopy cover, higher spray volumes are necessary, e.g. 15 – 20 gallons of water per acre. Use the proper spray volume and nozzles that will ensure thorough and uniform coverage of the targeted weeds. Use directed applications to avoid contacting sensitive crop foliage. Select nozzles that will provide optimum spray volume, distribution and coverage at a pressure (psi) that minimizes spray drift. Inspect nozzle distribution during application to avoid streaking and overspray.

Rope-wick or Wiper Applications

A typical rope-wick or wiper applicator consists of an absorbent material made of burlap, canvas, rope, sponge, or absorbent pad plumbed into a pipe reservoir filled with the aqueous herbicide mixture. Maintain the moisture on the absorbent materials to allow for leaf wetness on targeted weeds, but not to a moisture level that allows for excess moisture to drip from the absorbent material. Selected equipment must be maintained and capable of preventing all contact of the herbicide mixture with the crop or soil.

To ensure adequate contact with the weeds, adjust the height of the wiper applicator so that no wiper contact point is less than 2 inches above the desirable vegetation. For optimum performance, a minimum of 6 inches above the desirable vegetation of the wiper applicator will provide adequate exposed of the weeds to the herbicide mixture. Poor contact occurs when weeds are growing in dense clumps, in areas of severe weed infestation, when weed height varies dramatically or when operator speeds are too great. Terrain must be considered when making wiper applications. Sloping ground can cause herbicide solution to migrate to one side, causing dripping on the lower end and drying of the wiper on the upper end of the applicator.

Use Precautions:

Due to decreased efficacy do not apply this product when weeds are wet.

Only prepare enough herbicide mixture to be utilized that day.

Avoid leaks or dripping of the herbicide solution onto the crop as contact of this product to desirable vegetation could result in plant injury or destruction.

Keep wiper surfaces clean.

Clean wiper parts promptly after using this product by thoroughly flushing with water.

When using a surfactant refer to the Spray Additives section of this label.

When Using Motorized Ground Equipment:

Prior to application determine the per acre output of the applicator. If the output rate is unknown it may be obtained by evaluating the output at ~100% weed density. Apply a minimum of 1 oz HALOSULFURON 75WDG HERBICIDE per acre by mixing the desired per acre rate of this product, in ratio with your determined per acre output. Do not exceed the maximum labeled rate for the target crop.

The applicator device will physically wipe this product directly onto the weed in between rows of crop plants (row middles) or over the top of crops for selectively controlling weeds. Operate wiper applicators at a ground speed of no greater than 5 miles per hour. To maintain performance applicator should control chemical application rate by adjusting travel speed to match weed density. In areas of dense weeds better results can be obtained when two applications are made in opposite directions. Refer to the specific crop section of this label for rates and directions for use.

Spot Treatment:

For spot treatment or application with a hand held device, mix 1/4 oz – 1 oz this product per 1 gallon of water. For best results, when using a hand held applicator, wipe the desired target weeds in a back and forth motion to ensure proper contact and coverage.

Aerial Applications

When HALOSULFURON 75WDG HERBICIDE is applied by air, use in a minimum 3 - 15 gallons of water per acre. Properly calibrate the spray equipment. Follow the Spray Drift Management guidelines presented below. Inspect nozzle distribution during application to avoid streaking, overspray and spray drift.

SPRAY DRIFT**Aerial Applications:**

- Do not release spray at a height greater than 10 ft. above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES**SPRAY DRIFT****Handheld Technology Applications:**

- Take precautions to minimize spray drift.

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

- Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift. When applying aerially to crops, do not release spray at a height greater than 10 ft. above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Sensitive areas:

Use pesticide products adjacent to sensitive areas only when there is minimal potential for drift or off-target movement, e.g. wind is blowing away from non-target crops, residential areas, known habitats for threatened or endangered species, etc.

In California (only), particularly sensitive crops are identified as cotton and prunes. In applications near these sensitive crops utilize the following buffer zones:

- Do not apply aerial applications within 4 miles of sensitive crops.
- Do not apply ground applications within 1 mile of sensitive crops except when wind direction during the application is away from sensitive crops. When wind direction during the ground application is away from sensitive crops, do not apply within 0.5 miles of sensitive crops.
- Do not apply Direct Dry Applications on rice by air within 360 feet of sensitive crops.

Spray Equipment Cleanout

The mix tank and spray equipment cleanout is an important stewardship activity to avoid injury to desirable crops. It is important to clean all mixing and spraying equipment immediately after use and before using pesticide products including HALOSULFURON 75WDG HERBICIDE. This is especially important prior to spraying a crop other than grain sorghum and corn.

To clean the spraying equipment, follow the procedure outlined below:

- Completely drain the mix tank and/or sprayer, and then wash thoroughly the tank, sprayer, boom and nozzles with clean water. Drain the system again.
- Fill the mixing or spray tank half full with clean water and add domestic ammonium, normally a 3% solution, at a dilution rate of 1% vol/vol ammonium or 1 gallon per 100 gallons of rinsate.
- Completely fill the tank(s) with additional clean water. Agitate and recirculate and flush out the boom and hoses. Let the system run for 10 – 15 minutes. Drain the system completely.
- Remove nozzles and screens and dislodge any visible solid material. Then soak them in a 1% vol/vol ammonium solution. Inspect the nozzles and screen and remove any visual residues.
- Repeat the above procedure for a second time.
- Flush the mix tank and/or sprayer, boom and hoses with clean water. Drain the system again and inspect for any visible residues. If present, repeat the cleaning cycle again.
- If the rinsate cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Tank Mixtures

To improve this product's effectiveness, apply in combination with other pesticide products that are registered for the same crop and application techniques.

A list of potential herbicide tank mixture partners is provided in the use direction section under each crop. This list is an example of products used but is not an all inclusive list. For current information on the best tank mixture partner in your area, consult with the local dealer, distributor or State Agricultural Extension service.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

If HALOSULFURON 75WDG HERBICIDE is to be tank mixed with other herbicides, conduct a compatibility test prior to mixing. Use a small container and mix all components in a small amount, usually 0.5 to 1qt. of spray. Combine all products in the same ratio and order of addition as in the proposed spray mixture. Observe the mixture for indication of incompatibility which usual occurs in 10 to 30 minutes after mixing. If incompatibility is observed, try changing the order of addition of the components. The guideline on tank mixture partners is driven by formulation type. Start with wettable powders (WP's) including water soluble bags (WSB's), water dispersible granules (WDG's), suspension concentrated (SC's) or flowable (F's), all with very good agitation. Next follow with water miscible concentrates and emulsifiable concentrates (EC's) before adding drift control additives, nonionic surfactants (NIS's) or crop oil concentrates (COC's). After vigorous agitation, there must be a homogeneous suspension. Let the final tank mixture stand and observe for any rapid settling or floating of components. If any indications of physical incompatibility develop, do not use this mixture for spraying.

Application Restrictions

- Do not use air assisted (air blast) sprayers to apply this product.
- Do not apply this product through any type of irrigation system.
- Do not apply when wind speed exceeds 15 mph.
- Do not apply more than 2 oz. of this product per acre per 12-month period (includes applications to the crop and to row middles/furrows) on crops except on fallow ground, field corn, sugar cane, tree nuts and turf.
- Do not apply more than 2⅔ oz. of this product (0.125 lbs. halosulfuron-methyl) per acre per year on fallow ground, field corn, sugar cane and tree nuts.
- Do not apply more than 5⅓ oz. of this product (0.25 lbs. halosulfuron-methyl) per acre per year on turf.
- Do not allow this product to drift outside of targeted area.
- Do not apply tank mixtures if the crop is under heavy stress due to drought, water-saturated soils, poor fertility (especially low nitrogen levels), hail, frost, insects or when the maximum daytime temperature is above 92° F.
- Do not use this product if the target weeds or crop are under stress due to drought, water saturated soils, low fertility (especially low nitrogen levels) or other poor growing conditions.

Application Precautions

- Avoid spraying when conditions favor rainfall or using overhead sprinkler irrigation within 4 hours of this application.
- Significant crop injury may occur when spray residue from broadcast application of this product over plastic mulch is concentrated in the plant hole by irrigation or rainfall. To minimize this potential injury, ensure that planting beds are crowned properly.
- Under cool and wet growing conditions that delay early seedling emergence, vigor or growth, this product may cause injury or crop failure. These conditions are likely to occur during the first planting of the season.
- Loss in effectiveness or crop injury may result if weeds are under drought, stress, disease or insect damage.
- The maturity of the treated crops may be delayed by use of this product.
- Soil or foliar-applied organophosphate insecticides applied on crops treated with this product, may increase the potential for crop injury and/or the severity of the crop injury.
- Increase in crop injury may result if the seeding depth is too shallow and excessive amounts of water (greater than 1 inch) from rainfall or sprinkler irrigation occurs.
- Use nozzles and pressures that minimize the production of fine particles that drift outside of the targeted area
- Apply this product to labeled crops (including cultivars and/or hybrids of these). However, not all hybrids/varieties have been tested for sensitivity to this product. For untested varieties, treat a small amount of the field and determine potential sensitivity to its use. To the extent consistent with applicable law, the user assumes responsibility for such use and any plant injury that may occur.
- Applications of this product may cause temporary yellowing or stunting of the crop.
- Observe resistant management guidelines, especially on tolerant weeds.
- In California and Arizona due to environmental conditions that delay degradation of this product, extend the crop rotation intervals on drip irrigated crops.
- When this product is applied over-the-top of a blooming crop, bloom loss may occur under certain environmental conditions.
- If rainfall or irrigation occurs within 4 hours after application, reduce effectiveness may occur.
- Avoid disturbing (e.g. cultivation) treated areas for at least 7 days following application.

For Best Performance

Many factors such as application rate, weed species, weed pressure, conditions of weeds including size and climatic conditions impact the degree of weed control. Applications made to actively growing weeds at the early stages of development as described below will optimize performance. In post-emergent weed applications, early treatment is best to control the weeds vying (competing) with the crop. For residual control from early post-emergent treatments (in corn) a second application may be needed to control later germination of weeds.

HALOSULFURON 75WDG HERBICIDE is quick to act on targeted weeds by stunting growth allowing the crop to overtake the development of the targeted weeds. Once the development of the targeted weeds is stunted, the leaves and growing point begin to discolor and die. Complete control typically occurs within 7 to 14 days depending on the weed size, species and growing conditions. Depending on the stage and development of the targeted weeds, control generally takes place in 7 to 14 days.

When using spray additives, carefully follow the listed use instructions.

- In pre-emergence applications:
 - If the targeted weeds are present prior to crop emergence, use a nonionic surfactant identified in the "Spray Additives" section of the label.
 - For optimum pre-emergent weed control, activate the soil moisture.
 - Pre-emergent weed control is improved by incorporating this product with irrigations ($\frac{1}{4}$ – $\frac{1}{2}$ inch maximum).
- In post-emergence applications:
 - Better control is obtained when applied early to actively growing, small (1-3 inches in height) broadleaf weeds. Large broadleaf weeds may not be adequately controlled.
 - Nutsedge plants are best controlled at the actively growing, 3 - 5 leaf stage.
 - After a post-emergence application, delay overhead sprinkler irrigation for 2 to 3 days.
 - If weeds are under drought, stress, disease, or insect damage, do not use.
- Under heavy weed infestation, use early before the weeds become too competitive with the crop.

- To control suppressed weeds, large weeds that exceed the size limitations, weeds that emerge after an application, or weed species not listed, cultivate the treated soil 7 – 10 days after a post-emergence application unless specified otherwise.
- Avoid disturbing (e.g. cultivation) treated areas for at least 7 days following application.
- Annual weeds may have multiple flushes of seedlings, or treated perennials may sometimes re-grow from underground stems or roots, depending upon rainfall and other environmental conditions. To maximize control of such weeds, apply a sequential application of this product.

Pre-emergent Weed Activity Table
HALOSULFURON 75 WDG HERBICIDE
by Weed Species

Common Name	Scientific Name	Control	Suppression	Directions
Amaranth, Spiny	<i>Amaranth spinosus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Cocklebur, common	<i>Xanthium strumarium</i>	YES		
Corn Spurry	<i>Spergula arvensis</i>	YES		
Dayflower	<i>Commelina erecta</i>	YES		
Eclipta	<i>Ecilpta prostrate</i>	YES		
Flatsedge, Rice	<i>Cyperus iria</i>		YES	
Galinsoga	<i>Galinsoga</i>	YES		
Goosefoot		YES		
Groundsel, common	<i>Senecio vulgaris</i>	YES		
Horseweed/Marestail	<i>Erigeron canadensis</i>	YES		
Jimsonweed	<i>Datura stramonium</i>	YES		
Kochia	<i>Kochia scoparia</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Ladysthumb	<i>Polygonum persicaria</i>	YES		
Lambsquarter, common	<i>Chenopodium album</i>	YES		
Mustard, wild	<i>Sinapis arevensis</i>	YES		
Nutsedge, Yellow	<i>Cyperus esculentus</i>		YES	Use higher specified rates for suppression
Nutsedge, Purple	<i>Cyperus rotundus</i>		YES	Use higher specified rates for suppression
Pigweed, redroot	<i>Amaranthus retrofractus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Pigweed, smooth	<i>Amaranthus hybridus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Purslane	<i>Portulaca oleracea</i>		YES	
Radish, wild	<i>Raphanus raphanistrum</i>	YES		
Ragweed, common	<i>Ambrosia artemisiifolia</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Shepardspurse	<i>capsella bursapastoris (L.) medicus</i>	YES		
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	YES		
Sunflower	<i>Helianthus annuus</i>	YES		
Velvetleaf	<i>Abutilon theophrasti</i>	YES		

¹ If ALS resistant weeds are present, use another mode of action herbicide registered on the crop against the target weeds alone or as a tank mixture partner.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

**Post-emergent Weed Activity Table
HALOSULFURON 75 WDG HREBICIDE
by Weed Species**

Common Name	Scientific Name	Control	Suppression	Directions
Amaranth, Spiny	<i>Amaranth spinosus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Barnyardgrass	<i>Echinochloa crusgalli</i>	YES		
Bindweed	<i>Calystegia sepium</i>	YES		
Burcucumber	<i>Sicyas angulatus</i>	YES	YES	
California Arrowhead	<i>Sagittaria ontvedensis</i>	YES		1-1½ ounce rate required.
Cocklebur, common	<i>Xanthium strumarium</i>	YES		
Corn Spurry	<i>Spergula arvensis</i>	YES		
Cupgrass, Woolly	<i>Eriochloa villosa</i>	YES		
Dayflower	<i>Commelina erecta</i>		YES	
Dogbane Hemp	<i>Apocynum annabinum</i>		YES	
Eclipta	<i>Ecilpta prostrate</i>		YES	
Flatsedge, Rice	<i>Cyperus iria</i>	YES		
Fleabane, Philadelphia	<i>Erigeron philadelphicus</i>	YES		
Foxtail, giant, yellow, green bristly		YES		
Galinsoga	<i>Galinsoga</i>	YES		
Golden Crownbeard	<i>Verbesina encliodes</i>	YES		
Goosefoot		YES		
Horsenettle	<i>Solanum carolinense</i>	YES		
Horsetail	<i>Equisetum</i>		YES	
Itchgrass	<i>Rottboelliacochinchinensis</i>	YES		
Jointvetch	<i>Aeschynomene</i>	YES		
Johnsongrass rhizome, seedling	<i>Sorghum halepense</i>	YES		
Kochia	<i>Kochia scoparia</i>		YES	Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Ladysthumb	<i>Polygonum persicaria</i>	YES		
Mallow, Venice	<i>Hibiscus trionum</i>	YES		
Milkweed, Common	<i>Asclepias syriaca</i>		YES	
Milkweed, honeyvine	<i>Ampelamus albidus</i>		YES	
Millet, Wild Proso	<i>Panicum miliaceum</i>	YES		
Morningglory, Ivy leaf	<i>Ipomoea hederacea</i>		YES	Use higher rates for suppression.
Morningglory, Tall	<i>Ipomoea purpurea</i>		YES	Use higher rates for suppression.
Mustard, wild	<i>Sinapis arevensis</i>	YES		
Nightshade, Black	<i>Solanum americanum</i>	YES		
Nutsedge, Yellow	<i>Cyperus esculentus</i>	YES		Heavy infestation requires sequential applications.
Nutsedge, Purple	<i>Cyperus rotundus</i>	YES		Heavy infestation requires sequential applications.
Oats		YES		
Panicum, Fall	<i>Panicum dichotomiflorum</i>	YES		
Panicum, Texas	<i>Panicum texanum</i>	YES		
Passionflower, Maypop	<i>Passiflora incarnata</i>	YES		

Pigweed, redroot	<i>Amaranthus retrofractus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Pigweed, smooth	<i>Amaranthus hybridus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Pokeweed, common	<i>Phytolacca</i>	YES		
Quackgrass	<i>Elytrigia repense</i>	YES		
Radish, wild	<i>Raphanus raphanistrum</i>	YES		
Ragweed, common	<i>Ambrosia artemisiifolia</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Ragweed, giant	<i>Ambrosia trifida</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Redstem	<i>Ammannia auriculata</i>	YES		1-1½ ounce rate required.
Ricefield Bulrush	<i>Scirpus mucronatus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Ryegrass, Italian	<i>Lolium Multiflorum</i>	YES		
Sandbur		YES		
Sesbania, Hemp	<i>Sesbania exaltata</i>	YES		
Shattercane	<i>Sorghum bicolor</i>	YES		
Signalgrass, broadleaf		YES		
Shepardspurse	<i>capsella bursapastoris (L.) medicus</i>		YES	
Sida, prickly		YES		
Smallflower	<i>Umbrellaplant</i>	YES		1-1½ ounce rate required.
Smartweed, Pennsylvania	<i>Polyfomum Pensylvanisum</i>	YES		
Sorghum Almum		YES		
Thistle, Canada	<i>Cirsium arvense</i>	YES		
Sunflower	<i>Helianthus annuus</i>	YES		
Velvetleaf	<i>Abutilon theophrasti</i>	YES		

¹ If ALS resistant weeds are present, use another mode of action herbicide registered on the crop against the target weeds alone or as a tank mixture partner.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

HALOSULFURON 75WDG HERBICIDE Crop/Use Site Index [Pages to be added in final version]

	Page
Alfalfa	14
Bushberry	16
Caneberry	17
Corn:	
Field corn,	22
Field corn grown for seed,	22
Sweet corn and popcorn	40
Cotton	19
Fallow ground	22
Fruits:	
Cantaloupes,	19
Honeydews,	19
Crenshaw melons,	19
Watermelons	46

Grain sorghum (milo)	26
Lowbush blueberries	17
Millet, proso	28
Pasture rangeland, CRP grasses/hay	29
Pome fruit	30
Rice	32
Soybeans, soybean seed	35
Small fruit vine climbing	34
Succulent shelled pea & bean	38
Sugarcane	38
Tree nuts	42
Turfgrasses including turf and sod or turf seed farms	44
Vegetables:	
Asparagus,	15
Artichoke	15
Chile and bell peppers,	18
Cucurbit vegetables group,	20
Cucumbers,	19
Dry beans,	21
Eggplant	26
Fruiting vegetables,	26
Okra	28
Peppers	26
Pumpkins and winter squash,	31
Rhubarb	34
Succulent snap beans,	37
Summer squash for processing,	40
Tomatoes	41
Tuberous and corn vegetables	43

The use rate for HALOSULFURON 75 WDG is expressed in terms of the oz. of this product by weight per acre as Rate Oz. /Acre.

The pre-harvest interval (PHI) is the required days between the last application of HALOSULFURON 75WDG HERBICIDE and the harvesting of the crop.

For the minimum acceptable intervals between the last application of HALOSULFURON 75WDG and the planting of a rotational crops, see the Crop Rotation Guideline section of this label.

If HALOSULFURON 75WDG HERBICIDE is utilized with a tank mixture partner(s), refer to the specific partner label(s) and observe all the precautionary statements and use directions including pre-harvest intervals, crop rotation restrictions, mixing and application instructions. Observe the most restrictive of the labeling limitations, precautions, directions and restrictions of all products used in mixtures.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
ALFALFA	$\frac{2}{3}$ - 1	14	CA, NM and AZ only. Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl), per acre per year. Do not exceed $\frac{3}{4}$ oz. of product (0.035 lbs. halosulfuron-methyl) per treated acre for Post-emergence Spot Treatment and Sequential Post-emergence Treatment for Nutsedge Control. Do not apply this product by rope-wick or wiper applicators.
For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds. For ground equipment, use a minimum of 20 gallons of water per acre.			

Post-emergence Broadcast:

Post-emergent weed control in established alfalfa. For broadcast applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds. Alfalfa must be in the field for a minimum of six months before application of this product. Crop injury is reduced by applying as soon as possible after removal of hay from the field and prior to irrigation. Delay irrigation for a minimum of 48 hours after treatment.

Post-emergence Spot Treatment:

Use a spot treatment application for localized control of emerged nutsedge. Use sufficient water volume to allow for uniform coverage of the weeds.

Sequential Post-emergence Treatments for Nutsedge Control:

To maximize the control of nutsedge, a second post-emergent spot spray is applied to the areas where nutsedge has re-grown or emerged. In this case, use a spot treatment application for localized control of emerged nutsedge. Use sufficient water volume to allow for uniform coverage of the weeds. This sequential treatment has the greater potential for growth and yield reduction.

Data indicates that after application of this product, alfalfa growth and yields will be reduced for one or more cuttings. Where re-growth exceeds 6 inches, a greater yield reduction occurs. Symptoms may be temporary. Follow all directions carefully to minimize potential reduced plant growth and yield.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
ARTICHOKE	1 -2	5	Do not apply by air. Do not make more than 2 applications per 12 month period. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per 12 month period. Do not apply this product by rope-wick or wiper applicators.
For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.			
Directed Broadcast Application: Use Rate: 1 – 2 oz. Apply this product to the ground on either side of the row and winter ditches. Avoid contact of this product with the planted crop.			
Row Middle Applications: Use Rate: 1 – 2 oz. Apply this product between rows of perennial artichokes for the control of nutsedge and labeled broadleaf weeds. For optimum nutsedge control, use when plants are in the 3-5 leaf stage. Make application when oxalis is in full bloom. Avoid contact of this product with the planted crop.			
PRECAUTIONS: For best results in post-emergence applications, use a NIS spray additive. This product may not control ALS-resistant weeds Use rates are broadcast per acre. Reduce rate and spray volume in proportion to area actually sprayed. If sprayed directly, this product may cause significant, temporary stunting and delay maturity of artichokes if sprayed directly.			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
ASPARAGUS	½ - 1½	1	Do not apply more than 1½ oz. of this product (0.070 lbs. halosulfuron-methyl) per application. Do not make more than 2 applications per year. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per year. Do not use NIS west of the Rocky Mountains.

		Do not apply this product by rope-wick or wiper applicators.
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>For use in nursery, transplanted crowns and established beds.</p> <p>Post-emergence Post-transplant: Apply this product to asparagus before or during the harvesting season. For first year transplants, apply no sooner than six weeks after fern emergence. This product applied during this time period may increase the potential for crop injury. The addition of adjuvants or tank mixture partners may accentuate the potential crop injury. Use NIS east of the Rocky Mountains to improve weed control.</p> <p>Post-harvest: Apply this product at the end of the harvest season. Under heavy nutsedge pressure, use sequential applications. Avoid contact with the fern which may cause temporary yellowing. Use a nonionic surfactant or crop oil concentrate in this time period. Use drop nozzles to direct the spray below the fern to allow for more complete coverage of target weeds for better control of nutsedge and other broadleaf weeds while minimizing crop injury.</p> <p>To enhance the control of nutsedge, use sequential applications during the cutting/harvest season, when the first flush of nutsedge appears in the 3-5 leaf stage, apply $\frac{3}{4}$ to 1 oz. product per acre. Make a sequential application by applying $\frac{3}{4}$ to 1 oz. of this product per acre at least 21-30 days later and up to lay-by to control later flushes of nutsedge. Apply this product post-harvest during the fern stage. Avoid contact with the fern which may cause temporary yellowing. Use drop nozzles to direct the spray below the fern to allow for more complete coverage of target weeds for better control of nutsedge while minimizing crop injury.</p>		

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
BUSHBERRY (excluding lowbush blueberries) Crop Subgroup 13-07B	$\frac{1}{2}$ - $\frac{3}{4}$ 1 – 4 year old bushes $\frac{1}{2}$ - 1 > 4 year old bushes	14	<p>Do not overlap spray swath which will increase the application rate in the treated area.</p> <p>Do not use on bushes in established fields that are less than 1 year old or to plants under stress.</p> <p>Do not apply to areas where water is known to pond for periods of time following rainfall.</p> <p>Do not contact foliage, especially green wood renewal canes, with this product. Uptake via contacted foliage will result in plant injury.</p> <p>Do not make applications sooner than 45 days apart.</p> <p>Do not apply to 'Elliott' variety bushes established less than four years</p> <p>Do not make more than 2 applications per 12 month period.</p> <p>Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per 12 month period.</p> <p>Do not apply this product by rope-wick or wiper applicators.</p>
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Pre-emergence and Post-emergence Directed Applications for control of listed broadleaf and annual grass weeds. Use Rate: $\frac{1}{2}$ - 1 oz. Use this product as a single spray or sequential directed spray to ground on either side of row. If small weeds are present, tank mix with a post-emergence broad-spectrum type herbicide to maximize and enhance the spectrum of broadleaf and grass control. If ground cover prevents contact with the soil, reduced or no residual herbicidal activity will result.</p> <p>Post-emergence directed application for control of nutsedge. Use Rate: $\frac{1}{2}$ - 1 oz. Use this product as a single directed spray application when nutsedge is fully emerged or as two sequential directed spray applications. Make the first directed spray application to the initial nutsedge flush when it has reached the 3-5 leaf stage. If a second treatment is needed, it may be applied later in the season directed to secondary nutsedge emergence. For best results, use a minimum rate of $\frac{3}{4}$ oz. and when the nutsedge plants are in the 3-5 leaf stage.</p>			

PRECAUTIONS:

Contact of this product with the blueberry bushes will result in temporary leaf chlorosis.
 Use of a shielded boom is suggested.
 This product may not control ALS-resistant weeds.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
LOWBUSH BLUEBERRIES Crop Subgroup 13-07B (Except CA)	½ - 1	14	<p>Do not overlap spray swath which will increase the application rate in the treated area.</p> <p>Do not use on bushes in established fields that are less than 1 year old or to plants under stress.</p> <p>Do not apply to areas where water is known to pond for periods of time following rainfall.</p> <p>Do not apply this product after the crop has progressed into budbreak or significant injury will occur.</p> <p>Do not make more than 1 application per 12 month period.</p> <p>Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre per 12 month period.</p> <p>Do not apply this product by rope-wick or wiper applicators.</p>
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Pre-emergence and Post-emergence Applications in vegetative (non-crop) year and broadcast application prior to breaking dormancy for control of listed weeds.</p> <p>Use Rate: ½ - 1 oz.</p> <p>Use this product as a single broadcast spray.</p> <p>If small weeds are present, tank mix with a post-emergence broad-spectrum type herbicide to maximize and enhance the spectrum of broadleaf and grass control.</p> <p>If ground cover prevents contact with the soil, reduced or no residual herbicidal activity will result.</p> <p>Applications applied 1 to 2 months prior to breaking dormancy will allow for better weed control.</p> <p>PRECAUTIONS:</p> <p>Overlapping boom swaths increases the potential for phytotoxicity including leaf yellowing, reddening, and/or stunting</p> <p>This product may not control ALS-resistant weeds.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
CANEBERRY (Blackberry; loganberry; raspberry, black and red; wild raspberry; cultivars, varieties and/or hybrids of these) Crop Subgroup 13-07A (For use in OR and WA only)	¾ - 1½	14	<p>Do not make applications sooner than 45 days apart.</p> <p>Do not overlap spray swath which will increase the application rate in the treated area.</p> <p>Do not use on bushes established that are less than 1 year old or to plants under stress.</p> <p>Do not apply to areas where water is known to pond for periods of time following rainfall.</p> <p>Do not contact foliage, especially green wood renewal canes, with this product. Uptake via contacted foliage will result in plant injury.</p> <p>Do not apply to developing primocanes in season until hardened off.</p> <p>Do not apply this product for pre-emergence weed control if excessive weed growth prevents contact of spray with the soil.</p> <p>Do not apply by air.</p> <p>Do not make more than 2 applications per 12 month period.</p> <p>Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per 12 month period.</p>

	1		Rope-wick or Wiper Application Row middle/furrow application – Apply 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre.
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Pre-emergence Directed Applications for control of listed broadleaf and annual grass weeds. Use Rate: $\frac{3}{4}$ - 1 $\frac{1}{2}$ oz. Use this product as a broadcast directed spray application to ground on either side of row. Make application up to and including primocane burndown. If ground cover prevents contact with the soil, reduced or no residual herbicidal activity will result.</p> <p>Pre-emergence and Post-emergence Directed Applications for control of listed broadleaf and annual grass weeds. Use Rate: $\frac{3}{4}$ - 1 $\frac{1}{2}$ oz. Use this product as a single spray or sequential directed spray to ground on either side of row. If small weeds are present, tank mix with a post-emergence broad-spectrum type herbicide to maximize and enhance the spectrum of broadleaf and grass control. If ground cover prevents contact with the soil, reduced or no residual herbicidal activity will result.</p> <p>Post-emergence directed application for control of nutsedge. Use Rate: $\frac{3}{4}$ - 1 $\frac{1}{2}$ oz. Use this product as a single directed spray application when nutsedge is fully emerged or as two sequential directed spray applications. Make the first directed spray application to the initial nutsedge flush when it has reached the 3-5 leaf stage. If a second treatment is needed, it may be applied later in the season directed to secondary nutsedge emergence. For best results, use a minimum rate of $\frac{3}{4}$ oz. and when the nutsedge plants are in the 3-5 leaf stage.</p> <p>PRECAUTIONS: For best results in post-emergence applications, use a NIS spray additive. Contact of this product with the caneberry bushes will result in temporary leaf chlorosis. Use of a shielded boom is suggested. This product may not control ALS-resistant weeds.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
CHILE AND BELL PEPPERS	$\frac{1}{2}$ -1	30	AZ, CA, NM, TX and OK only. Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application. Do not make more than 2 applications per year. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl), per acre per year, (includes applications to the crop and to row middles/furrows). Not all pepper varieties have been tested.
	1		Rope-wick or Wiper Application Row middle/furrow application – Apply 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre.
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use 20 gallons of water per acre.</p> <p>Direct-seeded: <i>Post-emergence</i> – Apply as a directed spray 28 days after planting, or when the plants have reached a minimum of six inches in height, but prior to flowering. For lighter textured soils with low organic matter, use the lower rate.</p> <p>Transplanted: <i>Post-transplant</i> – Apply as a directed spray 21 days after transplanting, or when the plants have reached a minimum of six inches in height, but prior to flowering.</p>			

Direct-seeded and Transplant: <i>Row Middle/Furrow Applications</i> – For the treatment of nutsedge and labeled broadleaf weeds, use this product between rows of direct-seeded or transplanted crop. Avoid contact of this product with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.			
Crop	Rate Oz./Acre	PHI	RESTRICTIONS
COTTON	½ - 1½	28	Do not apply more than 1½ oz. of this product (0.062 lbs. halosulfuron-methyl) per application. Do not apply more than 1½ oz. of this product (0.062 lbs. halosulfuron-methyl), per acre per year. Do not apply this product by rope-wick or wiper applicators.
For post-emergent weed control in emerged cotton. Apply this product as a directed spray in hooded equipment. Make application any time after cotton emergence until row closure prohibits the use of hooded spray equipment. Use this product any time after cotton emergence until row closure inhibits use of hooded spray equipment. The applicator is responsible for maintaining proper spray speed and equipment position so spray mist does not contact cotton plants.			
Crop	Rate Oz./Acre	PHI	RESTRICTIONS
CUCUMBERS (including pickles), Cantaloupes, Honeydews , Crenshaw Melons	½ - 1	30 57 57 57	Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application. Do not apply more than 2 applications per crop cycle. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per year, (includes applications to the crop and to row middles/furrows). Rope-wick or Wiper Application Row middle/furrow application – Apply 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre.
	1		
For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.			
Direct-seeded Bare ground: <i>Pre-emergence</i> – Use this product after planting, but before cracking. For lighter textured soils with low organic matter, use the lower rate. <i>Post-emergence</i> – Use after the crop has reached at least 3-5 true leaves but before first female flowers appear. Apply this product as an over-the-top application, a directed spray application, or with crop shields to minimize contact of this product with the crop.			
Direct-seeded Plastic mulch: <i>Pre-seeding</i> – For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-plant application under the plastic mulch. After final bed shaping and just prior to the installation of the plastic mulch, apply this product. No sooner than 7 days after the application and the installation of the plastic mulch, plant the seed crops into this treated area unless local conditions demonstrate safety at an earlier interval. For lighter textured soils with low organic matter, use the lower rate. <i>Post-emergence</i> – Use after the crop has at least 3-5 true leaves but before first female flowers appear. Apply as an over-the-top application, a directed spray application, or with crop shields to minimize contact of this product with the crop. When applications are made over plastic, concentration of this product in the planting hole may occur resulting in additional phytotoxicity. Do not use over-the-top applications on plastic in the Northeastern and Midwestern states.			
Transplanted Bare ground: <i>Pre-transplant</i> – For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-plant application under the plastic mulch. No sooner than 7 days after the application and the installation of the plastic mulch, plant the seed crops into this treated area unless local conditions demonstrate safety at an earlier interval. For lighter			

textured soils with low organic matter, use the lower rate. Treated soil in the transplant hole may result in crop injury. During the transplant process, take care to limit movement of soil.

Post-transplant – Use this product to transplants that are established and actively growing. Do not apply until plants are actively growing and in the 3-5 true leaf stage or no sooner than 14 days after transplanting unless local conditions demonstrate safety at an earlier interval, but before first female flowers appear. Apply this product as an over-the-top application, a directed spray application, or with crop shields to minimize contact of this product with the crop.

Transplanted Plastic mulch;

Pre-transplant – For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-plant application under the plastic mulch. After final bed shaping and just prior to the installation of the plastic mulch, apply this product. No sooner than 7 days after the application and the installation of the plastic mulch, transplant the crop into this treated area unless local conditions demonstrate safety at an earlier interval. For lighter textured soils with low organic matter, use the lower rate. Soil treated with this product in the transplant hole may result in crop injury. During the transplant process, take care to limit movement of soil.

Post-transplant – Use this product on transplants that are established and actively growing. Do not apply until plants are established and actively growing in the 3-5 true leaf stage or no sooner than 14 days after transplanting unless local conditions demonstrate safety at an earlier interval, but before first female flowers appear. Apply as an over-the-top application, a directed spray application, or with the crop shields to minimize contact of this product with the crop. Additional phytotoxicity may occur when applications are made over plastic due to concentration of product in the transplant hole. Do not use over-the-top applications on plastic in the Northeastern and Midwestern states.

Pre-emergence Sequential Treatment for Nutsedge Control - To maximize the control of nutsedge, a post-emergent spot spray is applied to the areas where nutsedge has re-grown or emerged. In this case, use a spot treatment application for localized control of emerged nutsedge. Do not exceed 1 oz. product per treated acre in these areas. Use sufficient water volume to allow for uniform coverage of the weeds. Avoid contact of this product with the planted crop.

Sequential Post-emergence Treatments for Nutsedge Control - To maximize the control of nutsedge, a second post-emergent spot spray is applied to the areas where nutsedge has re-grown or emerged. In this case, use a spot treatment application for localized control of emerged nutsedge. Allow a minimum of 21 days between applications. Do not exceed 1 oz. product per treated acre in these areas. Use sufficient water volume to allow for uniform coverage of the weeds. Avoid contact of this product with the planted crop.

Direct-seeded and Transplant:

Row Middle/Furrow Applications – For the treatment of nutsedge and labeled broadleaf weeds, use this product between rows of direct-seeded or transplanted crop. Avoid contact of this product with the planted crop. If plastic is used on the planted rows, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
OTHER COMMODITIES IN THE CUCURBIT VEGETABLES GROUP (Including summer squash, gourd)	½ - 1	See Text	Including summer squash, gourd (See text for PHI). Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application. Do not apply within 30 days of harvest for the squash/cucumber subgroup. Do not apply within 57 days of harvest for the melon subgroup. Do not apply more than 2 applications per crop cycle. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl), per acre per year.
	1		Rope-wick or Wiper Application Row middle/furrow application – Apply 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre.

Direct-seeded and Transplant:

Row Middle/Furrow Applications – For the treatment of nutsedge and labeled broadleaf weeds, use this product between rows of direct-seeded or transplanted crop. Avoid contact of this product with the planted crop. If plastic is used on the

planted row, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.			
Crop	Rate Oz./Acre	PHI	RESTRICTIONS
DRY BEANS	½ - 1	30	<p>Do not apply more than 1 oz. of this product (0.047 lb. halosulfuron-methyl) per application.</p> <p>Do not make more than 2 applications per year.</p> <p>Do not make more than 1 post-emergence application.</p> <p>In CA, only apply as post-emergence directed spray to Middle Row/Furrows.</p> <p>For Direct -Seeded Pre-emergences and Post-emergence applications, do not apply more than ¾ oz. of this product (0.031 lbs. halosulfuron-methyl) per application.</p> <p>Do not apply more than 2 oz. (0.094 lb. halosulfuron-methyl) per acre per 12-month period (includes applications to the crop and to Row Middle/Furrows).</p> <p>For Row Middle/Furrow applications, do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application.</p> <p>Do not apply this product by rope-wick or wiper applicators.</p>
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Direct-Seeded Pre-emergence: Use Rate: ½ - ¾ oz. Use this product after planting, but before cracking. For lighter textured soils with low organic matter, use the lower rate.</p> <p>Direct-Seeded Post-emergence: Use Rate: ½ - ¾ oz. Use this product when plants have 1-3 trifoliate leaves, but before flowering. For best results, apply to weeds less than 6 inches high. Use a nonionic surfactant (NIS)</p> <p>Not all hybrids/varieties have been tested for sensitivity to this product. For untested varieties, a small amount of the field should be sprayed to determine potential sensitivity to its use. To the extent consistent with applicable law, the user assumes responsibility for such use and any plant injury that may occur.</p> <p>Row Middle/Furrow Applications: Use Rate: ½ - 1 oz. Apply this product between rows of crop for the control of nutsedge and labeled broadleaf weeds. Avoid contact of this product with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.</p> <p>Tank Mixture Partner A tank mixture of HALOSULFURON 75WDG HERBICIDE partnered with Eptam® 7-E, EPA Reg. No. 10163-283 (EPTC) will provide a broader spectrum of weed control than either product used separately.</p> <p>For post-emergence grass control, use partners including but not limited to Targa®, EPA Reg. No. 33906-9 (quizalofop-P-ethyl) or other graminicides.</p> <p>It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.</p> <p>Use HALOSULFURON 75WDG HERBICIDE at a rate of ½ - ¾ oz. with labeled rate of Eptam® 7-E, EPA Reg. No. 10163-283 (EPTC) and incorporate into the soil at a depth of approximately 2 inches before planting. If any crust appears on the soil, break it up by lightly rotary hoeing during or shortly after the emergence of the beans.</p> <p>For lighter textured soils with low organic matter, use the lower rate.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
FALLOW GROUND	$\frac{2}{3}$ – 1 $\frac{1}{3}$		Do not apply more than 1 $\frac{1}{3}$ oz. of this product (0.062 lbs. halosulfuron-methyl) per application. Do not make more than 2 applications per year. Do not apply more than 2 $\frac{2}{3}$ oz. of this product (0.125 lbs. halosulfuron-methyl) per acre per year. Do not apply this product by rope-wick or wiper applicators.
Apply this product as a broadcast spray to fallow ground. For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil.			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
FIELD CORN AND FIELD CORN GROWN FOR SEED	$\frac{2}{3}$ – 1 $\frac{1}{3}$	30	Do not apply more than 1 $\frac{1}{3}$ oz. of this product (0.062 lbs. halosulfuron-methyl) per application. Do not make more than 2 applications per crop year. Do not apply more than 2 $\frac{2}{3}$ oz. of this product (0.125 lbs. halosulfuron-methyl) per acre per year. After application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage. Do not apply this product by rope-wick or wiper applicators.

If used alone, apply a broadcast spray over-the-top or with drop nozzles from the spike through lay-by stage of field corn. For large corn or dense competing canopy, use drop nozzles.
Avoid spraying an excessive amount directly over the rows and into the whorl of the corn stalk.

Corn Weed Height Activity Table

Weed Activity	Control		Suppression	
	$\frac{2}{3}$ oz.	1 - 1 $\frac{1}{3}$ oz.	$\frac{2}{3}$ oz.	1 - 1 $\frac{1}{3}$ oz.
Rate of Product	Inches	Inches	Inches	Inches
Burcucumber			1 - 3	4 - 12
Cocklebur, common	1 - 9	9 – 14		
Fleabane, Philadelphia	1 - 3			
Kochia ¹	1 - 3			3 - 6
Lambsquarter, common			1 - 2	
Mallow, Venice	1 - 3	4 - 12		
Milkweed, common			3 - 5	6 - 12
Milkweed, honeyvine		1 - 6	1 - 3	
Morningglory				1 - 3
Mustard, wild		4 - 6		
Nutsedge: yellow ²	3 - 6	3 - 12		
purple	3 - 6	3 - 12		
Passionflower, maypop	1 - 3			
Pigweed, redroot ^{1,3}	1 - 3	4 - 6		
Pokeweed, common	1 - 6			
Radish, wild		4 - 6		
Ragweed: common ¹	1 - 9	9 - 12		
Giant ¹	1 - 3	4 - 6		
Smartweed, Pennsylvania	1 - 2			
Sunflower, common	1 - 12	12 - 15		
Velvetleaf	1 - 9	9 - 12 ³		

¹ See Pre-emergent and Post-emergent Weed Activity Tables

² Heavy infestations of nutsedge require sequential applications. To prevent nutsedge from competing with the crop an earlier application is required.

³ For large velvetleaf and pigweed control apply with liquid nitrogen fertilizer (2 to 4 quarts per acre) plus crop oil concentrate or nonionic surfactant is suggested.

Tank Mixture Partners It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.	
2,4-D	See tank mixture partner label for its rates. NIS Apply broadcast spray on corn up to 8 inches tall. If corn exceeds 8 inches, use directed spray with drop nozzles. Broadleaf weeds. Avoid sprays onto corn leaves just after unfolding, as injury may occur. Apply during the period from corn emergence through the 5 leaf stage or 8 inches tall, whichever comes first.
Accent® Herbicide EPA Reg. No. 352-560 (nicosulfuron)	See tank mixture partner label for its rates. COC or NIS or ammonium nitrogen fertilizer (e.g. 28%). Apply broadcast spray or with drop nozzles on emerged corn up to 24 inches tall. (free standing) For corn 24 to 36 inches tall, use directed spray with drop nozzles. Annual broadleaf weeds and annual grasses. Avoid sprays directly into the whorls of large cornstalks. Refer to Accent® label for use restrictions on corn varieties.
Accent Gold® Herbicide EPA Reg. No. 352-612 (clopyralid, flumetsulam, nicosulfuron, rimsulfuron)	See tank mixture partner label for its rates. COC or ammonium nitrogen fertilizer (e.g. 28%). Apply broadcast spray on corn up to 12 inches tall. Annual broadleaf weeds and annual grasses. Do not apply to seed corn.
atrazine (various formulations)	See tank mixture partner label for its rates. COC Apply broadcast spray on corn up to 12 inches tall. Apply when broadleaf weeds are small (3 inches or less). Post-emergence control of labeled broadleaf weeds. Aids in the burndown and control of many grass weeds (1.5 inches or less) which have escaped pre-emergence herbicide treatments. Atrazine mixtures may result in reduced control (antagonism) of larger broadleaf weeds. Smaller weeds are easier to control.
dicamba or Clarity® Herbicide, EPA Reg. No. 7969-137 (dicamba, diglycolamine salt)	See tank mixture partner label for its rates. NIS Apply broadcast spray on corn from emergence up to 36 inches tall. Use lower dicamba rates or directed sprays on corn taller than 8 inches. Broadleaf weeds. Avoid direct sprays into the whorls of large cornstalks. Do not make applications after corn exceeds 36 inches or 15 days before tassel emergence, whichever comes first. COC may cause crop injury, especially with higher dicamba or Clarity® rates.
Beacon® Herbicide EPA Reg. No. 352-560 (nicosulfuron)	See tank mixture partner label for its rates. COC or NIS or ammonium nitrogen fertilizer (e.g. 28%). Apply broadcast spray or with drop nozzles on corn from 4 - 20 inches tall. For corn 20 – 36 inches tall to pre-tassel, use drop nozzles. Broader spectrum. Avoid spraying directly into whorls of larger corn. See your dealer or seed supplier representative for a list of susceptible hybrids.
Buctril® Herbicide EPA Reg. No. 264-437	See tank mixture partner label for its rates. NIS Apply broadcast spray on corn up to tassel emergence. For post-emergence control of annual broadleaf weeds. Leaf burn may occur.

(bromoxynil octanoate)	Use of COC or ammonium nitrogen fertilizer (e.g. 28%) may cause additional leaf burn.
Buctril® Herbicide EPA Reg. No. 264-437 (bromoxynil octanoate) plus atrazine	See tank mixture partner label for its rates. NIS Apply broadcast spray on corn up to 12 inches tall. For post-emergence control of annual broadleaf weeds. Leaf burn may occur. Use of COC or ammonium nitrogen fertilizer (e.g. 28%) may cause additional leaf burn.
Callisto® Herbicide EPA Reg. No. 100-1131 (mesotrione)	See tank mixture partner label for its rates. COC or ammonium nitrogen fertilizer (e.g. 28%). Apply broadcast spray or with drop nozzles on seed or field corn up to 30 inches tall or 8 leaf collars, whichever is more restrictive. Broader spectrum.
Distinct® Herbicide EPA Reg. No. 7969-150 (dicamba, sodium salt, diflufenzopyr-sodium)	See tank mixture partner label for its rates. NIS Apply broadcast spray or with drop nozzles on corn 4 - 36 inches tall, e.g. V ₂ to V ₁₀ stage or 15 days prior to tassel emergence, whichever comes first. For corn taller than 20 inches, use drop nozzles. Broader spectrum. Avoid sprays directly into the whorls of large cornstalks. Do not use COC.
glyphosate (various formulations)	See tank mixture partner label for its rates. NIS or spray grade ammonium sulfate at 17 lb./100 gal. Apply broadcast spray or with drop nozzles on Glyphosate Tolerant (GT) field corn up to 30 inches tall or 8 leaf collars, whichever is more restrictive. For GT field corn between 24 – 36 inches, use drop nozzles. For corn taller than 20 inches, use drop nozzles. For burndown of emerged annual grasses, broadleaf weeds and nutsedge. Check product formulation label for specific restrictions. For use ONLY on corn hybrids tolerant to glyphosate herbicide.
glyphosate (various formulations)	Use HALOSULFURON 75WDG HERBICIDE at ⅔ oz. For glyphosate, see product formulation label. NIS Apply broadcast spray. For pre-plant burndown of emerged annual grasses, broadleaf weeds and nutsedge. To improve burndown of broadleaf weed control use dicamba or 2,4-D. Use only on Pioneer IR corn hybrids.
Impact® Herbicide EPA Reg. No. 5481-524 (topramezone)	See tank mixture partner label for its rates. NIS (preferred) or COC or ammonium nitrogen fertilizer (e.g. 28%). Apply broadcast spray or with drop nozzles on seed or field corn up to 36 inches tall. For a density canopy, drop nozzles are preferred. Broader spectrum.
Liberty® 280SL Herbicide EPA Reg. No. 7969-829 (glufosinate)	See tank mixture partner label for its rates. Spray grade ammonium sulfate (17lb./100 gallons of spray mix). Apply broadcast spray or with drop nozzles on field corn up to 24 inches tall or 7 leaf collars whichever is more restrictive. For field corn taller than 24 inches up to 36 inches tall, use drop nozzles. Broadleaf weeds and annual grasses. Do not add NIS or COC. For use ONLY on corn hybrids tolerant to Liberty® Herbicide.
Marksman® Herbicide EPA Reg. No. 7969-136 (dicamba, potassium salt, atrazine)	See tank mixture partner label for its rates. NIS Apply broadcast spray on corn up to 8 inches tall. Broader spectrum. COC may cause crop injury.
Option® Corn Herbicide	See tank mixture partner label for its rates. COC or ammonium nitrogen fertilizer (e.g. 28%) or spray grade ammonium sulfate (17 lb./100 gal.).

EPA Reg. No. 352-560 (nicosulfuron)	Apply broadcast spray or with drop nozzles on field corn 4 – 16 inches tall e.g. V ₂ to V ₆ . For field corn taller than 16 up to 36 inches e.g. V ₆ to V ₁₀ , use drop nozzles. Broader spectrum. Do not apply Option® to seed corn. Avoid spraying directly into the whorls of large cornstalks.
Status® Herbicide EPA Reg. No. 7969-242 (nicosulfuron, rimsulfuron)	See tank mixture partner label for its rates. NIS Apply broadcast spray or with drop nozzles on corn up to 20 inches tall. For corn taller than 20 inches use drop nozzles. Broader spectrum. Do not use COC.
Steadfast® Herbicide EPA Reg. No. 352-608 (nicosulfuron, rimsulfuron)	See tank mixture partner label for its rates. COC (preferred) or NIS or ammonium nitrogen fertilizer (e.g. 28%) or spray grade ammonium sulfate (17 lb./100 gal.). Apply broadcast spray or with drop nozzles on field corn up to 20 inches tall or 6 leaf collars whichever is more restrictive. Broader spectrum. Avoid spraying directly into the whorls of large cornstalks. Do not apply to seed corn.
Soil Residual Tank Mix Partners	Use HALOSULFURON 75WDG HERBICIDE at ⅔ oz. See tank mixture partner label for its rates. Micro-Tech®, EPA Reg. No. 524-344(alachlor) or Bullet®, EPA Reg. No. 524-418 (alachlor, atrazine) or Harness® Xtra, EPA Reg. No. 524-480 (acetochlor, atrazine) or Harness® Xtra 5.6L, EPA Reg. No. 524-485 (acetochlor, atrazine) or Degree Xtra®, EPA Reg. No. 524-511 (alachlor, atrazine). NIS (1 qt./100 gallons of spray) and 28% nitrogen fertilizer (4 gal/100 gallons of spray). Apply as broadcast spray in 15 – 30 gallons of spray/acre to emerged grasses at the 2 leaf stage or less and on corn less than 11 inches tall (5 inches tall for Micro-Tech® and Bullet®). For early post-emergence control of additional small broadleaf, nutsedge and emerged grasses and pre-emergence control or reduced competition of annual broadleaf weeds and grasses as listed on the partner product label. To control emerged lambsquarter less than 4 inches tall, use dicamba or Clarity® Herbicide, EPA Reg. No. 7969-137 (dicamba, diglycolamine salt)
	Use HALOSULFURON 75WDG HERBICIDE at ⅔ oz. and Accent®, EPA Reg. No. 352-560 (nicosulfuron). See tank mixture partner label for its rates. Micro-Tech®, EPA Reg. No. 524-344 (alachlor) or Bullet®, EPA Reg. No. 524-418 (alachlor, atrazine) or Harness® Xtra, EPA Reg. No. 524-480 (acetochlor, atrazine) or Harness® Xtra 5.6L, EPA Reg. No. 524-485 (acetochlor, atrazine) or Degree Xtra® EPA Reg. No. 524-511 (alachlor, atrazine). NIS (1 qt./100 gallons of spray) and 28% nitrogen fertilizer (4 gal/100 gallons of spray). Apply as broadcast spray in 15 – 30 gallons of spray/acre to emerged grasses at the 2 leaf stage or less, foxtail less than 2 inches tall and on corn less than 11 inches tall (5 inches tall for Micro-Tech® and Bullet®). For early post-emergence control of additional small broadleaf and emerged grasses, including foxtail and pre-emergence control or reduced competition of annual broadleaf weeds and grasses as listed on the partner product label. To control emerged lambsquarter less than 4 inches tall, use dicamba or Clarity® EPA Reg. No. 7969-137 [dicamba, diglycolamine salt].
	Use HALOSULFURON 75WDG HERBICIDE plus Accent®, EPA Reg. No. 352-560 (nicosulfuron), Beacon®, EPA Reg. No. 352-560 (nicosulfuron) Option®, EPA Reg. No. 352-560 (nicosulfuron) or Steadfast®, EPA Reg. No. 352-608 (nicosulfuron, rimsulfuron). See tank mixture partner label for its rates. Alachlor, acetochlor, metolachlor and dimethenamid. NIS (1 qt./100 gallons of spray) and 28% nitrogen fertilizer (4 gal/100 gallons of spray). Apply as broadcast spray in 15 – 30 gallons of spray/acre to emerged foxtails and other grasses. For early post-emergence and residual control of emerged foxtails and other grass weeds in seed and field corn. Provides residual control or reduced competition of annual grasses and certain broadleaf weeds as listed on the specific herbicide labels.

	Follow all directions and restrictions on maximum corn height for post applications on this label and the tank mix partner's label. Use the more restrictive guidelines.
Pioneer IR Field Corn Hybrids	Use HALOSULFURON 75WDG HERBICIDE at 1½ - 2 oz. Apply broadcast spray to soil. For residual control of velvetleaf, common cocklebur, common lambsquarter, common ragweed, pigweed, smartweed, sunflower and other difficult to control weeds. Use only on Pioneer IR corn hybrids.
Pre-plant, Pre-emergent.	Use HALOSULFURON 75WDG HERBICIDE plus Accent®, EPA Reg. No. 352-560 (nicosulfuron), Beacon®, EPA Reg. No. 352-560 (nicosulfuron), Option®, EPA Reg. No. 352-560 (nicosulfuron) or Steadfast®, EPA Reg. No. 352-608 (nicosulfuron, rimsulfuron). See tank mixture partner label for its rates. Such as Micro-Tech®, EPA Reg. No. 524-344(alachlor), Bullet®, EPA Reg. No. 524-418 (alachlor, atrazine), Harness®, EPA Reg. No. 524-423 (acetochlor), Harness® Xtra, EPA Reg. No. 524-480 (acetochlor, atrazine), Harness® Xtra 5.6L, EPA Reg. No. 524-485 (acetochlor, atrazine), Degree Xtra® EPA Reg. No. 524-511 (alachlor, atrazine), Lasso®, EPA Reg. No. 524-314 (alachlor), alachlor, acetochlor, metolachlor and dimethenamid. NIS (1 qt/100 gallons of spray) and 28% nitrogen fertilizer (4 gal/100 gallons of spray). Apply as broadcast spray in 15 – 30 gallons of spray/acre. Apply as an early pre-plant surface-applied, pre-plant incorporated or pre-emergence treatment. For effective broadleaf control in tank partner combinations with pre-emergence grass herbicides across all tillage systems. Follow all directions and restrictions on this label and the tank mix partner's label. Use the more restrictive guidelines.

Crop	Rate Oz./Acre	PHI	LIMITATIONS
FRUITING VEGETABLES (Including eggplant, peppers, tomatoes) Crop Group	½ - 1	30	Including eggplant, peppers, tomatoes. Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application. Do not apply more than 2 applications per crop cycle. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per year. Rope-wick or Wiper Application Row middle/furrow application – Apply 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre.
Direct-Seeded and Transplant: Row Middle/Furrow Applications – Avoid contact of this product with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.			
Crop	Rate Oz./Acre	PHI	RESTRICTIONS
GRAIN SORGHUM (Milo)	¾ – 1	30	Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application. Do not make more than 1 application per year. Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre per year. Following application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage. Do not apply this product by rope-wick or wiper applicators.
If used alone, apply at the 2-leaf through lay-by stage of grain sorghum (before the grain head emerges). If grain sorghum is under stress, temporary stature reduction occurs to the crop following application of this product. After application this effect will be evident in 7 – 10 days but under normally growing conditions will quickly recover.			
Sorghum Weed Height Activity Table			
Weed Activity	Control		Suppression
Rate of Product	¾ oz.	1 oz.	¾ oz.

Weed Height	Inches	Inches	Inches
Burcucumber			1 - 3
Cocklebur, common	1 - 9		
Fleabane, Philadelphia	1 - 3		
Kochia ¹	1 - 3		
Lambsquarter, common			1 - 2
Mallow, Venice	1 - 3		
Milkweed, common			3 - 5
Milkweed, honeyvine			1 - 3
Nutsedge: yellow ²	3 - 6	3 - 12	
purple	3 - 6	3 - 12	
Passionflower, maypop	1 - 3		
Pigweed, redroot	1 - 3		
Pokeweed, common	1 - 6		
Ragweed: common	1 - 9		
Giant	1 - 3		
Smartweed, Pennsylvania	1 - 2		
Sunflower, common	1 - 12		
Velvetleaf	1 - 9		

¹ See Pre-emergent and Post-emergent Weed Activity Tables.

² Heavy infestations of nutsedge require sequential applications. To prevent nutsedge from competing with the crop an earlier application is required.

Tank Mixture Partners

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

2,4-D	See tank mixture partner label for its rates. NIS Apply broadcast spray on sorghum 6 to 15 inches tall. If sorghum exceeds 8 inches, use directed spray with drop nozzles and avoid spray on foliage. Broadleaf weeds. Do not treat during the boot, flower or dough stage. Do not make applications when sorghum exceeds 15 inches.
atrazine (various liquid formulations)	See tank mixture partner label for its rates. COC Apply broadcast spray on sorghum up to 12 inches tall. Apply when broadleaf weeds are small (3 inches or less). Post-emergence control of labeled broadleaf weeds. Aids in the burndown and control of many grass weeds (1.5 inches or less) which have escaped pre-emergence herbicide treatments. Atrazine mixtures may result in reduced control (antagonism) of larger broadleaf weeds. Smaller weeds are easier to control.
Buctril® Herbicide EPA Reg. No. 264-437 (bromoxynil octanoate)	See tank mixture partner label for its rates. NIS Apply broadcast spray on sorghum. For post-emergence control of annual broadleaf weeds.
Buctril® Herbicide EPA Reg. No. 264-437 (bromoxynil octanoate) plus atrazine (various formulations)	See tank mixture partner label for its rates. NIS Apply broadcast spray on sorghum. For post-emergence control of annual broadleaf weeds.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
MILLET, PROSO	$\frac{1}{2}$ - $\frac{2}{3}$	See Text	Do not make more than 1 application per 12 month period. Do not apply more than $\frac{1}{4}$ oz. of this product (0.031 lbs. halosulfuron-methyl) per acre per 12 month period. Do not apply this product by rope-wick or wiper applicators. Do not apply within 0 days of harvest for millet forage. Do not apply within 50 days of harvest for millet grain and straw. Do not apply within 37 days of harvest for millet hay.

Post-emergence Broadcast Application

Use Rate: $\frac{1}{2}$ - $\frac{2}{3}$ oz.

Apply from the 2 leaf through layby stage (before grain head emergence).

If the proso millet is under stress, temporary stature reduction may occur to the crop following application of this product.

This effect will be most evident 7 to 10 days after application. The crop will quickly recover under normal growing conditions. Made applications after weed emergence and to actively growing weeds.

Tank Mixture Partners

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use.

Users must follow the most restrictive directions and precautionary language of the products in the mixture.

Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

For broader spectrum broadleaf weed control, use partners including but not limited to 2,4-D and dicamba.

Insecticide partners and fungicide products labeled for uses on millet, proso may be used with this product.

Listed day intervals following an application of this product are:

Crop	Lactating and Non-lactating Animals		
	Pre-Grazing Interval (PGI)	Pre-Harvest Interval (PHI)	Pre-Slaughter Interval (PSI)
millet forage	0	0	0
millet grain	N/A	50	0
millet straw	N/A	50	0
millet hay	N/A	37	0

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
OKRA	$\frac{1}{2}$ - 1	30	Do not make more than 2 applications per 12 month period. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per 12 month period. Do not apply this product by rope-wick or wiper applicators.
For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil.			
Direct-seeded and Transplant: <i>Row Middle/Furrow Application/Shielded Spray</i> – Use this product between rows of direct-seeded or transplanted crop. Avoid contact of this product with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS											
PASTURE, RANGELAND, CRP AND FORAGE GRASSES/HAY Crop Group 17	$\frac{2}{3}$ - $1\frac{1}{3}$	37	Do not make more than 2 applications per 12 month period. Do not apply more than $1\frac{1}{2}$ oz. of this product (0.062 lbs. halosulfuron-methyl) per acre per 12 month period. Do not apply this product by rope-wick or wiper applicators.											
<p>For broadcast spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 10 gallons of water per acre.</p> <p><u>Post-emergence Broadcast Applications to Established Fields</u> Use Rate: $\frac{2}{3}$ - $1\frac{1}{3}$ oz. Make application as soon as possible after removal of hay or before weeds exceed label height restriction. Delay irritation for at least 48 hours after application.</p> <p><u>Post-emergence Spot Treatment to Established Fields</u> Use Rate: $\frac{2}{3}$ - $\frac{3}{4}$ oz. Use at rates equivalent to broadcast field rates and not exceeding the maximum application rate of $\frac{3}{4}$ oz./acre. Use with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil.</p> <p><u>Post-emergence Broadcast followed by Post-emergence Spot Treat</u> Use Rate: $\frac{2}{3}$ - $\frac{3}{4}$ oz. To maximize control of nutsedge, it may be necessary to use a second post-emergence spot application to those areas where the nutsedge has emerged or re- grown. In this case, use a spot treatment method treating only those areas of emerged nutsedge. Use at rates equivalent to broadcast field rates and not exceeding the maximum application rate of $\frac{3}{4}$ oz./acre. Apply with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds.</p> <p><u>Tank Mixture Partners</u> It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.</p> <p>For broader spectrum broadleaf weed control, use partners including but not limited to 2,4-D, dicamba and, Grazon® P+D, EPA Reg. No. 62719-182 (picloram, potassium salt).</p> <p>Insecticide partners, including Confirm®, 2F EPA Reg. No. 8033-111 (tebufenozide) and fungicide products labeled for uses on Pasture, Rangeland, CRP and Forage Grasses/Hay section may be used with this product.</p> <p>Listed day intervals following an application of this product are:</p> <table border="1"> <thead> <tr> <th rowspan="2">Crop</th><th colspan="3">Lactating and Non-lactating Animals</th></tr> <tr> <th>Pre-Grazing Interval (PGI)</th><th>Pre-Harvest Interval (PHI)</th><th>Pre-Slaughter Interval (PSI)</th></tr> </thead> <tbody> <tr> <td>Pasture, Rangeland, CRP and Forage Grasses/Hay</td><td>0</td><td>37</td><td>0</td></tr> </tbody> </table>				Crop	Lactating and Non-lactating Animals			Pre-Grazing Interval (PGI)	Pre-Harvest Interval (PHI)	Pre-Slaughter Interval (PSI)	Pasture, Rangeland, CRP and Forage Grasses/Hay	0	37	0
Crop	Lactating and Non-lactating Animals													
	Pre-Grazing Interval (PGI)	Pre-Harvest Interval (PHI)	Pre-Slaughter Interval (PSI)											
Pasture, Rangeland, CRP and Forage Grasses/Hay	0	37	0											

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
POME FRUIT (East of the Rockies) (Apple; azarole; crab apple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these.) Crop Group 11-10	½ - 1	14	<p>Do not use this product if orchard temperatures are greater than 85° F at time of application.</p> <p>Do not make applications sooner than 45 days apart.</p> <p>Do not overlap spray swath to increase the application rate into the treated area.</p> <p>Do not use on trees in established orchards that are less than 1 year old or to plants under stress.</p> <p>Do not apply to nursery stock</p> <p>Do not apply to areas where water is known to pond for periods of time following rainfall.</p> <p>Do not contact foliage with this product. Uptake via contacted foliage will result in plant injury.</p> <p>Do not make more than 2 applications per 12 month period.</p> <p>Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per 12 month period.</p> <p>Do not apply this product by rope-wick or wiper applicators.</p>
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Pre-emergence and Post-emergence Directed Applications for listed broadleaf and annual grass weeds. Use Rate: ½ - 1 oz. Use this product as a single spray or sequential directed spray to orchard floor on either side of row. If small weeds are present, tank mix with a post-emergence broad-spectrum type herbicide to maximize and enhance the spectrum of broadleaf and grass control. If ground cover prevents contact with the soil, reduced or no residual herbicidal activity will result.</p> <p>Post-emergence directed application for control of nutsedge. Use Rate: ½ - 1 oz. Use this product as a single directed spray application when nutsedge is fully emerged or as two sequential directed spray applications. Make the first directed spray application to the initial nutsedge flush when it has reached the 3-5 leaf stage. If a second treatment is needed, it may be applied later in the season directed to secondary nutsedge emergence. For best results, use a minimum rate of ¾ oz. and when the nutsedge plants are in the 3-5 leaf stage.</p> <p>PRECAUTIONS: For best results in post-emergence applications, use a NIS spray additive. Contact of this product with the tree foliage or fruit by spray or drift may result in significant injury. Use of a shielded boom is recommended. This product may not control ALS-resistant weeds.</p>			
Crop	Rate Oz./Acre	PHI	RESTRICTIONS
POME FRUIT (West of the Rockies) (Apple; azarole; crab apple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these.) Crop Group 11-10	¾ - 2	14	<p>Do not use this product if orchard temperatures are greater than 85° F at time of application.</p> <p>Do not make applications sooner than 45 days apart.</p> <p>Do not overlap spray swath to increase the application rate into the treated area.</p> <p>Do not use on trees in established orchards that are less than 1 year old or to plants under stress.</p> <p>Do not apply to nursery stock</p> <p>Do not apply to areas where water is known to pond for periods of time following rainfall.</p> <p>Do not contact foliage with this product. Uptake via contacted foliage will result in plant injury.</p> <p>Do not make more than 2 applications per 12 month period.</p> <p>Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per 12 month period.</p>

			Do not apply this product by rope-wick or wiper applicators.
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Pre-emergence and Post-emergence Directed Applications for listed broadleaf and annual grass weeds. Use Rate: $\frac{3}{4}$ - 2 oz. Use this product as a single spray or sequential directed spray to orchard floor on either side of row. If small weeds are present, tank mix with a post-emergence broad-spectrum type herbicide to maximize and enhance the spectrum of broadleaf and grass control. If ground cover prevents contact with the soil, reduced or no residual herbicidal activity will result.</p> <p>Post-emergence broadcast application for control of nutsedge. Use Rate: $\frac{3}{4}$ - 2 oz. Use this product as a single broadcast spray application to orchard floor on either side of row when nutsedge is fully emerged (early to midsummer) or as two sequential broadcast spray applications. Make the first broadcast spray application to the initial nutsedge flush when it has reached the 3-5 leaf stage. If a second treatment is needed, it may be applied later in the season to secondary nutsedge emergence. For best results, make applications when nutsedge is less than 12 inches in height.</p> <p>PRECAUTIONS: For best results in post-emergence applications, use a NIS or penetrating spray additive. Contact of this product with the tree foliage or fruit by spray or drift, may result in significant injury. This product may not control ALS-resistant weeds.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
PUMPKINS AND WINTER SQUASH	$\frac{1}{2}$ - 1	30	<p>Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application. Do not make more than 2 applications per year. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl), per acre per year (includes applications to the crop and to row middles). Where possible, apply $\frac{1}{2}$ to $\frac{3}{4}$ inch of sprinkler irrigation to settle the soil after planting and prior to application.</p> <p>Rope-wick or Wiper Application Row middle/furrow application – Apply 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre.</p>
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Direct-seeded: Use Rate: $\frac{1}{2}$ - $\frac{3}{4}$ oz. <i>Pre-emergence</i> – Use this product after planting, but before cracking. For lighter textured soils with low organic matter, use the lower rate.</p> <p><i>Post-emergence</i> – Apply after the crop has reached at least 2-5 true leaf stage, preferably 4-5 true leaves, but before first female flowers appear. For lighter textured soils with low organic matter, use the lower rate.</p> <p>Transplanted: Use Rate: $\frac{1}{2}$ - $\frac{3}{4}$ oz. <i>Pre-transplant</i> – For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-transplant application under the plastic mulch. Crop may be transplanted into this treated area no sooner than 7 days after application unless local conditions demonstrate safety at an earlier interval. For lighter textured soils with low organic matter, use the lower rate. This product treated in soil in transplant hole may result in crop injury. During the transplant process, take care to limit movement of soil.</p>			

Post-transplant – Apply this product to transplants that are established and actively growing. Application should not be made until plants are actively growing and in the 3-5 true leaf stage or no sooner than 14 days after transplanting unless local conditions demonstrate safety at an earlier interval, but before first female flowers appear. Apply this product as an over-the-top application, a directed spray application, or with crop shields to minimize contact of the herbicide with the crop.

For Processing Only - Direct-seeded:

Use Rate: ½-1 oz.

Pre-emergence – Use this product after planting, but before cracking. For lighter textured soils with low organic matter, use the lower rate.

Post-emergence – Apply after the crop has reached at least 2-5 true leaf stage, but before first female flowers appear. For lighter textured soils with low organic matter, use the lower rate.

Direct-seeded and Transplant:

Use Rate: ½-1 oz.

Row Middle/Furrow Applications – For the treatment of nutsedge and labeled broadleaf weeds, use this product between rows of direct-seeded or transplanted crop. Avoid contact of this product with the planted crop. If plastic is used on the planted rows, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
RICE	⅔ – 1 ⅓	48*	<p>Do not apply more than 1½ oz. of this product (0.062 lbs. halosulfuron-methyl) per application.</p> <p>Do not make more than 3 applications (including pre-plant and at-planting applications) per year.</p> <p>Do not apply more than 1½ oz. of this product (0.062 lbs. halosulfuron-methyl) per acre per use year.</p> <p>After application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.</p> <p>*Do not apply within 69 days of harvest in California.</p> <p>For Direct Dry Applications by air:</p> <p>Do not apply to dry rice fields.</p> <p>Apply aerial applications at a maximum of no greater than ½ the wing span.</p> <p>Do not use a swath width greater than 120 feet.</p> <p>Do not mix this product with any other additives except as directed by this label.</p> <p>Do not apply within 360 feet of sensitive crops.</p> <p>Do not apply when wind speed is less than 3 mph or exceeds 15 mph.</p> <p>Do not apply this product by rope-wick or wiper applicators.</p>

Rice Weed Height Activity Table

Weed Activity	Control		Suppression	
Rate of Product	⅔ oz.	1 - 1⅓ oz.	⅔ oz.	1 - 1⅓ oz.
Weed Height	Inches	Inches	Inches	Inches
Burcucumber			1 - 3	4 - 12
California Arrowhead		Yes		
Cocklebur, common	1 - 9	9 – 14		
Dayflower	1 - 2	3 – 4		
Eclipta	1 – 4	4 – 8		
Flatsedge rice	1 - 9	9 – 12		
Fleabane, Philadelphia	1 - 3			
Jointvetch	1 – 2	3 – 4		
Kochia ¹	1 - 3			3 - 6
Lambsquarter, common			1 - 2	
Mallow, Venice	1 - 3	4 - 12		
Milkweed, common			3 - 5	6 - 12
Milkweed, honeyvine		1 - 6	1 - 3	

Morningglory				1 - 3
Mustard, wild		4 - 6		
Nutsedge: yellow ²	1 - 6	6 - 12		
purple	1 - 6	6 - 12		
Passionflower, maypop	1 - 3			
Pigweed, redroot	1 - 3	4 - 6		
Pokeweed, common	1 - 6			
Radish, wild		4 - 6		
Ragweed: common	1 - 9	9 - 12		
Giant	1 - 3	4 - 6		
Redstem	1 - 3	Yes		
Ricefield Bulrush		Yes		
Sesbania Hemp	1 - 3	3 - 6		
Sida, Prickly	1 - 2	3 - 4		
Smallflower Umbrellaplant		Yes		
Smartweed, Pennsylvania	1 - 2			
Sunflower, common	1 - 12	12 - 15		
Velvetleaf ³	1 - 9	9 - 12		

¹ See the Post-emergent Weed Activity Table.

² Heavy infestations of nutsedge require sequential applications. To prevent nutsedge from competing with the crop an earlier application is required.

³ For large velvetleaf and pigweed control, apply with liquid nitrogen fertilizer (2 – 4 qts./acre).

Pre-plant, at-plant, post-emergent and prior to emergence of rice through permanent flood:

Use ½ - 1 ½ oz. of this product per acre per year.

Apply foliar applications of this product at the 3 – 5 leaf stage of rice when weeds have 2 – 4 leaves.

For foliar applications, use nonionic surfactant at rate of 0.25 – 0.5% in the spray mixture.

For aerial foliar applications, use a minimum of 3 – 15 gallons of water per acre.

For ground foliar applications, use a minimum of 10 gallon of water per acre.

After mixing, apply spray suspensions the same day for best results.

PRECAUTIONS:

Best control of emerged weeds with foliar applications occurs when 70% - 80% of the weed foliage is exposed.

For best control of submerged weeds, apply when weeds have 2 leaves or less.

Check spray drift management section of this label.

Following the foliar applications of this product, do not reintroduce water into rice fields or checks for at least 24 hours.

To improve the spectrum of weed control, tank mix this product with Shark® EW, EPA Reg. No. 279-3242 (carfentrazone-ethyl) or Shark H2O, EPA Reg. No. 279-3194 (carfentrazone-ethyl).

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use.

Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

Sequential Applications:

To improve the spectrum of weed control, apply this product sequentially with Bolero®8 EC, EPA Reg. No. 59639-79 (thiobencarb), Clincher®, EPA Reg. No. 62719-357 (cyhalofop-butyl), Ordam®, EPA Reg. No. 10182-204 (molinate), Regiment®, EPA Reg. No. 59639-105 (bispaybac-sodium) or Shark® EW, EPA Reg. No. 279-3242 (carfentrazone-ethyl) or Shark H2O, EPA Reg. No. 279-3194 (carfentrazone-ethyl).

Direct Dry Application:

Apply this product post flood as a dry broadcast application at a rate of 1 – 1 ½ oz. per acre per year. When weeds have 2 leaves or less, apply the dry broadcast treatment of this product at 1 – 2 leaf stage of rice.

Water levels in rice fields and checks should remain static (3 – 6 inch depth) after dry broadcast applications of this product.

Do not reintroduce water into rice fields or checks for at least 5 days after dry broadcast treatments.

Rice fields and checks may be irrigated to maintain water level, but this may reduce weed control.

Co-application with Shark H2O, EPA Reg. No. 279-3194 (carfentrazone-ethyl) is allowed.

Tank Mixture Partners

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

Before using tank mixture partners, conduct a compatibility test according to the outline in the Tank Mixture section of the label.

For additional pre-emergence weed control in preemergence & preplant applications, tank partner including but not limited to Bolero®8 EC, EPA Reg. No. 59639-79 (thiobencarb), Command® 3ME, EPA Reg. No. 279-3158 (clomazone), glyphosate, pendimethalin or quinclorac may be used.

For additional post-emergence broadleaf weed control, tank partners including but not limited to Grandstand®, EPA Reg. No. 62719-215 (triclopyr, triethylamine salt), propanil, Aim®, EPA Reg. No. 279-3194 (carfentrazone-ethyl), Facet® EPA Reg. No. 7969-313 (quinclorac), Basagran®, EPA Reg. No. 7969-45 (sodium bentazon), Londax®, EPA Reg. No. 70506-147 (bensulfuron-methyl), Grasp®, EPA Reg. No. 62719-500 (penoxsulam), Regiment®, EPA Reg. No. 59639-105 (bispyrbac-sodium), NewPath®, EPA Reg. No. 241-412 (imazethapyr), Beyond®, EPA Reg. No. 241-441 (imazamox) and 2,4-D may be used.

For additional post-emergence grass control, tank partners including but not limited to NewPath®, EPA Reg. No. 241-412 (imazethapyr), Beyond®, EPA Reg. No. 241-441 (imazamox), propanil, Facet®, EPA Reg. No. 7969-313 (quinclorac), Grasp®, EPA Reg. No. 62719-500 (penoxsulam), and Regiments®, EPA Reg. No. 59639-105 (bispyrbac-sodium), may be used.

Glyphosate (various formulations)	Use HALOSULFURON 75WDG HERBICIDE at $\frac{2}{3}$ oz. See Glyphosate label for its rates. NIS Broadcast spray. For pre-plant or at-planting burndown of emerged annual grasses, broadleaf weeds and nutsedge. If applied as a pre-plant burn down treatment, consult the Crop Rotational Guidelines of this product and the Glyphosate label.
Stam® M4 EPA Reg. No. 71085-36 (propanil) and propanil (various liquid formulations)	Use HALOSULFURON 75WDG HERBICIDE at $\frac{2}{3}$ – 1 $\frac{1}{2}$ oz. See propanil labels for its rates. Broader spectrum weed control. If applied as a pre-plant burn down treatment, consult the Crop Rotational Guidelines.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
RHUBARB	$\frac{1}{2}$ - 1	60	Do not make more than 2 applications per 12 month period. Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre per 12 month period. Do not apply this product by rope-wick or wiper applicators.
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Broadcast Application: Use Rate: $\frac{1}{2}$ - 1 oz. Apply this product as a single broadcast application to dormant rhubarb.</p> <p>PRECAUTIONS: For best results in post-emergence applications, use a NIS spray additive. This product may not control ALS-resistant weeds Make the timing of the application as late as possible, or just prior to the breaking of rhubarb dormancy. This product may cause significant crop stunting. Test with the lower rate to determine potential sensitivity to its use along with speed and degree of recovery.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
SMALL FRUIT VINE CLIMBING (Except FUZZY KIWIFRUIT) (East of the Rockies) (Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these.) Crop Subgroup 13-07F	½ - 1	14	<p>Do not overlap spray swath which will increase the application rate in the treated area.</p> <p>Do not use on vines in established vineyards that are less than 1 year old or to plants under stress.</p> <p>Do not apply to areas where water is known to pond for periods of time following rainfall.</p> <p>Do not contact foliage, especially grape vines, with this product.</p> <p>Uptake via contacted foliage will result in plant injury.</p> <p>Do not make applications sooner than 45 days apart.</p> <p>Do not apply to nursery stock</p> <p>Do not make more than 2 applications per 12 month period.</p> <p>Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per 12 month period.</p> <p>Do not apply this product by rope-wick or wiper applicators.</p>
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Pre-emergence and Post-emergence Directed Applications for control of listed broadleaf and annual grass weeds. Use Rate: ½ - 1 oz. Use this product as a single spray or sequential directed spray to ground on either side of row. If small weeds are present, tank mix with a post-emergence broad-spectrum type herbicide to maximize and enhance the spectrum of broadleaf and grass control. If ground cover prevents contact with the soil, reduced or no residual herbicidal activity will result.</p> <p>Post-emergence directed application for control of nutsedge. Use Rate: ½ - 1 oz. Use this product as a single directed spray application when nutsedge is fully emerged or as two sequential directed spray applications. Make the first directed spray application to the initial nutsedge flush when it has reached the 3- leaf stage. If a second treatment is needed, it may be applied later in the season directed to secondary nutsedge emergence. For best results, use a minimum rate of ¾ oz. and when the nutsedge plants are in the 3-5 leaf stage.</p> <p>PRECAUTIONS: For best results in post-emergence applications, use a NIS spray additive. Contact of this product with the grape vines will result in leaf chlorosis and distortion with possible shortening of shoot internodes. Use of a shielded boom is suggested. This product may not control ALS-resistant weeds.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
SOYBEANS, Soybean Seed (Except CA)	⅔ - 1 ⅓	88	<p>Do not apply more than 1⅓ oz. of this product (0.062 lbs. halosulfuron-methyl) per application.</p> <p>Do not make more than 1 application per year.</p> <p>Do not apply more than 1⅓ oz. of this product (0.062 lbs. halosulfuron-methyl) per acre per year.</p> <p>Do not apply to frozen ground.</p> <p>After application to foliage, allow 30 days before grazing domestic livestock or harvest forage, silage and hay.</p> <p>Do not apply this product if plans include planting Adzuki beans since unacceptable crop injury may occur.</p> <p>Do not use post-emergence applications of this product to straight Roundup Ready or conventional soybean varieties as severe crop injury will occur.</p> <p>Do not use more than ⅔ oz. of this product (0.031 lbs. halosulfuron-methyl), in post-emergence application to sulfonyl-urea tolerant soybeans (STS).</p>

Pre-plant Burndown – Fall Application:

Use Rate: $\frac{3}{4}$ – 1 $\frac{1}{2}$ oz.

For control or suppression of listed broadleaf winter annual weeds prior to planting soybeans in the following Spring, apply this product as a fall burndown herbicide and/or preventative application.

If listed broadleaf weed are visible, use a high quality crop oil concentrates (1 – 2% vol/vol) and granular AMS (2 – 4 lbs./A) or UAN (1 -2% vol/vol) to the spray suspension to improve performance.

Apply this product from anytime from after harvest up until the ground freezes.
Apply this product by ground or air.

In research trials, no instances of crop injury from Fall applied applications have occurred but not all soybean varieties have been tested for crop tolerance to halosulfuron-methyl. For the latest halosulfuron-methyl tolerance information, consult the local seed agronomists and seed supplier.

For broadleaf winter annual weed activity, this product must contact the emerged weeds and reach the soil surface.
In reduce tillage systems to maintain the active ingredient in the top layer of soil where weed seeds germinate, apply this product after fall chisel, disking etc.

Tank Mixture Partners

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

2,4-D Amine or LV ester (various formulations)	See tank mixture partner label for its rates. For broader spectrum control in pre-plant burndown of emerged annual broadleaf weeds or under heavy weed infestation.
Glyphosate (various formulations)	See tank mixture partner label for product rates. For control in pre-plant burn down of emerged grass weeds.

Pre-emergence or Pre-plant Spring Application to Soybean Varieties Tolerant to Sulfonyl-urea Herbicides (STS) Only:

Use Rate: $\frac{3}{4}$ oz.

For contact and residual control or suppression of listed broadleaf winter and early germinating summer annual weeds, apply this product from 21 days before planting until prior to emergence (i.e. cracking).

For best performance, apply this product to actively growing weeds free from environmental stress.

If listed broadleaf weed are visible, use a high quality crop oil concentrates (1% vol/vol) and granular AMS (2 – 4 lbs./A) or UAN (1 -2% vol/vol) to the spray suspension to improve performance.

For use on any soybean varieties tolerant to sulfonyl-urea herbicides (STS) unless prohibited by the seed supplier.

In research trials, no instances of crop injury from Spring pre-plant or pre-emergence applied applications have occurred but not all sulfonyl-urea tolerant soybean (STS) varieties have been tested for crop tolerant to halosulfuron-methyl. For the latest halosulfuron-methyl tolerance information, consult the local seed agronomists and seed supplier.

In reduce tillage systems to maintain the active ingredient in the top layer of soil where weed seeds germinate, apply this product after all tillage operations.

Tank Mixture Partners

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.	
2,4-D LV ester (various formulations)	See tank mixture partner label for its rates. For control in early germinating summer weeds.
Glyphosate (various formulations)	See tank mixture partner label for product rates. For broader spectrum control in pre-plant burndown of emerged annual broadleaf weeds or grasses.

Post-emergence Application to Soybean Varieties Tolerant to Sulfonyl-urea Herbicides (STS) Only: Use Rate: $\frac{2}{3}$ oz. For contact and residual control of listed broadleaf weeds and nutsedge, apply this product post-emergent from V1 through R2 stages of sulfonyl-urea tolerant soybean (STS) varieties only. If the tolerant soybean variety is also stacked with glyphosate or glufosinate tolerant trait, then glyphosate or glufosinate respectively may be used as a tank mixture partner. For best performance, apply this product to actively growing weeds free from environmental stress. For use on any soybean varieties tolerant to sulfonyl-urea herbicides (STS) unless prohibited by the seed supplier. Always use a NIS (1 to 2 qts./100 gallons of spray) or high quality crop oil concentrates (1 % vol/vol) and granular AMS (2 – 4 lbs./A) or UAN (1 -2% vol/vol) to the spray suspension to improve performance. Some phytotoxicity from post-emergent applications may occur on susceptible sulfonyl-urea tolerant soybeans (STS) varieties. These symptoms may include stunting (seen as reduction in leaf size or internodal length), yellowing of leaves and or red veins and necrosis of leaves and petioles. Sulfonyl-urea tolerant soybeans (STS) that have exhibited these symptoms tend to recover after the product is metabolized by the plant. Soybean injury is most noticeable when the plants are under environmental stress conditions such as hot, humid conditions, or wide fluctuations in climatic conditions, drought, etc. For the latest halosulfuron-methyl tolerance information, consult the local seed agronomists and seed supplier.	
Tank Mixture Partners It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.	
Glyphosate (various formulations)	See tank mixture partner label for product rates. For broader spectrum control of emerged annual broadleaf and grass weeds or under heavy weed infestation.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
SUCCULENT SNAP BEANS (Including lima beans)	$\frac{1}{2}$ - 1	30	Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application. Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl), per acre per crop cycle, not to exceed 2 oz. (0.094 lbs. halosulfuron-methyl) per acre per 12-month period (includes applications to the crop and to row middles/furrows). Application of this product may cause significant, temporary stunting and delay maturity of snap beans resulting in delayed harvest. This product is available to the end-user/grower solely to the extent that the benefit and utility, in the sole opinion of the end-user/grower, outweigh the extent of potential injury associated with the use of this product. To the extent consistent

			with applicable law, the risk of crop damage, is at the end-user/grower's risk. Do not apply this product by rope-wick or wiper applicators.
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Direct-seeded: Use Rate: ½-1 oz. <i>Pre-emergence</i> – Use this product after planting, but before cracking. For lighter textured soils with low organic matter, use the lower rate.</p> <p><i>Row Middle/Furrow Applications</i> – Apply this product for the control of nutsedge and listed broadleaf weeds. Avoid contact of this product with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.</p> <p>Direct-seeded: Use Rate: ½-¾ oz. <i>Post-emergence</i> – Apply after the crop has reached the 2-4 trifoliate leaf stage, but before flowering. For lighter textured soils with low organic matter, use the lower rate. Use directed sprays to limit crop injury.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
SUCCULENT SHELLED PEA AND BEAN (Any succulent shelled cultivar of bean (<i>Phaseolus</i>) including lima bean, green; broad bean, succulent; (<i>vigna</i>) including blackeyed pea, cowpea, southern pea,) Crop Subgroup 6B	½ - 1	30	<p>For Pre-emergence application, do not make more than 1 application per 12 month period.</p> <p>For Post-emergence applications, do not make more than 2 applications per 12 month period.</p> <p>Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre per 12 month period.</p> <p>Do not apply this product on Adzuki beans, English peas and garden peas.</p> <p>Do not feed to livestock.</p> <p>Do not apply this product by rope-wick or wiper applicators.</p>
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p><u>Pre-emergence Applications for control of listed broadleaf weeds.</u> Use Rate: ½ oz. Use this product as a single broadcast spray after planting but before crop emergence.</p> <p>PRECAUTIONS: This product may not control ALS-resistant weeds. Use of this product may cause significant, temporary stunting and delay maturity of peas resulting in delayed harvest. This product is available to the end-user/grower solely to the extent that the benefit and utility, in the sole opinion of the end-user/grower, outweigh the extent of potential injury associated with the use of this product.</p> <p><u>Post-emergence directed application for control of nutsedge.</u> Use Rate: ½ - 1 oz. Use this product as a single directed spray application when plants have 2 to 4 trifoliate leaves and before flowering. Make one broadcast application. To limit crop injury, directed sprays are suggested.</p> <p>Not all varieties have been tested for tolerance. Under adverse growing conditions (dry or excessive moisture, cool weather, etc.), maturity of the treated crop may be delayed which can influence harvest date, yield, and quality. For untested varieties, a small area of the field should be sprayed to determine potential sensitivity to its use.</p> <p>PRECAUTIONS: For best results in post-emergence applications, use a NIS spray additive. Contact of this product with the grape vines will result in leaf chlorosis and distortion with possible shortening of shoot internodes. This product may not control ALS-resistant weeds.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
SUGARCANE	$\frac{2}{3}$ – 1 $\frac{1}{2}$	30	Do not apply more than 1 $\frac{1}{4}$ oz. of this product (0.062 lbs. halosulfuron-methyl) per application. Do not make more than 3 applications (including pre-plant applications) per year. Do not apply more than 2 $\frac{2}{3}$ oz. of this product (0.125 lbs. halosulfuron-methyl) per acre per year. After application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage. Do not apply this product by rope-wick or wiper applicators.

If used alone, apply this product prior to planting, prior to emergence or after the emergence of the sugarcane and until row closure. Use mechanical cultivation to control weed species not on this label. If mechanical cultivation is used, apply a sequential treatment to control weeds in areas of disturbed soil.

Sugarcane Weed Height Activity Table

Weed Activity	Control		Suppression	
Rate of Product	$\frac{2}{3}$ oz.	1 - 1 $\frac{1}{2}$ oz.	$\frac{2}{3}$ oz.	1 - 1 $\frac{1}{2}$ oz.
Weed Height	Inches	Inches	Inches	Inches
Burcucumber			1 - 3	4 - 12
Cocklebur, common	1 - 9	9 - 14		
Fleabane, Philadelphia	1 - 3			
Kochia ¹	1 - 3			3 - 6
Lambsquarter, common			1 - 2	
Mallow, Venice	1 - 3	4 - 12		
Milkweed, common			3 - 5	6 - 12
Milkweed, honeyvine		1 - 6	1 - 3	
Morningglory				1 - 3
Mustard, wild		4 - 6		
Nutsedge: yellow ²	3 - 6	3 - 12		
purple	3 - 6	3 - 12		
Passionflower, maypop	1 - 3			
Pigweed, redroot ³	1 - 3	4 - 6		
Pokeweed, common	1 - 6			
Radish, wild		4 - 6		
Ragweed: common	1 - 9	9 - 12		
Giant	1 - 3	4 - 6		
Smartweed, Pennsylvania	1 - 2			
Sunflower, common	1 - 12	12 - 15		
Velvetleaf ³	1 - 9	9 - 12		

¹ See Pre-emergent and Post-emergent Weed Activity Tables.

² Heavy infestations of nutsedge require sequential applications. To prevent nutsedge from competing with the crop an earlier application is required.

³ For large velvetleaf and pigweed control, apply with liquid nitrogen fertilizer (2 – 4 qts. /acre) plus NIS or COC.

Tank Mixture Partners

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

2,4-D amine	See tank mixture partner label for its rates. NIS Apply to sugarcane before crop emergence or post-emergence until 6 weeks before harvest. Broadleaf weeds. Do not make more than 3 applications per year.
asulam, sodium salt (various	See tank mixture partner label for its rates. NIS or COC

			Do not apply this product to sweet corn or popcorn previously treated with soil applied organophosphate insecticides. Do not apply an organophosphate insecticide within 7 days before or 3 days after any application of this product. Do not apply this product to sweet corn or popcorn unless the seed company, processor or State Agricultural Extension service has tested this product on the particular hybrid or variety and specifically approves and supports the use. Do not apply this product if the sweet corn and popcorn is under severe stress due to drought, water-saturated soils, low fertility (especially low nitrogen levels) or other poor growing conditions. Do not apply this product by rope-wick or wiper applicators.
Spray Applications If used alone, apply a broadcast spray over-the-top or with drop nozzles from the spike through lay-by stage of corn. Use mechanical cultivation to control weeds species not on this label. Apply a sequential treatment, if necessary, only with drop nozzles semi-directed or directed to avoid application into the corn plant whorl.			
PRECAUTIONS: Not all sweet corn and popcorn hybrids or varieties have been tested for sensitivity to this product. To the extent consistent with applicable law, the user assumes all responsibility for such use. After application, avoid cultivation for at least 7 days.			
Sweet Corn and Popcorn Weed Height Activity Table			
Weed Activity	Control	Suppression	
Rate of Product	¾ oz.	¾ oz.	
Weed Height	Inches	Inches	
Burcucumber		1 - 3	
Cocklebur, common	1 - 9		
Fleabane, Philadelphia	1 - 3		
Kochia ¹	1 - 3		
Lambsquarter, common		1 - 2	
Mallow, Venice	1 - 3		
Milkweed, common		3 - 5	
Milkweed, honeyvine		1 - 3	
Nutsedge: yellow ²	3 - 6		
purple	3 - 6		
Passionflower, maypop	1 - 3		
Pigweed, redroot	1 - 3		
Pokeweed, common	1 - 6		
Ragweed: common	1 - 9		
Giant	1 - 3		
Smartweed, Pennsylvania	1 - 2		
Sunflower, common	1 - 12		
Velvetleaf	1 - 9		
¹ See Pre-emergent and Post-emergent Weed Activity Tables. ² Heavy infestations of nutsedge require sequential applications. To prevent nutsedge from competing with the crop an earlier application is required.			
Crop	Rate Oz./Acre	PHI	RESTRICTIONS
TOMATOES	½ - 1	30	Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application. Do not make more than 2 applications per crop cycle. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per year, (includes applications to the crop and to row middles/furrows).

	1		Rope-wick or Wiper Application Row middle/furrow application – Apply 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre.
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 20 gallons of water per acre.</p> <p>Direct-seeded: <i>Post-emergence</i> – Apply this product over-the-top once tomatoes have reached the 4-leaf stage through first bloom. After bloom, applications must be made as a directed spray or with crop shields to minimize contact of this product with the crop.</p> <p>Transplanted: <i>Pre-transplant on Bareground</i> – Apply this product as a pre-transplant application to bareground for the suppression of nutsedge and control of listed broadleaf weeds. Seven (7) days after the application, transplant tomatoes into this treated area unless local conditions demonstrate safety at an earlier interval. For lighter textured soils with low organic matter, use the lower rate. This product treated in soil in transplant hole may result in crop injury. During the transplant process, take care to limit movement of soil.</p> <p><i>Pre-transplant under Plastic Mulch Applications</i> – For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-plant application under the plastic mulch. After final bed shaping and just prior to the installation of the plastic mulch, apply this product. Seven (7) days after the application and the installation of the plastic mulch, transplant the tomatoes into this treated area unless local conditions demonstrate safety at an earlier interval. Soil treated with this product in the transplant hole may result in crop injury. During the transplant process, take care to limit movement of soil.</p> <p><i>Post-transplant</i> – Apply this product to tomato transplants that are established and actively growing. Apply to tomato transplants a minimum of 14 days after transplanting unless local conditions demonstrate safety at an earlier interval but before first bloom. Following bloom, apply this product only as a directed spray or with crop shields to minimize contact of this product with the crop.</p> <p>Direct-seeded and Transplant: <i>Pre-transplant followed by post-emergence for nutsedge control</i>– To maximize the control of nutsedge, use a sequential post-emergence application to those areas where the nutsedge has broken through the plastic mulch. For these situations, use a spot treatment method treating only those areas of emerged nutsedge. Do not exceed ¾ oz. product per treated acre in these areas. Use a water volume that will allow for good coverage of the plants. Soil treated with this product in the transplant hole may result in crop injury. During the transplant process, take care to limit movement of soil.</p> <p><i>Post-emergence Sequential Treatments for Nutsedge Control</i> - To maximize the control of nutsedge, a second sequential post-emergent spray is applied to the areas where nutsedge has re-grown or emerged. In this case, use a spot treatment application for localized control of emerged nutsedge. Allow a minimum of 21 days between applications. Do not exceed 1 oz. product per treated acre in these areas.</p> <p><i>Row Middle/Furrow Applications</i> – For the treatment of nutsedge and labeled broadleaf weeds, use this product between rows of direct-seeded or transplanted crop. Avoid contact of this product with the planted crop. If plastic is used on the planted rows, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
TREE NUTS (almonds, beechnuts, Brazil nuts, butternuts, cashews, chestnuts, chinquapins, filberts, hickory nuts, macadamia nuts, pecans, pistachios, walnuts {black and English})	¾ -1½	1	Do not apply more than 1½ oz. of this product (0.062 lbs. halosulfuron-methyl) per application. Do not make more than 2 applications per acre per year. Do not apply more than 2¾ oz. of this product (0.125 lbs. halosulfuron-methyl) per acre per year. On coarse textured soils classified as sand, loamy sand, and sandy loam with less than 18% clay and more than 65% sand, or on soils with less than 1% organic matter, apply up to 2 applications of this product with a total of all

			<p>applications not to exceed 2 oz. (0.094 lbs. halosulfuron-methyl) of this product per acre per year.</p> <p>Do not apply to gravelly soils.</p> <p>Do not over apply, since excessive application rates can result in severe tree injury or death.</p> <p>Do not use in controlled droplet application, irrigation, or chemigation equipment due to variations in the actual application rate.</p> <p>Do not apply this product by rope-wick or wiper applicators.</p>
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. Use this product as a directed spray to the soil in established tree nut crops. Established tree nut crops are defined as those that have been transplanted into their final growing location for a period of at least 12-months, and where the soil has firmly settled around the roots from packing and rainfall or irrigation.</p> <p>Specified rates are based on broadcast treatment. For band applications, adjust the rate and spray volume proportionally to the actual treated area. For all applications, adjust the rate of this product to account for high volume output nozzles, such as off-center nozzles, and overlaps in the spray pattern.</p> <p>Use a maximum of 1 oz. (0.047 lbs. halosulfuron-methyl) of this herbicide per acre on coarse textured soils classified as sand, loamy sand, and sandy loam with less than 18% clay and more than 65% sand, or on soils with less than 1% organic matter.</p> <p>Use mechanical cultivation or mowing to control weed species not listed on this label. If the soil is disturbed, use a sequential spot treatment for continued control.</p> <p>PRECAUTIONS:</p> <p>Avoid contact of the spray containing this product with trunk, stems, roots, or foliage of tree nut crops, as severe injury or death may result.</p> <p>For the best results, apply this product in the spring when nutsedge is not drought stressed and maximize the interval between application and subsequent irrigation. If this product is applied to trees that have been weakened by or recovering from stress caused by, but not limited to, excessive fertilizer or soil salts, disease, nematodes, frost, wind injury, drought, flooding, previously applied pesticides, insects, winter injury, soil pan of any type, nutrient deficiency, or mechanical damage, severe injury or death may result. Application of this product to weakened or stressed trees as described, especially in soils with less than 1% organic matter, significantly increases the probability of severe injury or death. To the extent consistent with applicable law, all such risks shall be assumed by the user.</p> <p>Tank Mixture Partner:</p> <p>It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.</p> <p>Use this product at labeled rate with glyphosate agricultural herbicides. This will aid in the burndown and control of emerged annual grasses, broadleaf weeds and nutsedge.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
TUBEROUS AND CORM VEGETABLES (Arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; potato; sweet potato; tanier; turmeric; yam	$\frac{1}{2}$ - 1	45	<p>Do not make more than 2 applications per 12 month period.</p> <p>Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre per 12 month period.</p> <p>Do not apply this product by rope-wick or wiper applicators.</p>

bean; yam, true.) Crop Subgroup 1C			
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil.</p> <p>Pre-emergence and Post-emergence applications for control of labeled broadleaf weeds and nutsedge. Use Rate: ½ - 1 oz. Apply a single broadcast application after planting but prior to crop emergence. If needed, make a second post-emergence foliar application 45 days before harvest.</p> <p>For second application, add NIS (1 to 2 quarts) per 100 gal of spray solution.</p> <p>PRECAUTIONS: This product may not control ALS-resistant weeds Use of this product may cause significant, temporary stunting and delay maturity of potatoes resulting in delayed harvest. This product is available to the end-user /grower solely to the extent that the benefit and utility, in the sole opinion of the end-user/grower, outweigh the extent of potential injury associated with the use of this product.</p>			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS	
TURFGRASSES (established lawns, ornamental turfgrass, landscaped areas, commercial and residential turfgrass), AND OTHER NON-CROP SITES (including airports, cemeteries, fallow non-crop areas, golf courses, landscaped areas, public recreation areas, residential property, roadsides, school grounds, sod or turf seed farms, sports fields, landscaped areas with established woody ornamentals, fairgrounds, race tracks, tennis courts, campgrounds and rights-of-way)	⅔ – 1½		Do not apply more than 1½ oz. of this product (0.062 lbs. halosulfuron-methyl) per application. Do not make more than 4 applications per year. Do not apply more than 5½ oz. of this product (0.25 lbs. halosulfuron-methyl) per acre per year. Do not apply this product through any type of irrigation system. Do not apply this product by air. In California: Do not make more than 2 applications per year. Do not apply more than 2½ oz. of this product (0.125 lbs. halosulfuron-methyl) per acre per year. Do not mow turfgrass for 2 days before or 2 days after application for best results. Do not apply this product to golf course putting greens. Do not exceed the specified amount of spray additive due to the potential for turf injury at higher rates. Do not apply this product by rope-wick or wiper applicators.	
Broadcast Treatment: Cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds. Use 0.25 - 0.5% nonionic surfactant (1 - 2 qts. per 100 gallons of spray suspension) for broadcast applications. For high volume applications, do not exceed 1 qt. of spray additive per acre.				
Spot Applications: Add 2 teaspoons (⅓ fl. oz.) of nonionic surfactant per gallon of water. Use only nonionic surfactants which contain at least 80% active material. Refer to the spray additive label and observe all precautions, restrictions, mixing and application instructions.				
Post-emergent Weed Activity Table HALOSULFURON 75 WDG HREBICIDE by Weed Species				
Common Name	Scientific Name	Control	Suppression	Comments
	<i>Kyllinga spp.</i>		YES	

Nutsedge, Yellow	<i>Cyperus esculentus</i>	YES		Heavy infestation requires sequential applications.
Nutsedge, Purple	<i>Cyperus rotundus</i>	YES		Heavy infestation requires sequential applications.

[Mixing Instruction for additional packaging options.]

[INSTRUCTIONS FOR WATER DISPERSIBLE GRANULE IN 1.3 OUNCE BOTTLE] Mix 0.03 ounces (0.9 gram) of this product (using the measuring scoop provided) in 1 - 2 gallons of water to treat 1,000 sq. ft. Add 2 teaspoons (1/8 fluid ounce) of nonionic surfactant per gallon of water. Measure this product as a level and not a rounded scoop. Mix or shake thoroughly for at least two minutes to completely disperse this product. To ensure that this product remains thoroughly mixed while spraying, occasionally shake the spray suspension.

[INSTRUCTIONS FOR WATER DISPERSIBLE GRANULE PREMEASURED IN WATER SOLUBLE BAG:] Use one (1) water soluble bag of this product per 1 - 2 gallons of water in a hand-held or backpack sprayer to treat 1,000 sq. ft. Add 2 teaspoons (1/8 fl. oz.) of a nonionic surfactant per gallon of spray solution. Mix or shake thoroughly for at least two minutes to completely disperse this product. To ensure that this product remains thoroughly mixed while spraying, occasionally shake the spray suspension.

Turfgrass – Use this product on well established seeded, sodded or sprigged turfgrass for the post-emergent control of nutsedge, e.g. yellow and purple. The turf needs to develop a good root system and uniform stand before application. If needed, overseed treated areas with annual or perennial ryegrass or bermudagrass 2 weeks after application.

Broadcast Treatments - After nutsedge has reached the 3 - 8 leaf stage of growth, apply 2/3 - 1 1/2 oz., of this product per acre. For light infestations use the lower rate and heavy infestations use the higher rate.

Sequential Treatments – To maximize the control of nutsedge, a second post-emergent spot or broadcast spray is applied 6 - 10 weeks after the initial treatment to the areas where nutsedge has re-grown or emerged. After nutsedge has reached the 3 - 8 leaf stage of growth, apply 2/3 - 1 1/2 oz., of this product per acre. For light infestations use the lower rate and heavy infestations use the higher rate. Use a spot treatment application for localized control of newly emerged nutsedge. For spot treatments, mix 0.03 oz. (0.9 gram) of this product in 1 - 2 gallons of water to treat 1,000 sq. ft.

Woody Ornamentals in Landscaped Areas

Use this product as a post-directed spray at the specified use rates around established woody ornamental plants in landscaped areas. If applications are to be made to transplanted woody ornamentals, allow 3 months after transplanting before applying this product.

Fallow Treatments

This product may be used on fallow areas prior to establishing turfgrass plants. Wait 4 weeks between application and seeding or sodding of turfgrass.

Precautions:

This product is effective if no rainfall occurs within 3 hours, but best results are obtained with no rainfall or irrigation for at least 4 hours.

When transplanted into landscaped areas treated with this product, flowers, ornamentals plants and shrubs may be injured. Avoid contact of the spray containing this product to desirable flowers, ornamentals, shrubs or trees as discoloration, severe foliar injury or death may result.

Avoid application of this product when turfgrass or nutsedge is under stress since turf injury and poor nutsedge control may occur.

Turfgrass Renovation

For turfgrass renovations, apply at 2/3 oz. per acre in combination with glyphosate herbicide formulations labeled for turfgrass renovation. This is for a non-selective pre-plant burndown of emerged annual grasses, broadleaf weeds and nutsedge.

Wait 4 weeks between application and seeding or sodding of turfgrass.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
ROADSIDES, RIGHTS-OF-WAY, TANK FARMS, LUMBERYARDS, FUEL STORAGE AREAS, FALLOW NON-CROP LAND, AND FENCE ROWS	2⅔		Do not apply more than 2⅔ oz. of this product (0.125 lbs. halosulfuron-methyl) per application. Do not make more than 2 applications per 12-month period. Do not apply more than 5½ oz. of this product (0.25 lbs. halosulfuron-methyl) per acre per 12-month period. Do not apply this product through any type of irrigation system. Do not apply this product by air. Do not apply this product by rope-wick or wiper applicators.

For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds. Use 0.25 - 0.5% nonionic surfactant (1 - 2 quarts per 100 gallons of spray solution) for broadcast applications.

Post-emergent Weed Activity Table
HALOSULFURON 75 WDG HREBICIDE
by Weed Species

Common Name	Scientific Name	Control	Suppression	Comments
Cocklebur, common	<i>Xanthium strumarium</i>		YES	
Horsetail	<i>Equisetum arvense</i>	YES	YES	Control if weeds are less than 6 inches tall. Suppression if weeds are greater than 6 inches tall
Pigweed, redroot	<i>Amaranthus retrofractus</i>		YES	
Pigweed, smooth	<i>Amaranthus hybridus</i>		YES	
Ragweed, common	<i>Ambrosia artemisiifolia</i>		YES	
Ragweed, giant	<i>Ambrosia trifida</i>		YES	
Sunflower	<i>Helianthus annuus</i>		YES	
Velvetleaf	<i>Abutilon theophrasti</i>		YES	

For post-emergence control of horsetail (*Equisetum arvense*), apply 2⅔ oz. of this product per acre or 0.06 oz. (1.8 grams) of this product per 1,000 square feet (0.125 lbs. halosulfuron-methyl per acre) after horsetail has leafed out. Within 14 days after application, signs of herbicide effect will appear as a necrotic ring at the base of the plant, even though the leaves and stems remain green and a deep leathery green in color.

For a non-selective burndown of emerged annual grasses, broadleaf weeds and nutsedge, use this product in combination with glyphosate herbicide formulations labeled for these same uses.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
WATERMELONS	½ - 1	57	For use only in: AL, AR, AZ, CA, CT, DE, FL, GA, IL, IN, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, NH, NJ, NY, OH, OK, PA, RI, SC, TN, TX, VA, VT, WA, WV, WI. Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application per crop cycle. Do not apply more than 2 applications of this product (0.5 oz. (0.023 lbs. halosulfuron-methyl per application) per acre per crop cycle.

	1	<p>Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per year, (includes applications to the crop and to row middle).</p> <p>Rope-wick or Wiper Application Row middle/furrow application – Apply 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre.</p>
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 20 gallons of water per acre.</p> <p>For the best control of nutsedge, apply to nutsedge no larger than 3- 5 leaves. To improve the control of nutsedge, use 2 applications of 0.5 oz. (0.0235 lbs. halosulfuron-methyl) per acre. The 2nd application should be applied 5-7 days after the first application.</p> <p>Direct-seeded: Bare Ground Use Rate: ½ - ¾ oz. <i>Pre-emergence</i> – Apply this product pre-emergence for the suppression of nutsedge and control of listed broadleaf weeds. Use this product after planting, but before cracking. For lighter textured soils with low organic matter, use the lower rate. Where soil is fumigated prior to planting, allow at least five days after soil fumigation before application of this product.</p> <p>Direct-seeded: Plastic mulch Use Rate: ½ - ¾ oz. <i>Pre-seeding</i> – For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-seeding application under the plastic mulch. After final bed shaping and just prior to the installation of the plastic mulch, apply this product. No sooner than 7 days after the application and the installation of the plastic mulch, plant watermelon seeds into this treated area unless local conditions demonstrate safety at an earlier interval. For lighter textured soils with low organic matter, use the lower rate. Soil treated with this product in the planting hole may result in crop injury. During the seeding process, take care to limit movement of soil.</p> <p>Transplanted: Bare ground Use Rate: ½ - ¾ oz. <i>Pre-transplant</i> – For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-transplant application under the plastic mulch. No sooner than 7 days after the application, transplant watermelons into this treated area unless local conditions demonstrate safety at an earlier interval. For lighter textured soils with low organic matter, use the lower rate. This product treated in soil in transplant hole may result in crop injury. During the transplant process, take care to limit movement of soil.</p> <p>Transplanted: Plastic mulch Use Rate: ½ - ¾ oz. <i>Pre-transplant</i> – For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-transplant application under the plastic mulch. After final bed shaping and just prior to the installation of the plastic mulch, apply this product. No sooner than 7 days after the application and the installation of the plastic mulch, transplant watermelons into this treated area unless local conditions demonstrate safety at an earlier interval. Treated soil in the transplant hole may result in crop injury. During the transplant process, take care to limit movement of soil.</p> <p>Direct-seeded and Transplant: Use Rate: ½-1 oz. <i>Row Middle Applications</i> – For the treatment of nutsedge and labeled broadleaf weeds, use this product between rows of direct-seeded or transplanted crop. Avoid contact of the herbicide with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Adjust the rate within the specified rate range and spray volume proportionally to the actual treated area.</p>		

CROP ROTATIONAL GUIDELINES

Following applications of this product, the crop rotational intervals listed below provide for adequate safety to newly planted crops. If the crop is planted in a shorter interval, crop injury may result. If the degradation of halosulfuron-methyl is slowed down by the conditions such as drought, cool conditions or drip irrigation in Arizona and California, the time lines need to be extended. Since all possible environmental and application scenarios, have not been tested, Aceto Life Sciences, [L.L.C. d/b/a Actylis](#) suggests that the end user test this product in order to determine its suitability for such intended use. In

areas where local experience has demonstrated crop safety, use the shorter intervals. In the event of crop failure, labeled crops may be planted back into the treated area at the user's risk for potential phytotoxicity to the subsequent crop.

**TIME INTERVAL(MONTHS) BEFORE PLANTING
AFTER USE OF HALOSULFURON 75WDG HERBICIDE**

CROP	MONTHS	EXCEPTIONS
CROP NOT SPECIFICALLY LISTED	36	
Alfalfa	9	
Apple*	9	
Barley (winter)	2	
Beans, Dry	0	
Beans, Snap	9	In the northeast, midwest and southeast: 2 months; In TX :3 months.
Broccoli	18	In muck soils areas of FL: 3 months.
Blueberry*	9	
Cabbage	15	In muck soils areas of FL: 3 months.
Caneberry*	9	
Canola	15	
Carrot	15	
Cauliflower	18	In muck soils areas of FL: 3 months.
Cereal crops, Spring	2	
Clovers	9	
Collards	18	
Corn, IR/IMR Field	0	
Corn, IT Field	1	
Corn, Normal Field	1	
Corn, Seed	2	
Corn, Sweet and Popcorn	3	For sweet corn and popcorn, the application rates of this product are specific to those crops. For re-planting sweet corn and popcorn crops in those treated areas, that are lost, terminated or harvested, the crop rotational interval must be adhered to.
Cotton	4	
Cucumbers	9	In the northeast and southeast: 2 months; In TX :3 months.
Eggplant	12	For FL transplants: 4 months.
Forage Grasses	2	
Grapes*	9	
Lettuce Crops	18	In muck soils areas of FL: 3 months.
Melons	9	In southeast and TX: 2 months.
Mint	15	
Oats	2	
Onions and Leeks	18	
Peanuts	6	
Pears*	9	
Peas	9	
Peas, Fields	9	
Peppers	10	For FL transplants: 4 months and for TX transplants: 3 months.
Peppers	4	
Potatoes	9	
Pumpkins	9	In southeast: 2 months.
Proso Millet	2	
Radish	12	In muck soils areas of FL: 3 months.
Red Beet	24	If irrigation is required or rainfall is sparse, the time interval is 36 months.
Rice	2	
Rye (winter)	2	
Sorghums	2	
Soybeans	9	
Spinach	24	In muck soils areas of FL: 3 months.
Squash	9	In southeast: 2 months.

Strawberries	36	For annual FL transplants: 6 months.
Sugar beet	24	If irrigation is required or rainfall is sparse, the time interval is 36 months; In MI: 21 months; In MN, ND, Red River Valley: 36 months.
Sugarcane	0	
Sunflowers	18	
Tomato	8	In the northeast and southeast: 2 months; In TX: 3 months.
Tree Nut*	9	
Wheat (winter)	2	

* After application of this product, the soiled must be plowed and cross disked before rotation of crop.

When used with tank mixture partners, consult the partner product labels to determine rotational crop restrictions. Follow the most restrictive label when planning and applying the tank mixture combinations. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

Southeast: AL, FL, GA, LA, MS, NC, Puerto Rico, SC, TN.
Northeast: CT, DE, IA, IL, IN, KY, MA, MD, ME, MI, MN, MO, ND, NE, NH, NJ, NY, OH, PA, RI, SD, VA, VT, WI, WV.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry and secure location. [Add for water soluble packaging] Do not store at temperatures below 32° F (0° C). Rough handling may cause breakage, especially at low temperatures. Do not allow inner bag to become wet during storage. Do not handle inner bag with wet gloves.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

[Plastic bottle packaging:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once triple rinsed, recycle if available. Some agricultural pesticide containers can be taken to a container collection site or pick up for recycling. To find the nearest site, contact your chemical dealer or manufacturer. If recycling is not available, dispose of in a sanitary landfill or by incineration if allowed by state and local ordinances.

[Water Soluble Packaging:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely use water soluble bags in application equipment. Then offer for recycling if available, or dispose of empty foil pouch and cardboard box in a sanitary landfill, or by incineration, or by burning, if allowed by state and local authorities. If burned, stay out of smoke.

[Fiber Drums with Liners:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into the handling or application equipment. Then offer for recycling if available, or dispose of liner in a sanitary landfill, or by incineration, or by burning, if allowed by state and local authorities. If burned, stay out of smoke. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner.

WARRANTY DISCLAIMER AND NOTICE

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aceto Life Sciences, L.L.C. d/b/a Actylis. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ACETO LIFE SCIENCES, L.L.C. D/B/A ACTYLIS MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aceto Life Sciences, L.L.C. d/b/a Actylis is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, ACETO LIFE SCIENCES, L.L.C. D/B/A ACTYLIS DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ACETO LIFE SCIENCES, L.L.C. D/B/A ACTYLIS'S ELECTION, THE REPLACEMENT OF PRODUCT.

CHEMTREC is a registered trademark of American Chemical Society.

All trademarks are the property of their respective owners.

Made in China, packaged in USA

[Advertising claims that may be present on the label:

- Your sedge solution
- Easy Handling Formulation
- Excellent Nutsedge Control
- Tank Mix Compatible
- Economical and Effective
- Easy on Crop Tough on Nutsedge]

[Graduated Scale for Plastic Containers]

[10 oz. Container

SCALE*

OZ.

___ 8

___ 6

___ 4

___ 2

___]

[20 oz. Container

SCALE*

OZ.

___ 18

___ 14

___ 10

___ 6

___ 2]

[*Quantity scale is for approximation only, use weight scale for exact measurement]

002749-00528.20250211.Notification

Page 50 of 65

Formatted: List Paragraph

Turf Sub Label

HALOSULFURON-METHYL	GROUP	2	HERBICIDE
----------------------------	--------------	----------	------------------

HALOSULFURON 75WDG HERBICIDE

HALOSULFURON 75WDG HERBICIDE is an herbicide for selective pre-emergent and post-emergent control of listed weeds including both broadleaf weeds and nutsedge in turfgrasses (established lawns, ornamental turfgrass, landscaped areas, commercial and residential turfgrass), and other non-crop sites (including airports, cemeteries, fallow areas, golf courses, landscaped areas, public recreation areas, residential property, roadsides, school grounds, sod or turf seed farms, sports fields, landscaped areas with established woody ornamentals, fairgrounds, race tracks, tennis courts, campgrounds and rights-of-way).

Read the entire label before using this product.
Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

ACTIVE INGREDIENT:	% BY WT.
Halosulfuron-methyl.....	75%
OTHER INGREDIENTS:.....	25%
TOTAL	100%

EPA Reg. No. 2749-528	Net Contents: 0.03 ounces (0.9 grams) [water-soluble bag]
EPA Est. No. 065387-AR-001	1.3 ounces

Manufactured for:
Aceto Life Sciences, L.L.C. d/b/a Actylis LLC
4 Tri Harbor Court
Port Washington, NY 11050

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUSES MODERATE EYE IRRITATION. HARMFUL IF SWALLOWED. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call poison control center or physician for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call poison control center or physician immediately for treatment advice. • Remove visible particles from mouth. • Have person rinse mouth thoroughly with water, spit out rinse water. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or physician, or going for treatment. FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- long-sleeved shirt and long pants, and
- shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS:

When handlers use closed systems, or enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should:

- Remove PPE immediately after handling this product.
- Wash the outside of the gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FOR CHEMICAL SPILL, LEAK, FIRE, EXPOSURE OR MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887.

ENVIRONMENTAL HAZARDS

This product is toxic to non-target vascular plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Ground Water Advisory: Halosulfuron-methyl is known to leach through soil into ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of halosulfuron-methyl from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Non-target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Aceto Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forest, nurseries and green houses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during this restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves, made of nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils or polyethylene.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

Product Information

HALOSULFURON 75WDG HERBICIDE is a sulfonylurea herbicide that works by inhibition of acetolactate synthase (ALS). Many factors such as application rate, weed species, weed pressure, conditions of weeds including size and climatic factors impact the degree of weed control. Applications made to actively growing weeds at the early stages of development as described below will optimize performance. In post-emergent weed applications, early treatment is best to control the weeds vying (competing) with the crop.

HALOSULFURON 75WDG HERBICIDE is quick to act on targeted weeds by stunting growth allowing the crop to over take the development of the targeted weeds. Once the development of the targeted weeds is stunted, the leaves and growing point begin to discolor and die. Complete control typically occurs within 7 to 14 days depending on the weed size, species and growing conditions. Depending on the stage and development of the targeted weeds, control generally takes place in 7 to 14 days.

Windblown Soil Particles

HALOSULFRUON 75WDG HERBICIDE has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affects the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying HALOSULFRUON 75WDG HERBICIDE if prevailing local conditions may be expected to result in off-site movement.

Weed Resistance Management

For resistance management, HALOSULFRUON 75WDG HERBICIDE is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to HALOSULFRUON 75WDG HERBICIDE and other Group 2 herbicides. The

resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

See specific crop Use Directions for maximum single application rate, annual maximum number of applications and amount of active ingredients.

To delay herbicide resistance take one or more of the following steps:

Rotate the use of HALOSULFRUON 75WDG HERBICIDE or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.

Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

Users should scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

Users should report lack of performance to the registrant or their representative.

Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

Mixing Instructions

HALOSULURON 75WDG HERBICIDE is a water dispersible granule designed to be diluted with water at the rates listed in the specific crop use directions. Fill the spray tank with approximately ½ of the desired volume with water or carrier. With the agitation operating, add the specified amount of the formulation as listed in the targeted crop use directions. Complete the filling process while maintaining agitation. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the carrier source. Add nonionic surfactant and other spray additives as the last ingredients in the tank. Allow time to fully disperse.

Since this product forms a suspension in water, it is important to maintain good agitation during mixing and spraying. If the spray suspension is allowed to settle for a short period of time, be sure to agitate the spray suspension for a minimum 10 minutes. Apply spray solutions within 24 hours after mixing.

[Mixing Instruction for additional packaging options.]

[INSTRUCTIONS FOR WATER DISPERSIBLE GRANULE IN 1.3 OUNCE BOTTLE] Mix 0.03 ounces (0.9 gram) of this product (using the measuring scoop provided) in 1 - 2 gallons of water to treat 1,000 sq. ft. Add 2 teaspoons (⅓ fluid ounce) of nonionic surfactant per gallon of water. Measure this product as a level and not a rounded scoop. Mix or shake thoroughly for at least two minutes to completely disperse this product. To ensure that this product remains thoroughly mixed while spraying, occasionally shake the spray suspension.

[INSTRUCTIONS FOR WATER DISPERSIBLE GRANULE PREMEASURED IN WATER SOLUBLE BAG:] Use one (1) water soluble bag of this product per 1 - 2 gallons of water in a hand-held or backpack sprayer to treat 1,000 sq. ft. Add 2 teaspoons (⅓ fl. oz.) of a nonionic surfactant per gallon of spray solution. Mix or shake thoroughly for at least two minutes

to completely disperse this product. To ensure that this product remains thoroughly mixed while spraying, occasionally shake the spray suspension.

Spray Additives

Spray additives such as nonionic surfactant (NIS) are used with HALOSULFURON 75 WDG HERBICIDE to improve performance. The typical nonionic surfactant contains a minimum of 80% NIS and is accepted by the EPA for use on food crops. The use rate is 0.25 to 0.5% NIS concentrate (1 to 2 quarts per 100 gallons of spray mixture). Always use NIS in the spray mixture.

For specific details, consult the use direction in crop section

Use Rate Equivalency

Since HALOSULFURON 75WDG HERBICIDE contains 75% active ingredient per lb. of product, the following table expresses the use rate equivalency of oz. of this product in term of lbs. halosulfuron-methyl on an acre basis.

oz. of Product per acre	lbs. halosulfuron- methyl per acre
$\frac{1}{2}$	0.0235
$\frac{2}{3}$	0.031
$\frac{3}{4}$	0.035
1	0.047
$1\frac{1}{3}$	0.062
$1\frac{1}{2}$	0.070
2	0.094
$2\frac{2}{3}$	0.125
$5\frac{1}{3}$	0.250

Application Methods

Apply this product by ground to produce uniform coverage on growing weeds or soil to achieve consistent weed control.

Uniform, thorough spray coverage is important to achieve consistent weed control. Calibrate application equipment according to manufacturer's specifications. Use nozzle type arrangements that provide optimum spray distribution and maximum coverage while avoid contact to sensitive crop foliage.

Thoroughly clean application equipment immediately after use and prior to spraying a crop other than corn or grain sorghum. See Spray Equipment Cleanout section of this label for complete details.

Ground Applications

When HALOSULFURON 75WDG HERBICIDE is applied by ground equipment, use in a minimum of 10 gallons of water per acre for a broadcast application. In dense weed populations and thick canopy cover, higher spray volumes are necessary, e.g. 15 – 20 gallons of water per acre. Use the proper spray volume and nozzles that will ensure thorough and uniform coverage of the targeted weeds. Use directed applications to avoid contacting sensitive crop foliage. Select nozzles that will provide optimum spray volume, distribution and coverage at a pressure (psi) that minimizes spray drift. Inspect nozzle distribution during application to avoid streaking and overspray.

SPRAY DRIFT

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

SPRAY DRIFT

Handheld Technology Applications:

- Take precautions to minimize spray drift.

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

- Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Sensitive areas:

Use pesticide products adjacent to sensitive areas only when there is minimal potential for drift or off-target movement, e.g. wind is blowing away from non-target crops, residential areas, known habitats for threatened or endangered species, etc.

In California (only), particularly sensitive crops are identified as cotton and prunes. In applications near these sensitive crops utilize the following buffer zones:

- Ground application shall not be made closer than 1 mile from sensitive crops unless wind direction during the application is away from sensitive crops. When wind direction during the ground application is away from sensitive crops, ground application shall not be made closer than 0.5 miles from sensitive crops.

Spray Equipment Cleanout

The mix tank and spray equipment cleanout is an important stewardship activity to avoid injury to desirable crops. It is important to clean all mixing and spraying equipment immediately after use and before using pesticide products including HALOSULFURON 75WDG HERBICIDE. This is especially important prior to spraying a crop other than grain sorghum and corn.

To clean the spraying equipment, follow the procedure outlined below:

- Completely drain the mix tank and/or sprayer, and then wash thoroughly the tank, sprayer, boom and nozzles with clean water. Drain the system again.
- Fill the mixing or spray tank half full with clean water and add domestic ammonium, normally a 3% solution, at a dilution rate of 1% vol/vol ammonium or 1 gallon per 100 gallons of rinsate.
- Completely fill the tank(s) with additional clean water. Agitate and recirculate and flush out the boom and hoses. Let the system run for 10 – 15 minutes. Drain the system completely.
- Remove nozzles and screens and dislodge any visible solid material. Then soak them in a 1% vol/vol ammonium solution. Inspect the nozzles and screen and remove any visual residues.
- Repeat the above procedure for a second time.
- Flush the mix tank and/or sprayer, boom and hoses with clean water. Drain the system again and inspect for any visible residues. If present, repeat the cleaning cycle again.
- If the rinsate cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Tank Mixtures

To improve this product's effectiveness, apply in combination with other pesticide products that are registered for the same crop and application techniques.

A list of potential herbicide tank mixture partners is provided in the use direction section under each crop. This list is an example of products used but is not an all inclusive list. For current information on the best tank mixture partner in your area, consult with the local dealer, distributor or State Agricultural Extension service.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

If HALOSULFURON 75WDG HERBICIDE is to be tank mixed with other herbicides, conduct a compatibility test prior to mixing. Use a small container and mix all components in a small amount, usually 0.5 to 1 qt. of spray. Combine all products in the same ratio and order of addition as in the proposed spray mixture. Observe the mixture for indication of incompatibility which usually occurs in 10 to 30 minutes after mixing. If incompatibility is observed, try changing the order of addition of the components. The guideline on tank mixture partners is driven by formulation type. Start with wettable powders (WP's) including water soluble bags (WSB's), water dispersible granules (WDG's), suspension concentrated (SC's) or flowable (F's), all with very good agitation. Next follow with water miscible concentrates and emulsifiable concentrates (EC's) before adding drift control additives, nonionic surfactants (NIS's) or crop oil concentrates (COC's). After vigorous agitation, there

must be a homogeneous suspension. Let the final tank mixture stand and observe for any rapid settling or floating of components. If any indications of physical incompatibility develop, do not use this mixture for spraying.

Application Restrictions

- Do not use air assisted (air blast) sprayers to apply this product.
- Do not apply this product through any type of irrigation system.
- Do not apply more than 5½ oz. of this product (0.25 lbs. halosulfuron-methyl) per acre per year on turf.
- Do not allow this product to drift outside of targeted area.
- Do not apply tank mixtures if the turf is under severe stress due to drought, water-saturated soils, poor fertility (especially low nitrogen levels), hail, frost, insects or when the maximum daytime temperature is above 92° F.
- Do not use if the target weeds or turf are under stress due to drought, water saturated soils, low fertility (especially low nitrogen levels) or other poor growing conditions.

Application Precautions

- Increase in turf injury may result if the seeding depth is too shallow and excessive amounts of water (greater than 1 inch) from rainfall or sprinkler irrigation occurs.
- Avoid spraying when conditions favor rainfall or using overhead sprinkler irrigation within 4 hours of this application.
- Loss in effectiveness or crop injury may result if weeds are under drought, stress, disease or insect damage.
- Under cool and wet growing conditions that delay early seedling emergence, vigor or growth, this product may cause injury or crop failure. These conditions are likely to occur during the first planting of the season.
- The maturity of the turf may be delayed by use of this product.
- Use nozzles and pressures that minimize the production of fine particles that drift outside of the targeted area.
- Applications of this product may cause temporary yellowing or stunting of the turf.
- In California and Arizona due to environmental conditions that delay degradation of this product, extend the crop rotation intervals on drip irrigated crops.
- When this product is applied over-the-top of a blooming turf, bloom loss may occur under certain environmental conditions.
- If rainfall or irrigation occurs within 4 hours after application, reduce effectiveness may occur.
- Avoid disturbing (e.g. cultivation) treated areas for at least 7 days following application.

For Best Performance

Many factors such as application rate, weed species, weed pressure, conditions of weeds including size and climatic conditions impact the degree of weed control. Applications made to actively growing weeds at the early stages of development as described below will optimize performance. In post-emergent weed applications, early treatment is best to control the weeds vying (competing) with the crop. For residual control from early post-emergent treatments a second application may be needed to control later germination of weeds.

HALOSULFURON 75WDG HERBICIDE is quick to act on targeted weeds by stunting growth allowing the crop to overtake the development of the targeted weeds. Once the development of the targeted weeds is stunted, the leaves and growing point begin to discolor and die. Complete control typically occurs within 7 to 14 days depending on the weed size, species and growing conditions. Depending on the stage and development of the targeted weeds, control generally takes place in 7 to 14 days.

When using spray additives, carefully follow the listed use instructions.

- In post-emergence applications:
 - Better control is obtained when applied early to actively growing, small (1-3 inches in height) broadleaf weeds. Large broadleaf weeds may not be adequately controlled.
 - Nutsedge plants are best controlled at the actively growing, 3 - 5 leaf stage.
 - After a post-emergence application, delay overhead sprinkler irrigation for 2 to 3 days.
 - If weeds are under drought, stress, disease, or insect damage, do not use.
- Under heavy weed infestation, use early before the weeds become too competitive with the crop.
- Annual weeds may have multiple flushes of seedlings, or treated perennials may sometimes re-grow from underground stems or roots, depending upon rainfall and other environmental conditions. To maximize control of such weeds, apply a sequential application of this product.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
TURFGRASSES (established lawns, ornamental turfgrass, landscaped areas, commercial and residential turfgrass), AND OTHER NON-CROP SITES (including airports, cemeteries, fallow non-crop areas, golf courses, landscaped areas, public recreation areas, residential property, roadsides, school grounds, sod or turf seed farms, sports fields, landscaped areas with established woody ornamentals, fairgrounds, race tracks, tennis courts, campgrounds and rights-of-way)	2/3 – 1 1/3		Do not apply more than 1 1/3 oz. of this product (0.062 lbs. halosulfuron-methyl) per application. Do not make more than 4 applications per year. Do not apply more than 5 1/3 oz. of this product (0.25 lbs. halosulfuron-methyl) per acre per year. Do not apply this product through any type of irrigation system. Do not apply this product by air. In California: Do not make more than 2 applications per year. Do not apply more than 2 2/3 oz. of this product (0.125 lbs. halosulfuron-methyl) per acre per year. Do not mow turfgrass for 2 days before or 2 days after application for best results. Do not apply this product to golf course putting greens. Do not exceed the specified amount of spray additive due to the potential for turf injury at higher rates. Do not apply this product by rope-wick or wiper applicators.

Broadcast Treatment:
Cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds. Use 0.25 - 0.5% nonionic surfactant (1 - 2 qts. per 100 gallons of spray suspension) for broadcast applications. For high volume applications, do not exceed 1 qt. of spray additive per acre.

Spot Applications:
Add 2 teaspoons (1/3 fl. oz.) of nonionic surfactant per gallon of water. Use only nonionic surfactants which contain at least 80% active material.
Refer to the spray additive label and observe all precautions, restrictions, mixing and application instructions.

Post-emergent Weed Activity Table
HALOSULFURON 75 WDG HREBICIDE
by Weed Species

Common Name	Scientific Name	Control	Suppression	Comments
	<i>Kyllinga spp.</i>		YES	
Nutsedge, Yellow	<i>Cyperus esculentus</i>	YES		Heavy infestation requires sequential applications.
Nutsedge, Purple	<i>Cyperus rotundus</i>	YES		Heavy infestation requires sequential applications.

Turfgrass – Use this product on well established seeded, sodded or sprigged turfgrass for the post-emergent control of nutsedge, e.g. yellow and purple. The turf needs to develop a good root system and uniform stand before application. If needed, oversee d treated areas with annual or perennial ryegrass or bermudagrass 2 weeks after application.

Broadcast Treatments - After nutsedge has reached the 3 - 8 leaf stage of growth, apply 2/3 - 1 1/3 oz., of this product per acre. For light infestations use the lower rate and heavy infestations use the higher rate.

Sequential Treatments – To maximize the control of nutsedge, a second post-emergent spot or broadcast spray is applied 6 - 10 weeks after the initial treatment to the areas where nutsedge has re-grown or emerged. After nutsedge has reached the 3 - 8 leaf stage of growth, apply 2/3 - 1 1/3 oz., of this product per acre. For light infestations use the lower rate and heavy infestations use the higher rate. Use a spot treatment application for localized control of newly emerged nutsedge. For spot treatments, mix 0.03 oz. (0.9 gram) of this product in 1 - 2 gallons of water to treat 1,000 sq. ft.

Woody Ornamentals in Landscaped Areas

Use this product as a post-directed spray at the specified use rates around established woody ornamental plants in landscaped areas. If applications are to be made to transplanted woody ornamentals, allow 3 months after transplanting before applying this product.

Fallow Treatments

This product may be used on fallow areas prior to establishing turfgrass plants. Wait 4 weeks between application and seeding or sodding of turfgrass.

Precautions: This product is effective if no rainfall occurs within 3 hours, but best results are obtained with no rainfall or irrigation for at least 4 hours.

When transplanted into landscaped areas treated with this product, flowers, ornamentals plants and shrubs may be injured. Avoid contact of the spray containing this product to desirable flowers, ornamentals, shrubs or trees as discoloration, severe foliar injury or death may result.

Avoid application of this product when turfgrass or nutsedge is under stress since turf injury and poor nutsedge control may occur.

Turfgrass Renovation

For turfgrass renovations, apply at $\frac{2}{3}$ oz. per acre in combination with glyphosate herbicide formulations labeled for turfgrass renovation. This is for a non-selective pre-plant burndown of emerged annual grasses, broadleaf weeds and nutsedge.

Wait 4 weeks between application and seeding or sodding of turfgrass.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use.

Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
ROADSIDES, RIGHTS-OF-WAY, TANK FARMS, LUMBERYARDS, FUEL STORAGE AREAS, FALLOW NON-CROP LAND, AND FENCE ROWS	2 $\frac{1}{2}$		Do not apply more than 1 $\frac{1}{2}$ oz. of this product (0.062 lbs. halosulfuron-methyl) per application. Do not make more than 2 applications per 12-month period. Do not apply more than 5 $\frac{1}{2}$ oz. of this product (0.25 lbs. halosulfuron-methyl) per acre per 12-month period. Do not apply this product through any type of irrigation system. Do not apply this product by air. Do not apply this product by rope-wick or wiper applicators.

For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds. Use 0.25 - 0.5% nonionic surfactant (1 - 2 quarts per 100 gallons of spray solution) for broadcast applications.

**Post-emergent Weed Activity Table
HALOSULFURON 75 WDG HERBICIDE
by Weed Species**

Common Name	Scientific Name	Control	Suppression	Comments
Cocklebur, common	<i>Xanthium strumarium</i>		YES	
Horsetail	<i>Equisetum arvense</i>	YES	YES	Control if weeds are less than 6 inches tall. Suppression if weeds are greater than 6 inches tall
Pigweed, redroot	<i>Amaranthus retrofractus</i>		YES	
Pigweed, smooth	<i>Amaranthus hybridus</i>		YES	
Ragweed, common	<i>Ambrosia artemisiifolia</i>		YES	
Ragweed, giant	<i>Ambrosia trifida</i>		YES	
Sunflower	<i>Helianthus annuus</i>		YES	
Velvetleaf	<i>Abutilon theophrasti</i>		YES	

For post-emergence control of horsetail (*Equisetum arvense*), apply 2½ oz. of this product per acre or 0.06 oz. (1.8 grams) of this product per 1,000 square feet (0.125 lbs. halosulfuron-methyl per acre) after horsetail has leafed out. Within 14 days after application, signs of herbicide effect will appear as a necrotic ring at the base of the plant, even though the leaves and stems remain green and a deep leathery green in color.

For a non-selective burndown of emerged annual grasses, broadleaf weeds and nutsedge, use this product in combination with glyphosate herbicide formulations labeled for these same uses.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry and secure location. [Add for water soluble packaging] Do not store at temperatures below 32° F (0° C). Rough handling may cause breakage, especially at low temperatures. Do not allow inner bag to become wet during storage. Do not handle inner bag with wet gloves.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

[Plastic bottle packaging:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once triple rinsed, recycle if available. Some agricultural pesticide containers can be taken to a container collection site or pick up for recycling. To find the nearest site, contact your chemical dealer or manufacturer. If recycling is not available, dispose of in a sanitary landfill or by incineration if allowed by state and local ordinances.

[Water Soluble Packaging:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely use water soluble bags in application equipment. Then offer for recycling if available, or dispose of empty foil pouch and cardboard box in a sanitary landfill, or by incineration, or by burning, if allowed by state and local authorities. If burned, stay out of smoke.

WARRANTY DISCLAIMER AND NOTICE

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aceto Life Sciences, L.L.C. d/b/a Actylis~~L.L.C.~~. To the extent consistent with applicable law all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ACETO LIFE SCIENCES, L.L.C. D/B/A 002749-00528.20250211.Notification

| ~~ACTYLIS^{LLC}~~ MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF
| FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE
| ON THIS LABEL. No agent of Aceto Life Sciences, ~~L.L.C. d/b/a Actylis^{LLC}~~ is authorized to make any warranties beyond
| those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, ACETO
| LIFE SCIENCES, ~~L.L.C. D/B/A ACTYLIS^{LLC}~~ DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL,
| INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

| **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE
| USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR
| HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT
| LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ACETO LIFE SCIENCES,
| ~~L.L.C. D/B/A ACTYLIS^{LLC}~~'S ELECTION, THE REPLACEMENT OF PRODUCT.

CHEMTREC is a registered trademark of American Chemical Society.

Made in China, packaged in USA

Advertising claims that may be present on the label:

- Your sedge solution
- Easy Handling Formulation
- Excellent Nutsedge Control
- Tank Mix Compatible
- Economical and Effective
- Easy on Crop Tough on Nutsedge

Supplemental Labeling

HALOSULFURON 75WDG HERBICIDE

~~This supplemental labeling includes the use on watermelons and dry beans.~~

~~The labeling must be in possession of the user at the time of application.~~

~~This supplemental label expires on September 27, 2024 and must not be used or distributed after this date.~~

~~Read the label affixed to the container for [Halosulfuron 75WDG Herbicide] before applying.~~

~~Read the entire label before using this product. Use only as according to label instructions.~~

~~Read "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" before buying or using. If terms are not acceptable, return product at once unopened.~~

~~It is a violation of Federal law to use this product in a manner inconsistent with its labeling.~~

~~Use of [HALOSULFURON 75WDG HERBICIDE] according to this labeling is subject to the use precautions and restrictions imposed by the label affixed to the container for [HALOSULFURON 75WDG HERBICIDE].~~

ACTIVE INGREDIENT: _____ % BY WT.
Halosulfuron-methyl.....75%
OTHER INGREDIENTS:.....25%
TOTAL.....100%

~~See product label for complete Directions for Use, Precautions, Restrictions, First Aid, Storage and Disposal.~~

EPA Reg. No. 2749-528

CAUTION

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
WATERMELONS	$\frac{1}{2}$ –1	57	For use only in: AL, AR, AZ, CA, CT, DE, FL, GA, IL, IN, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, NH, NJ, NY, OH, OK, PA, RI, SC, TN, TX, VA, VT, WA, WV, WI. Do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application per crop cycle. Do not apply more than 2 applications of this product (0.5 oz. (0.023 lbs. halosulfuron-methyl per application) per acre per crop cycle. Do not apply more than 2 oz. of this product (0.094 lbs. halosulfuron-methyl) per acre per year, (includes applications to the crop and to row middle). Rope-wick or Wiper Application Row middle/furrow application—Apply 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per acre.
For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 20 gallons of water per acre. For the best control of nutsedge, apply to nutsedge no larger than 3–5 leaves. To improve the control of nutsedge, use 2 applications of 0.5 oz. (0.0235 lbs. halosulfuron-methyl) per acre. The 2nd application should be applied 5–7 days after the first application. Direct-seeded: Bare Ground Use Rate: $\frac{1}{2}$ – $\frac{3}{4}$ oz.			

Pre-emergence—Apply this product pre-emergence for the suppression of nutsedge and control of listed broadleaf weeds. Use this product after planting, but before cracking. For lighter textured soils with low organic matter, use the lower rate. Where soil is fumigated prior to planting, allow at least five days after soil fumigation before application of this product.

Direct-seeded: Plastic mulch

Use Rate: $\frac{1}{2}$ — $\frac{3}{4}$ oz.

Pre-seeding—For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-seeding application under the plastic mulch. After final bed shaping and just prior to the installation of the plastic mulch, apply this product. No sooner than 7 days after the application and the installation of the plastic mulch, plant watermelon seeds into this treated area unless local conditions demonstrate safety at an earlier interval. For lighter textured soils with low organic matter, use the lower rate. Soil treated with this product in the planting hole may result in crop injury. During the seeding process, take care to limit movement of soil.

Transplanted: Bare ground

Use Rate: $\frac{1}{2}$ — $\frac{3}{4}$ oz.

Pre-transplant—For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-transplant application under the plastic mulch. No sooner than 7 days after the application, transplant watermelons into this treated area unless local conditions demonstrate safety at an earlier interval. For lighter textured soils with low organic matter, use the lower rate. This product treated in soil in transplant hole may result in crop injury. During the transplant process, take care to limit movement of soil.

Transplanted: Plastic mulch

Use Rate: $\frac{1}{2}$ — $\frac{3}{4}$ oz.

Pre-transplant—For the suppression of nutsedge and control of labeled broadleaf weeds, use this product as a pre-transplant application under the plastic mulch. After final bed shaping and just prior to the installation of the plastic mulch, apply this product. No sooner than 7 days after the application and the installation of the plastic mulch, transplant watermelons into this treated area unless local conditions demonstrate safety at an earlier interval. Treated soil in the transplant hole may result in crop injury. During the transplant process, take care to limit movement of soil.

Direct-seeded and Transplant:

Use Rate: $\frac{1}{2}$ –1 oz.

Row-Middle Applications—For the treatment of nutsedge and labeled broadleaf weeds, use this product between rows of direct-seeded or transplanted crop. Avoid contact of the herbicide with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Adjust the rate within the specified rate range and spray volume proportionally to the actual treated area.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
DRY BEANS	$\frac{1}{2}$ –1	30	<p>Do not apply more than 1 oz. of this product (0.047 lb. halosulfuron-methyl) per application.</p> <p>Do not make more than 2 applications per year.</p> <p>Do not make more than 1 post-emergence application.</p> <p>In CA, only apply as post-emergence directed spray to Middle Row/Furrows.</p> <p>For Direct-Seeded Pre-emergences and Post-emergence applications, do not apply more than $\frac{2}{3}$ oz. of this product (0.031 lbs. halosulfuron-methyl) per application.</p> <p>Do not apply more than 2 oz. (0.094 lb. halosulfuron-methyl) per acre per 12-month period (includes applications to the crop and to Row-Middle/Furrows).</p> <p>For Row-Middle/Furrow applications, do not apply more than 1 oz. of this product (0.047 lbs. halosulfuron-methyl) per application.</p> <p>Do not apply this product by rope-wick or wiper applicators.</p>
<p>For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.</p> <p>Direct-Seeded Pre-emergence: Use Rate: $\frac{1}{2}$—$\frac{3}{4}$ oz.</p>			

~~Use this product after planting, but before cracking. For lighter textured soils with low organic matter, use the lower rate.~~

~~Direct-Seeded Post-emergence:~~

~~Use Rate: 1/2–3/4 oz.~~

~~Use this product when plants have 1-3 trifoliate leaves, but before flowering. For best results, apply to weeds less than 6 inches high. Use a nonionic surfactant (NIS).~~

~~Not all hybrids/varieties have been tested for sensitivity to this product. For untested varieties, a small amount of the field should be sprayed to determine potential sensitivity to its use. To the extent consistent with applicable law, the user assumes responsibility for such use and any plant injury that may occur.~~

~~Row-Middle/Furrow Applications:~~

~~Use Rate: 1/2–1 oz.~~

~~Apply this product between rows of crop for the control of nutsedge and labeled broadleaf weeds. Avoid contact of this product with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.~~

~~Tank Mixture Partner~~

~~A tank mixture of HALOSULFURON 75WDG HERBICIDE partnered with Eptam® 7-E, EPA Reg. No. 10163-283 (EPTC) will provide a broader spectrum of weed control than either product used separately.~~

~~For post-emergence grass control, use partners including but not limited to Targa®, EPA Reg. No. 33906-9 (quizalofop-P-ethyl) or other graminicides.~~

~~It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.~~

~~Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.~~

~~Use HALOSULFURON 75WDG HERBICIDE at a rate of 1/2–3/4 oz. with labeled rate of Eptam® 7-E, EPA Reg. No. 10163-283 (EPTC) and incorporate into the soil at a depth of approximately 2 inches before planting. If any crust appears on the soil, break it up by lightly rotary hoeing during or shortly after the emergence of the beans.~~

~~For lighter textured soils with low organic matter, use the lower rate.~~