



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (7505P)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

**EPA Reg. Number:**  
 2749-594

**Date of Issuance:**  
 8/25/20

**NOTICE OF PESTICIDE:**

Registration  
 Reregistration  
 (under FIFRA, as amended)

**Term of Issuance:**  
 Conditional

**Name of Pesticide Product:**  
 Aceto Etoxazole 72% WSP  
 Miticide

**Name and Address of Registrant (include ZIP Code):**

John F. Wright  
 Agent for Aceto Life Sciences, LLC  
 c/o Product & Regulatory Assoc. LLC  
 8595 Collier Blvd., Suite 107-51  
 Naples, FL 34114

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

**Signature of Approving Official:**

Venus Eagle, Product Manager 01  
 Invertebrate and Vertebrate Branch 3  
 Registration Division (7505P)

**Date:**

8/25/20

2. You are required to comply with the data requirements described in the DCI identified below:

a. Etoxazole GDCI-107091

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:

- Revise the EPA Registration Number to read, "EPA Reg. No. 2749-594."

4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 7/23/2020

If you have any questions, please contact Marianne Lewis by phone at (703) 308-8043 or via email at [lewis.marianne@epa.gov](mailto:lewis.marianne@epa.gov).

Enclosure

ETOXAZOLE	GROUP	<b>10B</b>	INSECTICIDE
-----------	-------	------------	-------------

## Aceto Etoxazole 72% WSP Miticide


Miticide for use on Caneberry, Christmas Trees, Cotton, Cucurbit Vegetables, Field Corn, Popcorn, Corn (Grown for Seed), Hops, Low Growing Berry, Mint, Non-Bearing Fruit Trees, Pepper and Eggplant, Pome Fruit, Small Fruit Vine Climbing (except Fuzzy Kiwifruit), Stone Fruit, Tree Nuts, and Tropical and Sub-Tropical Fruit (Inedible Peel).

	% BY WT
<b>ACTIVE INGREDIENT:</b>	
Etoxazole* .....	72.0%
<b>OTHER INGREDIENTS:</b> .....	28.0%
Total	100.0%
*2-(2,6-difluorophenyl)-4-[4-(1,1-dimethylethyl)-2-ethoxyphenyl]-4,5-dihydrooxazole	

EPA Reg. No. 2749-XXX  
EPA Est. No.

Net Contents:

Manufactured for:  
Aceto Life Sciences, LLC  
4 Tri Harbor Court  
Port Washington, NY 11050



See inside booklet for Precautionary Statements and Directions for Use

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>Call poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

**Emergency Assistance:** Have the product container or label with you when calling a poison control center or doctor or going for treatment. FOR EMERGENCY MEDICAL HELP, CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887.

FOR CHEMICAL SPILL, LEAK, FIRE, EXPOSURE OR MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION.** Causes moderate eye irritation. Harmful if inhaled. Avoid contact with eyes, skin and clothing. Avoid Breathing dust.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made out of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride ≥14 mils, or viton ≥14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. The wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product, Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENGINEERING CONTROLS

Water soluble packets, when used correctly, qualify as a closed mixing/loading system under the Worker Protection Standard [40 CFR 170.607(d)]. Mixers and loaders handling this product while it is enclosed in intact water-soluble packets may elect to wear reduced PPE of long-sleeved shirt, long pants, shoes, socks, a chemical-resistant apron, and chemical-resistant gloves. When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater and marine/estuarine aquatic invertebrates, including oysters and shrimp. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Do not apply this product in a way that it will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

#### **TANK MIXES**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### **CHEMIGATION**

**RESTRICTION: Do not apply this product through any type of irrigation or chemigation system.**

#### RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance-management, **ACETO ETOXAZOLE 72% WSP MITICIDE** contains a Group 10B insecticide. Any insect population may contain individuals naturally resistant to **ACETO ETOXAZOLE 72% WSP MITICIDE** and other Group 10B insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed. To reduce the potential for developing insect resistance, rotate to an insecticide with a different mode of action. Monitor treated pest populations for resistance development. Read product label before applying any insecticide and follow label directions.

To delay insecticide resistance, take the following steps:

- Rotate the use of **ACETO ETOXAZOLE 72% WSP MITICIDE** or other Group 10B insecticides within a growing season, or among growing seasons, with different groups that control the same pests. Avoid application of more than the maximum seasonal use rate or the total number of consecutive sprays of **ACETO ETOXAZOLE 72% WSP MITICIDE** per season.
- Use insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
  - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Aceto Life Sciences, LLC representatives at [aceto.com](http://aceto.com).

## WATER SOLUBLE PACAKGING INFORMATION

### MIXING INSTRUCTIONS

Follow the steps below when mixing this product, including if tank mixed with other pesticide products. If being tank mixed, the mixing directions 1 through 9 below take precedence over the mixing directions of the other tank mix products. WSPs may, in some cases, be mixed with other pesticide products so long as the directions for use of all mixed products do not conflict. Do not tank mix this product with products that prohibit tank mixing or have conflicting mixing directions.

1. If a basket or strainer is present in the tank hatch, remove prior to adding the WSP to the tank.
2. Fill tank with water to approximately one-third to one-half of the desired final volume of spray.
3. Stop adding water and stop any agitation.
4. Place intact/unopened WSP(s) into the tank.
5. Do not spray water from a hose or fill pipe to break or dissolve the WSP(s).
6. Start mechanical and recirculation agitation from the bottom of tank without using any overhead recirculation, if possible. If overhead recirculation cannot be turned off, close the hatch before starting agitation.
7. Dissolving the WSP(s) may take up to 5 minutes or longer, depending on water temperature, water hardness and intensity of agitation.
8. Stop agitation before tank lid is opened.
9. Open the lid to the tank, exercising caution to avoid contact with dusts or spray mix, to verify that the WSPs have fully dissolved and the contents have been thoroughly mixed into the solution.
10. Do not add other allowed products or complete filling the tank until the bags have fully dissolved and pesticide is thoroughly mixed.
11. Once the WSP have fully dissolved and any other products have been added to the tank, resume filling the tank with water to the desired level, close the tank lid, and resume agitation.
12. Use the spray solution when mixing is complete.
13. Maintain agitation of the diluted pesticide mix during transport and application.
14. It is unlawful to use any registered pesticide, including WSPs, in a manner inconsistent with its label.

<b>Number of ACETO ETOXAZOLE 72% WSP MITICIDE Water Soluble Packets Required to Treat</b>		
<b>Rate (oz/A)</b>	<b>Number of Acres to be Treated</b>	<b>Number of 3 oz. Packets Required</b>
0.66	45	10
1.0	30	10
2.0	15	10
3.0	10	10
4.0	7.5	10

### Aceto Etoxazole 72% WSP Miticide Crop/Use Site Index

<b>Crop</b>	<b>Page</b>
Caneberry	<b>7</b>
Christmas Trees,	<b>8</b>
Cotton	<b>9</b>
Cucurbit Vegetables	<b>10</b>
Field Corn, Popcorn, Corn (Grown for Seed Production)	<b>11</b>
Hops	<b>12</b>

Low Growing Berry	<b>13</b>
Mint (Peppermint and Spearmint)	<b>14</b>
Non-Bearing Fruit Trees	<b>15</b>
Pepper and Eggplant	<b>16</b>
Pome Fruit	<b>17</b>
Small Fruit Vine Climbing (except Fuzzy Kiwifruit)	<b>18</b>
Stone Fruit	<b>19</b>
Tree Nuts	<b>20</b>
Tropical and Sub-Tropical Fruit (Inedible Peel).	<b>21</b>



<b>CANEBERRY (Subgroup 13-07 A)</b>			
<b>CROP</b>	<b>PESTS</b>	<b>PRODUCT RATE OZ/ACRE</b>	<b>INSTRUCTIONS</b>
Blackberry Loganberry Raspberry, Black Raspberry, Red Raspberry, Wild Cultivars, varieties and/or hybrids of these	European Red Spider Mite McDaniel Spider Mite Two-spotted Spider Mite Yellow Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	<p>Apply by ground as a full coverage spray in a minimum of 50 gals/a of water. applications of this product must be done using enough carrier (water) to ensure thorough coverage of the crop's vegetative and reproductive parts, which mites are using for dispersal, feeding and reproduction. Use higher water volumes on older trees and varieties that have more compact and dense foliage. Higher volumes of water coupled with well calibrated equipment will help to provide better coverage.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide. Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Do not use below use rate 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
			<p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 0 days.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 3.0 oz (0.135 lb. ai) of this product per acre per calendar year.</li> </ul>

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state, and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

FIELD CORN, POPCORN, CORN (Grown for Seed Production)			
CROP	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Field Corn Popcorn Corn (grown for seed production)	Banks Grass Mite <sup>A</sup> Carmin Spider Mite Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite	1.0 to 3.0 (0.045 to 0.135 lb. ai/A)	<p>Apply with [air* or] ground equipment in adequate water for uniform coverage ([a minimum of 3 gals/A by air* or] a minimum of 10 gals/A [by ground]). Coverage is essential for good control. Use of higher water volume will assure better coverage. Applications of this product are recommended before tasseling stage to allow good coverage and provide better control.</p> <p>For field corn utilized for seed production apply this product before or at tasseling growth stage. if a second application is needed wait for 14 days to re-treat with this product</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>Best results are achieved when mite populations are below threshold or beginning to build up on the plants. This product is predominately an ovicide/larvicide and should be used early in the life cycle of mites. Always follow the recommended threshold for your area.</p> <p>Do not use rates below 1.0 oz/A (0.045 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
[*Not approved for aerial application in New York.]			
<p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>• Do not apply within twenty-one (21) days of harvest.</li> <li>• Do not make more than two (2) this product applications per season.</li> <li>• Do not apply treatments less than 14 days apart.</li> <li>• Do not apply more than 6.0 oz (0.27 lb. ai) of this product per acre per calendar year.</li> </ul>			

<sup>A</sup>Applications targeting exclusively Banks grass mites may require higher rates within the rate range of 1.0 to 3.0 oz/A particularly if populations have exceeded the established threshold in the fields to be treated with this product. Also, under extreme drought conditions and higher populations Banks grass mites might require more than one application of a miticide.

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with other miticides that have different modes of action and/or with other chemical classes of miticides. Follow local, state and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

COTTON			
CROP	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Cotton	Carmine Spider Mite Pacific Spider Mite Two- spotted Spider Mite	0.66 to 1.0 (0.03 to 0.045 lb. ai/A)	<p>Apply with air or ground equipment in adequate water for uniform coverage (3 to 10 gals/A by air or 10 to 50 gals/A by ground). Coverage is essential for good control. Use of higher water volume will assure better coverage.</p> <p>Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide and should be used early in the life cycle of mites. Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>Do not use below use rate 0.66 oz/A (0.03 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
<p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 28 days.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 1.0 oz (0.045 lb. ai) of this product per acre per calendar year.</li> </ul>			

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

CUCURBIT VEGETABLES (Crop Group 9)			
CROP	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Acorn Squash Balsam Apple Balsam Pear Bitter Melon Butternut Squash Calabaza Squash Cantaloupe Chayote (fruit) Chinese Cucumber Chinese Okra Chinese Waxgourd (Chinese Preserving Melon) Citron Melon Cucumber Cucuzza Gherkin Gourd, Edible Hechima Hubbard Squash Hyotan <i>Momordica</i> spp. Muskmelon Pumpkin Spaghetti Squash Summer Squash Watermelon Winter Squash	Two-spotted Spider Mite Carmine Spider Mite Pacific Spider Mite Strawberry Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	Apply with [air* or] ground equipment in adequate water for uniform coverage ([a minimum of 3 gals/A by air* or] a minimum of 10 gals/A [by ground]). Coverage is essential for good control. Use of higher water volume will assure better coverage.  Applications of this product must be made using enough carrier (water) to ensure thorough coverage of the crop's vegetative and reproductive parts, which mites use for dispersal, feeding and reproduction. Use higher water volumes on more mature plants and varieties that have more compact and dense foliage. Higher volumes of water coupled with well calibrated equipment will help to provide better coverage.  Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.  Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide. Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.  Do not use below use rate 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.
[*Not approved for aerial application in New York.]			
<b>Restrictions</b> <ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 7 days.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 3.0 oz (0.135 lb. ai) of this product per acre per calendar year.</li> </ul>			

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

HOPS			
CROP	PEST	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Hops	Two-spotted Spider Mite	3.0 to 4.0 (0.135 to 0.18 lb. ai/A)	<p>Apply with ground equipment in a minimum of 50 gals/A of water. Coverage is essential for good control. Use of higher water volume will assure better coverage.</p> <p>Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide.</p> <p>Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>Do not use below use rate of 3.0 oz/A (0.135 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
<p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 7 days.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 4.0 oz (0.18 lb. ai) of this product per acre per calendar year.</li> </ul>			

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

<b>LOW GROWING BERRY (Subgroup 13-07G)</b>			
<b>CROP</b>	<b>PEST</b>	<b>PRODUCT RATE OZ/ACRE</b>	<b>INSTRUCTIONS</b>
Bearberry Bilberry Blueberry, Lowbush Cloudberry Cranberry Lingonberry Muntries Partridgeberry Strawberry Cultivars, varieties and/or hybrids of these	European Red Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	<p>Apply with ground equipment in a minimum of 100 gals/A of water. Coverage is essential for good control. Use of higher water volume will assure better coverage.</p> <p>Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide and should be used early in the life cycle of mites. Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>This product will not control Cyclamine Mite. Another miticide registered for this pest should be used if these mites are a problem.</p> <p>Do not use below use rate of 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
	Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A) + Labeled rate of registered fenpropathrin product	<p>Apply with ground equipment in adequate water for uniform coverage (using a minimum of 100 gals/A.)</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>Alternate with other non-pyrethroid insecticides if retreatment is needed in less than 30 days to comply with local IPM programs.</p> <p>Comply with all applicable directions, restrictions and precautions on the registered label for any product used in a tank mix.</p>

	<p><b>Restrictions</b></p> <ul style="list-style-type: none"><li>• Preharvest interval (PHI) is 1 day.</li><li>• Do not make more than one (1) this product application per season.</li><li>• Do not apply more than 3.0 oz (0.135 lb. ai) of this product per acre per calendar year.</li></ul>
--	--

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state, and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

MINT			
CROP	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Mint (Peppermint and Spearmint)	Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite	2.0 to 4.0 (0.09 to 0.18 lb. ai/A)	<p>Apply with [air* or] ground equipment in adequate water for uniform coverage (minimum of [10 gals/A by air* or] 50 gals/A [by ground]). Coverage is essential for good control. Use of higher water volume will assure better coverage.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide. Apply this product miticide at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Do not use below use rate of 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
[*Not approved for aerial application in New York.]			
<b>Restrictions</b> <ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 7 days.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 4.0 oz (0.18 lb. ai) of this product per acre per calendar year.</li> </ul>			

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.



PEPPER AND EGGPLANT (Subgroup 8-10B)			
CROP	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
African Eggplant Bell Pepper Eggplant Martynia Non-Bell Pepper Okra Pea Eggplant Pepino Roselle Scarlet Eggplant Cultivars, varieties and/or hybrids of these	Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	<p>Apply with ground equipment in a minimum of 20 gals/A of water. Applications of this product miticide must be made using enough carrier (water) to ensure thorough coverage of the crop's vegetative and reproductive parts, which mites use for dispersal, feeding and reproduction. Use higher water volumes on more mature plants and varieties that have more compact and dense foliage. Higher volumes of water coupled with well calibrated equipment will help to provide better coverage.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide. Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Do not use below use rate 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
<p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 7 days.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 3.0 oz (0.135 lb. ai) of this product per acre per calendar year.</li> </ul>			

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state, and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

POME FRUIT (Crop Group 11)			
CROPS	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Apple Crabapple Loquat Mayhaw Pear Pear, Asian Quince	European Red Mite McDaniel Spider Mite Pacific Spider Mite Two- spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	<p>Apply by ground with airblast equipment in a minimum of 100 gals/A. Coverage is essential for good control. Use of higher water volume will assure better coverage. Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide and should be used early in the life cycle of mites. Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>This product will not control Rust Mites or Blister Mites. If these pests are a problem, use an alternative miticide registered for that use.</p> <p>Do not use below use rate 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
<b>Restrictions</b>			
<ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 14 days.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 3.0 oz (0.135 lb. ai) of this product per acre per calendar year.</li> </ul>			

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state, and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

SMALL FRUIT VINE CLIMBING, EXCEPT FUZZY KIWIFRUIT (Subgroup 13-07 F)			
CROP	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Amur River Grape Gooseberry Grape Kiwifruit, Hardy Maypop Schisandra Berry Cultivars, varieties, and/or hybrids of these	European Red Mite Pacific Spider Mite Two- spotted Spider Mite Willamette Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	<p>Apply by ground as a full coverage spray in a minimum of 25 gals/A of water. Coverage is essential for good control. Use of higher water volume will assure better coverage. Best results are achieved when mite populations are low. This product is predominately an ovicide /larvicide and should be used early in the life cycle of mites. Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>This product will not control Rust Mites or Blister Mites. If these pests are a problem, use an alternative miticide registered for that use.</p> <p>Do not use below use rate of 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the Development of resistance to etoxazole among mite populations.</p>
<p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 14 days</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 3.0 oz (0.135 lb. ai) of this product per acre per calendar year.</li> </ul>			

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

STONE FRUIT (Crop Group 12)			
CROP	PEST	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Apricot Cherry (sweet and tart) Nectarine Peach Plum Plum, Chickasaw Plum, Damson Plum, Japanese Plumcot Prune (fresh)	European Red Mite Pacific Spider Mite Two- spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	Apply with ground equipment in a minimum of 50 gals/A of water. Coverage is essential for good control. Use of higher water volume will assure better coverage. Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide. Apply this product Miticide at or prior to threshold for your area but not greater than the maximum rate listed.  Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.  Do not use below use rate of 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.
<b>Restrictions</b> <ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 7 days.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 3.0 oz (0.135 lb. ai) of this product per acre per calendar year.</li> </ul>			

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

NON-BEARING FRUIT TREES			
CROP	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Non-Bearing Fruit Trees	European Red Mite McDaniel Spider Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	<p>Apply by ground with airblast equipment as a full coverage spray.</p> <p>Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide. Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>Do not use below use rate 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
	<p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>• Do not harvest fruit or nuts from treated trees within one (1) year of application.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 3.0 oz (0.135 lb. ai) this product per acre per calendar year.</li> </ul>		

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with other miticides having different modes of action and/or that belong to different chemical classes. Follow local, state and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

CHRISTMAS TREES			
CROP	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Christmas Trees	McDaniel Spider Mite Pacific Spider Mite Spruce Spider Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	<p>Apply by ground as full coverage spray or by air in a minimum of 10 gals/A. Coverage is critical for effective control.</p> <p>Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide. Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>Do not use below use rate 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
	<p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>Do not make more than one (1) this product application per season.</li> <li>Do not apply more than 3.0 oz (0.135 lb. ai) this product per acre per calendar year.</li> </ul>		

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state, and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

TREE NUTS			
CROPS	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Almond Beech Nut Brazil Nut Butternut Cashew Chestnut Chinquapin Filbert (Hazelnut) Hickory Nut Macadamia Nut Pecan Pistachio Walnut Black English	European Red Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	Apply by ground with airblast equipment as a full coverage spray. Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide. Apply this product at or prior to threshold for your area but not greater than the maximum rate listed.  Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.  Do not use below use rate of 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.
<b>Restrictions</b> <ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 28 days.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 3.0 oz (0.135 lb. ai) this product per acre per calendar year.</li> </ul>			

**MANAGING RESISTANCE:** Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state, and federal Integrated Pest Management (IPM) and Resistance Management (RM) recommendations. Read and follow all product labels before applying any miticide.

TROPICAL AND SUB-TROPICAL FRUIT (INEDIBLE PEEL)			
CROP	PESTS	PRODUCT RATE OZ/ACRE	INSTRUCTIONS
Avocado Canistel Mango Papaya Sapodilla Sapote, Black Sapote, Mamey Star Apple	Avocado Brown Mite Persea Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb. ai/A)	<p>Apply with [air* or] ground equipment in adequate water for uniform coverage ([minimum 20 gals/A by air* or] minimum 50 gals/A [by ground]). Applications of this product must be made using enough carrier (water) to ensure thorough coverage of the crop's vegetative and reproductive parts, which mites use for dispersal, feeding and reproduction. Use higher water volumes on more mature plants and varieties that have more compact and dense foliage. Higher volumes of water coupled with well calibrated equipment will help to provide better coverage.</p> <p>Best results are achieved when mite populations are low. This product is predominately an ovicide/larvicide. Apply this product Miticide at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Use the higher rates for moderate to heavy infestations and choose the lower rate for light infestations.</p> <p>Do not use below use rate of 2.0 oz/A (0.09 lb. ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p>
[*Not approved for aerial application in New York.]			
<p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>• Preharvest interval (PHI) is 1 day.</li> <li>• Do not make more than one (1) this product application per season.</li> <li>• Do not apply more than 3.0 oz (0.135 lb. ai) of this product per acre per calendar year.</li> </ul>			

MANAGING RESISTANCE: Repeated use of the same class of miticides or miticides with similar modes of action can lead to the buildup of resistant mite strains. Use this product in alternation with miticides having different modes of action and/or that belong to different chemical classes. Follow local, state, and federal Integrated Pest Management (IPM) and Resistance Management. Read and follow all product labels before applying any miticide.



**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

**PESTICIDE STORAGE**

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Do not store diluted spray.

**PESTICIDE DISPOSAL**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited.

**CONTAINER HANDLING**

Nonrefillable outer bag. Do not reuse or refill the outer bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration.

Batch Code: \_\_\_\_\_

**WARRANTY DISCLAIMER AND NOTICE**

---

**IMPORTANT: READ BEFORE USE**

---

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aceto Life Sciences, LLC. To the extent consistent with applicable law all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ACETO LIFE SCIENCES, LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aceto Life Sciences, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, ACETO LIFE SCIENCES, LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ACETO LIFE SCIENCES, LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

All trademarks are proprietary of their respective owners.

[Label for Container]

## Aceto Etoxazole 72% WSP Miticide

	% BY WT
<b>ACTIVE INGREDIENT:</b>	
Etoxazole* .....	72.0%
<b>OTHER INGREDIENTS:</b> .....	28.0%
Total	100.0%
*2-(2,6-difluorophenyl)-4-[4-(1,1-dimethylethyl)-2-ethoxyphenyl]-4,5-dihydrooxazole	

### KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>Emergency Assistance:</b> Have the product container or label with you when calling a poison control center or doctor or going for treatment. FOR EMERGENCY MEDICAL HELP, CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887.	

FOR CHEMICAL SPILL, LEAK, FIRE, EXPOSURE OR MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION.** Causes moderate eye irritation. Harmful if inhaled. Avoid contact with eyes, skin and clothing. Avoid Breathing dust.

See inside booklet for Precautionary Statements and Directions for Use.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

EPA Reg. No. 2749-XXX  
EPA Est. No.

Net Contents:

Manufactured for:  
Aceto Life Sciences, LLC  
4 Tri Harbor Court  
Port Washington, NY 11050

Batch Code:\_\_\_\_\_

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

**PESTICIDE STORAGE**

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Do not store diluted spray.

**PESTICIDE DISPOSAL**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited.

**CONTAINER HANDLING**

Nonrefillable outer bag. Do not reuse or refill the outer bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration.