



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

2749-648

Date of Issuance:

10/8/25

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

AG36424 C 1SL Herbicide

Name and Address of Registrant (include ZIP Code):

John F. Wright
Authorized Representative of Aceto Life Sciences, LLC, d/b/a Actylis
Product and Regulatory Associates, LLC
8595 Collier Blvd.
Naples, FL 34114

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Elizabeth Fertich, Product Manager 04
Invertebrate-Vertebrate Branch 1
Registration Division (7505T)

Date:

10/8/25

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 2749-648."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSFs:

- Basic CSF dated 09/18/2025
- Alternate CSF 1 dated 09/18/2025

If you have any questions, you may contact Robert Mitchell at 202-566-2842 or via email at mitchell.robert@epa.gov.

Enclosure

[Note to reviewer: [Text] in brackets denotes optional or explanatory language]
[Note to reviewer: {Text} in braces denotes where in the final label text will appear].
{BOOKLET FRONT PANEL LANGUAGE}

| | | | |
|----------|-------|---|-----------|
| IMAZAMOX | GROUP | 2 | HERBICIDE |
|----------|-------|---|-----------|

AG36424 C 1SL Herbicide

[For The Control of Vegetation in and Around Aquatic and Noncropland Sites] [Including
Areas That May Be Grazed or Cut for Hay]

ACTIVE INGREDIENT: (% by weight)

Ammonium salt of imazamox: 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-
-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic
acid*12.1%

OTHER INGREDIENTS:87.9%

TOTAL.....100.0%

*Equivalent to 11.4% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-methoxymethyl)-3- pyridinecarboxylic acid
1 gallon contains 1.0 pound of active ingredient as the free acid.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not
understand the label, find someone to explain it to you in detail.)

See inside label booklet for additional Precautionary Statements, and Directions for Use [including] [Storage
and Disposal] [Instructions].


| FIRST AID | |
|---|---|
| If on skin or clothing: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice. |
| If in eyes: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after first 5 minutes; then continue rinsing.• Call a poison control center or doctor for treatment advice. |
| If inhaled: | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth to mouth if possible.• Call a poison control center or doctor for further treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or physician or going for treatment. FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887 (24 Hours per Day, 7 Days per Week). | |

EPA Reg. No.: 2749-XXX

EPA Est. No.:

Net Contents: [1 qt., 1, & 2.5 gal. and bulk]

Manufactured [for][by]
Aceto Life Sciences, L.L.C., d/b/a Actylis
4 Tri Harbor Court
Port Washington, NY 11050-4661



Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer and Notice at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887 (24 Hours per Day, 7 Days per Week).

{LANGUAGE INSIDE BOOKLET}

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants.
2. Chemical-resistant gloves including barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber (includes natural rubber blends and laminates) \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, or viton \geq 14 mils.
3. Shoes plus socks.

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washable exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide may be hazardous to plants outside the treated area. DO NOT apply to water except as specified in this label. DO NOT contaminate water when disposing of equipment washwater and rinsate.

Non-Target Organism Advisory Statement

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Groundwater Advisory

Imazamox has properties and characteristics associated with chemicals detected in groundwater. Imazamox may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of [name of chemical] from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT mix or allow contact with oxidizing agents, as hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application.

USE RESTRICTIONS:

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Ensure spray drift to nontarget susceptible species does not occur.

DO NOT apply AG36424 C 1SL Herbicide in any manner not specifically described in this label.

Observe all cautions and limitations on this label and on the labels of products used in combination with AG36424 C 1SL Herbicide. DO NOT use AG36424 C 1SL Herbicide other than in accordance with the instructions set forth on this label. Keep containers closed to avoid spills and contamination.

PRODUCT INFORMATION

AG36424 C 1SL Herbicide is an aqueous formulation that may be diluted in water and either applied directly to water for the control/suppression of certain submerged aquatic vegetation or applied as a broadcast or spot spray to floating and emergent vegetation. Aquatic sites that may be treated include estuarine and marine sites, ponds, lakes, reservoirs, wetlands, marshes, swamps, bayous, arroyos, ditches, canals, streams, rivers, creeks and other slow-moving or quiescent bodies of water. AG36424 C 1SL Herbicide may also be used during drawdown conditions. AG36424 C 1SL Herbicide may also be applied for terrestrial and riparian vegetation control in industrial noncropland sites, and railroad, utility, and highway rights-of-way. Industrial noncropland sites include utility plant sites, tank farms, pumping installations, storage areas, fence rows and ditch banks. AG36424 C 1SL Herbicide may also be used for establishment and maintenance of wildlife openings. AG36424 C 1SL Herbicide may also be used on those sites listed above that may be grazed or cut for hay.

AG36424 C 1SL Herbicide is quickly absorbed by foliage and/or plant roots and rapidly translocated to the growing points stopping growth. Susceptible plants may develop a yellow appearance or general discoloration and will eventually die or be severely growth-inhibited.

AG36424 C 1SL Herbicide is herbicidally active on many submerged, emergent and floating broadleaf and monocot aquatic plants. The relative levels of control and selectivity can be manipulated by using a choice of rates and herbicide placement (water-injected or floating/emergent foliar application).

To help maintain the utility of herbicide programs, the use of herbicides with different modes of action is effective in managing weed resistance.

WEED RESISTANCE MANAGEMENT

For resistance management, AG36424 C 1SL Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to AG36424 C 1SL Herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of AG36424 C 1SL Herbicide or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected areas by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicides with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed management recommendations for specific crops and weed biotypes.

Report any incidence of non-performance of this product against a particular weed species to your Aceto Life Science, L.L.C., d/b/a Actylis retailer, representative or call 516-627-6000 (M-F 9:00 AM to 4:30 PM). If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemicals means to remove escapees, as practical, with the goal of preventing further seed production.

SPRAY ADJUVANTS

Applications of AG36424 C 1SL Herbicide to emergent, floating or shoreline species require the use of a spray adjuvant. Always use a spray adjuvant that is appropriate for aquatic sites.

Nonionic Surfactant

Use a nonionic surfactant at 0.25% volume/volume (v/v) or higher (see manufacturer's label) of the spray solution (0.25% v/v is equivalent to 1 quart in 100 gallons). For best results, select a nonionic surfactant with an HLB (hydrophilic to lipophilic balance) ratio between 12 and 17 with at least 70% surfactant in the formulated product (alcohols, fatty acids, oils, ethylene glycol or diethylene glycol should not be considered as surfactants to meet the above requirements).

Methylated Seed Oils or Vegetable Oil Concentrates

Instead of a surfactant, a methylated seed oil or vegetable-based seed oil concentrate may be used at 1.5 to 2 pints per acre. When using spray volumes greater than 30 gallons per acre, mix methylated seed oil or vegetable-based seed oil concentrates at 1 % of the total spray volume, or alternatively use a nonionic surfactant as described above. Research indicates that these oils may aid in AG36424 C 1SL Herbicide deposition and uptake by plants under stress.

Silicone-based Surfactants

See manufacturer's label for specific rate directions. Silicone-based Surfactants may reduce the surface tension of the spray droplet allowing greater spreading on the leaf surface as compared to conventional nonionic surfactants. However, some silicone-based surfactants may dry too quickly, limiting herbicide uptake.

Invert Emulsion

AG36424 C 1SL Herbicide can be applied as an invert emulsion. The spray solution results in an invert (water-in-oil) spray emulsion designed to minimize spray drift and spray runoff, resulting in more herbicide on the target foliage. The spray emulsion may be formed in a single tank (batch mixing) or injected (in-line mixing). Consult the invert chemical label for proper mixing directions.

Other

An antifoaming agent, spray pattern indicator, sinking agent or drift-reducing agent may be applied at the product labeled rate if necessary or desired.

SPRAY DRIFT

Aerial Applications

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

Ground Boom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size- Ground Boom:

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a

- higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size- Aircraft:

- **Adjust Nozzles** - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT- Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

AQUATIC USE DIRECTIONS

AG36424 C 1SL herbicide may be applied directly to the water for the control of submerged aquatic plant species and some emergent and floating species, or as a foliar application specifically for emergent and floating species.

AG36424 C 1SL Herbicide may be applied by surface and aerial equipment including both fixed-wing aircraft and helicopter.

USE RESTRICTIONS:

- DO NOT apply AG36424 C 1SL Herbicide to achieve a total active ingredient concentration in the water greater than 500 ppb (173 fl. oz.; 1.35 lb. ae*) per surface acre.
- For aquatic applications, DO NOT apply more than 500 ppb (173 fl. oz.; 1.35 lb. ae) per surface acre.
- For aquatic applications, DO NOT apply more than 48.66 gals (48.66 lb. ae) per surface acre per year.
- For aquatic applications, DO NOT apply more than 36 applications per year.
- For foliar broadcast applications, DO NOT apply more than 1 gallon (1 lb. ae) of AG36424 C 1SL Herbicide per surface acre per application for the control of emergent and floating vegetation.
- For surface applications, DO NOT apply more than 36 gals (36 lb. ae) per surface acre per year.

- For surface applications, DO NOT apply more than 36 applications per year.
- For foliar spot applications, DO NOT apply more than 5% v/v (0.05 lb. ae/gal) of AG36424 C 1SL Herbicide per surface acre per application. DO NOT apply more than 1 lb. ae per surface acre.
- The retreatment interval is 10 days.

*ae = acid equivalent

Foliar Application

Targeted Emergent and/or Floating Vegetation Application.

To make surface applications targeting emergent or floating vegetation, uniformly apply with properly calibrated broadcast or spot treatment equipment in 10 or more gallons of water per surface acre. Spot treatments can be made with up to 5% (0.05 lb. ae/surface acre) AG36424 C 1SL Herbicide by volume. To ensure through spray coverage, higher spray volumes may be required when treating areas with large and/or dense vegetation. Use an appropriate spray pressure to minimize the drift potential depending upon spray equipment, conditions and application objectives.

Foliar Treatment of Emergent and Floating Vegetation Guidelines

- Always use a surfactant for foliar applications of emergent and floating weeds.
- Foliar applications of AG36424 C 1SL Herbicide may be made as a broadcast spray or as a spot spray, with a percent spray solution ranging from 0.25% to 5% (0.0025-0.05 lb. ae/gal; maximum 1lb. ae/surface acre) AG36424 C 1SL Herbicide by volume
- Control will be reduced if spray is washed off foliage by wave actions.

In aquatic sites, those application techniques described in the Terrestrial Use Directions section may be used to treat emergent vegetation.

Application to Water

Water Application to Target Submerged and/or Emergent/Floating Vegetation

AG36424 C 1SL Herbicide may be broadcast-applied to the water surface or injected below the water surface. AG36424 C 1SL Herbicide may be applied as undiluted product or diluted with water prior to application. Under surface-matted conditions, inject AG36424 C 1SL Herbicide below the water surface to achieve better product distribution.

Apply AG36424 C 1SL Herbicide to water to achieve a final concentration of the active ingredient of no more than 500 ppb (173 fl. oz. ; 1.35 lb. ae) per surface acre. Multiple applications of AG36424 C 1SL Herbicide may be made during the annual growth cycle to maintain the desired vegetation response

AG36424 C 1SL Herbicide Rates per Treated Surface Acre

| Average Water Depth of Treatment Site (feet) | Desired Active Ingredient Concentration (ppb)* | | | |
|--|--|-----|-----|------|
| | 50 | 100 | 200 | 500 |
| | AG36424 C 1SL Herbicide Rate per Treated Surface Acre (fl. oz.) | | | |
| 1 | 17 | 35 | 69 | 173 |
| 2 | 35 | 69 | 138 | 346 |
| 3 | 52 | 104 | 207 | 518 |
| 4 | 70 | 138 | 277 | 691 |
| 5 | 87 | 173 | 346 | 864 |
| 6 | 104 | 207 | 415 | 1037 |
| 7 | 122 | 242 | 484 | 1210 |
| 8 | 139 | 277 | 553 | 1382 |
| 9 | 157 | 311 | 622 | 1555 |
| 10 | 174 | 346 | 691 | 1728 |
| * AG36424 C 1SL Herbicide contains 1.0 pound of active ingredient per gallon. There are 128 fl. oz. in one gallon. | | | | |

Aerial Application

AG36424 C 1SL Herbicide may be applied by both fixed-wing aircraft and helicopter. There is no minimum sprat volume when making applications directly to the water. For applications targeting emergent and/or floating vegetation, uniformly apply with properly calibrated equipment in 5 or more gallons of water per surface acre. For best results, make aerial applications using a minimum of 20 gallons per surface acre.

Drawdown Application

AG36424 C 1SL Herbicide may be used in drawdown situations to provide postemergence and/or preemergence control/suppression of aquatic vegetation. Apply AG36424 C 1SL Herbicide as a broadcast spray at rates up to 1 gallon/A (1 lb. ae/A), or as a spot spray treatment with up to 5% (0.05 lb. ae/gal; maximum 1 lb. ae/surface acre) AG36424 C 1SL Herbicide by volume. Make applications when water has receded and exposed soil is moist to dry. For postemergence (foliar) applications, wait at least two weeks after application before reintroducing water. When treating irrigation canals, the initial flush of recharge water after application must not be used for irrigation purposes.

Irrigation Restrictions

- DO NOT use treated water to irrigate greenhouses, nurseries, or hydroponics until the imazamox concentration has been determined by an acceptable method to be less than or equal to 1.0 ppb.
- DO NOT plant sugar beets, onions, potatoes or non-Clearfield®* canola in soils that have been previously irrigated with AG36424 C 1SL Herbicide treated water until a soil bioassay successfully demonstrates acceptable levels of crop safety.
- DO NOT use AG36424 C 1SL Herbicide treated waters resulting in a concentration greater than 50 ppb for irrigation until residue levels have been shown to be, less than or equal to 50 ppb by an acceptable method.
- DO NOT make applications of this product in and around golf course irrigation, sod farm irrigation, and vineyard irrigation waterbodies without testing potential irrigation water prior to irrigation and confirming the imazamox concentration to be less than or equal to 1.0 ppb.
- In still or quiescent waters, do not use AG36424 C 1SL Herbicide treated water resulting in a concentration greater than 10 ppb for irrigation of newly seeded or newly established plants until residue levels have been shown to be less than or equal to 10 ppb by an acceptable method.
- Wait 24 hours before irrigating from still or quiescent waters after making a AG36424 C 1SL Herbicide application for submerged vegetation less than 100 feet from an irrigation intake.
- Wait 24 hours before irrigating from still and quiescent waters after making a AG36424 C 1SL Herbicide application to emergent and/or floating vegetation if greater than 25% of the surface area of the water body has been treated or application was made less than 100 feet from an irrigation intake.
- Flowing waters may be used to irrigate allowable sites with no restrictions when AG36424 C 1SL Herbicide is applied at less than or equal to 2 quarts (0.5 lb. ae) per surface acre to waters with an average depth of greater than or equal to 4 feet.
- After application of AG36424 C 1SL Herbicide to dry irrigation canals/ditches, the initial flush of water during recharge must not be used for irrigation purposes unless the imazamox concentration has been determined by an acceptable method to be less than 25 ppb.

AG36424 C 1SL Herbicide applied at less than or equal to 2 quarts (0.5 lb. ae) per surface acre in or on waters with a minimum average depth greater than or equal to 4 feet will result in AG36424 C 1SL Herbicide concentrations less than 25 ppb.

* Clearfield is a registered trademark of BASF.

Other Water Use Restrictions

There are no restrictions on livestock watering, swimming, fishing, domestic use, or use of treated water for agricultural sprays.

Potable Water

AG36424 C 1SL Herbicide may be applied to potable water sources at concentrations up to 500 ppb (173 fl. oz.; 1.35 lb. ae) per surface acre to within a distance of 1/4 mile from an active potable water intake. Within 1/4 mile of an active potable water intake, AG36424 C 1SL Herbicide may be applied, but water concentrations resulting from injection and/or foliar

applications may not exceed 50 ppb. If water concentrations greater than 50 ppb are required, the potable water intake must be shut and, if necessary, an alternate water supply be made available until the water concentration can be shown to be less than 50 ppb by an acceptable method.

Endangered Plant Species

To prevent potential negative impacts to endangered plant species, DO NOT apply AG36424 C 1SL Herbicide in a way that adversely affects federally listed endangered and threatened species.

Weeds Controlled or Suppressed by AG36424 C 1SL Herbicide

Efficacy and selectivity of AG36424 C 1SL Herbicide is dependent upon many factors including: dose, time of year, stage of plant growth, plant susceptibility, method of application, and water movement. Rate selection will be partially dependent on characteristics of the treatment area and whether growth regulation or control is desired. Some areas may require a repeat application to control or suppress regrowth. Consult Aceto Life Sciences, L.L.C., d/b/a Actylis to determine the best treatment protocols to manage individual species and to meet specific aquatic plant management objectives.

Emergent, Floating, and Shoreline Species Controlled with Foliar Application

| Common Name | Scientific Name | Rate fl. oz./surface A (lb. ae/surface acre) | Comments |
|--------------------|------------------------------------|--|---|
| Alligatorweed | <i>Alternanthera philoxeroides</i> | 64 to 128 (0.5-1) | Repeat applications may be necessary. Add an aquatic glyphosate Herbicide for quicker brownout. See tank mix partner label for rates*. |
| American lotus | <i>Nelumbo lutea</i> | 64 to 128 (0.5-1) | |
| Arrowhead | <i>Sagittaria</i> spp. | 32 to 64 (0.25-0.5) | |
| Cattail** | <i>Typha</i> spp. | 32 to 64 (0.25-0.5) | Apply after full greenup through killing frost. |
| Chinese tallowtree | <i>Sapium sebiferum</i> | 64 to 128 (0.5-1) | |
| Common reed | <i>Phragmites</i> spp. | 96 to 128 (0.75-1) | Use 1 qt/A methylated seed oil (MSO); apply in late vegetative stage up to killing frost. Also apply as a spot treatment using 1 % to 2% (0.01-0.02 lb. ae/gal; maximum 1 lb. ae/surface acre) AG36424 C 1SL Herbicide per spray volume. Older stands of phragmites and stands growing in water may be more difficult to control and will require follow-up applications. |
| Common salvinia | <i>Salvinia minima</i> | 32 to 64 (0.25-0.5) | Apply with MSO or MSO + silicone-based surfactant; retreatment will be necessary. |
| Floating heart | <i>Nymphoides</i> spp. | 64 to 128 (0.5-1) | Also apply as a spot treatment using 2% to 5% (0.02-0.05 lb. ae/gal; maximum 1 lb. ae/surface acre) AG36424 C 1SL Herbicide and 1% MSO per spray volume. |
| Floating pennywort | <i>Hydrocotyle ranunculoides</i> | 32 to 64 (0.25-0.5) | Repeat applications may be necessary. |
| Flowering rush | <i>Butomus umbellatus</i> | 64 to 128 (0.5-1) | |
| Four-leaf clover | <i>Marsilea</i> spp. | 32 to 64 (0.25-0.5) | |
| Frog's bit | <i>Lymnobia spongia</i> | 16 to 32 (0.125-0.25) | |

| | | | |
|--|--|--------------------------|---|
| Giant cane | <i>Arundo donax</i> | 64 to 128 (0.5-1) | |
| Japanese knotweed | <i>Polygonum cuspidatum</i> | 64 to 128 (0.5-1) | |
| Mexican lily | <i>Nymphaea mexicana</i> | 32 to 64 (0.25-0.5) | |
| Mosquito fern | <i>Azolla</i> spp. | -- | Apply using 2% to 5% (0.02-0.05 lb. ae/gal; maximum 1 lb. ae/surface acre) AG36424 C 1SL Herbicide and 1% MSO by volume. |
| Parrotfeather | <i>Myriophyllum aquaticum</i> | 64 to 128 (0.5-1) | Apply only to emergent vegetation. |
| Pickeralweed | <i>Pontederia cordata</i> | 32 to 64 (0.25-0.5) | |
| Saltcedar | <i>Tamarix</i> spp. | 64 to 128 (0.5-1) | Also apply using 2% to 5% (0.02-0.05 lb. ae/gal; maximum 1 lb. ae/surface acre) AG36424 C 1SL Herbicide and 1 %. MSO per spray volume. |
| Smartweed, ladythumb Pennsylvania swamp | <i>Polygonum persicaria pennsylvanicum coccineum</i> | 64 to 128 (0.5-1) | |
| Spatdock | <i>Nuphar lutea</i> | 64 to 128 (0.5-1) | |
| Umbrella plant** | <i>Cyperus involucratus</i> | 64 (0.5) | Apply with MSO or COC. Also apply as a spot treatment using 5% (0.05 lb. ae/gal; maximum 1 lb. ae/surface acre) AG36424 C 1SL Herbicide per spray volume. |
| Variable-leaf milfoil | <i>Myriophyllum heterophyllum</i> | 64 to 128 (0.5-1) | Apply with MSO (1 % v/v) as an emergent foliar treatment when plants have emerged on the surface. Also apply as a spot treatment using 1% to 3% (0.01-0.03 lb. ae/gal; maximum 1 lb. ae/surface acre) AG36424 C 1SL Herbicide per spray volume. |
| Water chestnut | <i>Trapa natans</i> | 64 to 128 (0.5-1) | Apply with MSO to emergent part of plant. Also apply as a spot treatment using 2% to 5% (0.02-0.05 lb. ae/gal; maximum 1 lb. ae/surface acre) AG36424 C 1SL Herbicide per spray volume. |
| Water hyacinth | <i>Eichhornia crassipes</i> | 16 to 32 (0.125-0.25) | |
| Water lettuce | <i>Pistia stratiotes</i> | 48 to 96 (0.375-0.75) | |
| Water lily | <i>Nymphaea</i> spp. | 32 to 64 (0.25-0.5) | |
| Water primrose | <i>Ludwigia</i> spp. | 32 to 64 (0.25-0.5) | Add an aquatic glyphosate herbicide for quicker brownout. See tank mix partner label for rates*. |
| Watershield | <i>Brasenia schreberi</i> | 48 to 64 (0.375-0.5) | |
| Wild taro | <i>Colocasia esculenta</i> | 96 to 128 (0.75-1) | |

*It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

** Not Registered for Use by California.

Species Susceptible to Water-injected Applications

The following categories are provided to define species' that may be growth regulated or controlled with 50 to 500ppb (17-173 fl. oz.; 0.133-1.35 lb. ae) per surface acre AG36424 C 1SL Herbicide following in-water applications: susceptible, moderately susceptible, and less susceptible. The rates associated with each susceptibility category, including the Special Weed Control section, are provided as guidance with the overriding allowance that an application rate from 50 to 500 ppb (17-173 fl. oz.; 0.133-1.35 lb. ae) per surface may be used depending on the aquatic vegetation management objective and the characteristics of the aquatic vegetation and water body being treated.

Some species that are susceptible to foliar applications of AG36424 C 1SL Herbicide may be less susceptible to in-water applications. Use of higher rates are necessary to achieve desired control/suppression in areas of greater water exchange; when treating more mature or less susceptible plants; when targeting more difficult-to-control aquatic species; and when treating small areas in larger bodies of water (partial or spot treatments). Lower concentrations are generally used when conducting early season large-scale treatments; when greater selectivity is desired; and treating larger areas, more immature or susceptible plants, and areas with less potential for rapid water exchange.

Use of lower rates may increase selectivity on some species within the same category. Effects on susceptible plants can range from control to growth regulation depending on treatment site characteristics, exposure time, and application rate. Susceptible plant species may exhibit herbicide stress or reduced growth during active treatment phases. Whole lake applications with lower rates may provide plant growth regulation or greater selectivity while higher rates will generally provide broader activity.

Susceptible Vascular Aquatic Plant (50 to 200ppb; 17-69 fl. oz.; 0.133-0.539 lb. ae/surface acre)

| Common Name | Scientific Name |
|-----------------------|------------------------------|
| Curlyleaf pondweed | <i>Potamogeton crispus</i> |
| Eurasian watermilfoil | <i>Myriophyllum spicatum</i> |
| Hydrilla | <i>Hydrilla verticillata</i> |
| Sago pondweed | <i>Stuckenia pectinata</i> |
| Water hyacinth | <i>Eichhornia crassipes</i> |
| Water stargrass | <i>Heteranthera dubia</i> |

Moderately Susceptible Vascular Aquatic Plants (100 to 300 ppb; 35-103.8 fl. oz.; 0.273-0.811 lb. ae/surface acre)

| Common Name | Scientific Name |
|-----------------------|-----------------------------------|
| American pondweed | <i>Potamogeton nodosus</i> |
| Bladderwort | <i>Utricularia spp.</i> |
| Frog's bit | <i>Lymnobia spongia</i> |
| Illinois pondweed | <i>Potamogeton illinoensis</i> |
| Pickrelweed | <i>Pontederia cordata</i> |
| Salvinia | <i>Salvinia spp.</i> |
| Spikerush | <i>Eleocharis baldwinii</i> |
| Variable-leaf milfoil | <i>Myriophyllum heterophyllum</i> |
| Wigeon grass | <i>Ruppia maritima</i> |

Less Susceptible Vascular Aquatic Plants (200 to 500 ppb; 69-173 fl. oz.; 0.539-1.35 lb. ae/surface acre)

| Common Name | Scientific Name |
|--------------------|------------------------------------|
| Bulrush | <i>Schoenoplectus californicus</i> |
| Cattail* | <i>Typha spp.</i> |
| Coontail | <i>Ceratophyllum demersum</i> |
| Eelgrass, Japanese | <i>Zostera japonica</i> |
| Egeria | <i>Egeria densa</i> |
| Flowering rush | <i>Butomus umbellatus</i> |
| Southern naiad | <i>Najas guadalupensis</i> |
| Spatterdock | <i>Nuphar lutea</i> |
| Water lily | <i>Nymphaea odorata</i> |
| Watershield | <i>Brasenia schreberi</i> |

* Not Registered for Use by California.

Eurasian Watermilfoil. Apply AG36424 C 1SL Herbicide at 100 to 200 ppb (35-69 fl. oz.; 0.273-0.539 lb. ae) per surface acre to actively growing plants early in the growing season. Applications made to mature Eurasian watermilfoil (vegetation topped out) may require multiple applications.

Hydrilla. Apply AG36424 C 1SL Herbicide at 150 to 200 ppb (51.9-69 fl. oz.; 0.405-0.539 lb. ae) per surface acre to actively growing plants early in the growing season. Applications made prior to topped-out hydrilla may require repeat application. A single application of 50 to 75 ppb (17-25.95 fl. oz.; 0.133-0.203 lb. ae) per surface acre can be used to suppress and growth-regulate hydrilla for up to 10 to 12 weeks. If desired, an additional 50 to 75 ppb (17-25.95 fl. oz.; 0.133-0.203 lb. ae) per surface acre can be applied to extend the period of growth suppression when normal hydrilla growth resumes.

Japanese Eelgrass. Japanese eelgrass is a submerged aquatic plant which can be found in tidal and intertidal areas. AG36424 C 1SL Herbicide may be applied directly to the water or directly to the plant (e.g. at low tide).

TERRESTRIAL USE DIRECTIONS

USE RESTRICTIONS:

- The maximum amount of this product that can be applied is 1 gallon (equivalent to 1 pound of active ingredient as the free acid) per acre per year.
- DO NOT exceed 1 gallon (128 fl. oz.; 1lb.ae/A*) of product per acre per application.
- DO NOT exceed 2 applications of this product per year when using reduced application rates.
- The retreatment interval is 7 days.

*ae = acid equivalent

AG36424 C 1SL Herbicide may be applied with ground and aerial equipment including both fixed-wing aircraft and helicopter. Applications may be made using foliar broadcast spray, foliar spot spray, injection (hack and squirt), frill and girdle, cut stump, or basal methods.

Broadcast Spray Application

DO NOT apply more than 1 gallon (1 lb.ae/A) of AG36424 C 1SL Herbicide per acre per year.

Foliar Spot Application

Apply AG36424 C 1SL Herbicide as a percent solution, containing up to 5% AG36424 C 1SL Herbicide by volume and not to exceed 1 gallon (1 lb.ae/A).

Injection (Hack and Squirt), Frill and Girdle, and Cut Stump Application

Treatments may be made using up to 100% AG36424 C 1SL Herbicide by volume.

Basal Application

Treatments can be made using up to 25% AG36424 C 1SL Herbicide by volume. Basal applications require the use of a good emulsion system to maintain AG36424 C 1SL Herbicide in a stable emulsion with the penetrating agent being used. All foliar applications of AG36424 C 1SL Herbicide require the use of a spray adjuvant. Refer to Spray Adjuvants section for additional information.

To the extent consistent with the applicable law, the applicator is responsible for any loss or damage which results from spraying AG36424 C 1SL Herbicide in a manner other than directed in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

AG36424 C 1SL Herbicide may be used for the control of the following plant species. AG36424 C 1SL Herbicide may be effective for the control or suppression of additional plant species not listed below. The use of AG36424 C 1SL Herbicide for the control or suppression of undesirable plants not listed below may be done at the discretion of the user.

To the extent consistent with applicable law, the user assumes responsibility for any lack of control or suppression associated with application to weeds not listed on this label.

Weeds Controlled

| Common Name | Scientific Name | Rate Foliar fl. oz./A (lb. ae/A) | Comments |
|--|---|--|---|
| Alligator weed | <i>Alternanthera philoxeroides</i> | 64 to 128 (0.5-1) | Add an aquatic glyphosate herbicide for quicker brownout. See tank mix partner label for rates.** |
| Annual ryegrass | <i>Lolium multiflorum</i> | 16 to 32 (0.125-0.25) | |
| Artichoke, Jerusalem | <i>Helianthus tuberosus</i> | 64 to 128 (0.5-1) | |
| Bedstraw | <i>Galium aparine</i> | 64 to 128 (0.5-1) | |
| Beet, wild | <i>Beta procumbens</i> | 64 to 128 (0.5-1) | |
| Brazilian pepper[***] Christmasberry[***] | <i>Schinus terebinthifolius</i> | 96 to 128 (0.75-1) | Also apply using 2% to 5% (0.02-0.05 lb. ae/gal; maximum 1 lb.ae/A) AG36424 C 1SL Herbicide per spray volume. |
| Buckwheat, wild | <i>Polygonum convolvulus</i> | 64 to 128 (0.5-1) | |
| Buttercup | <i>Ranunculus spp.</i> | 64 to 128 (0.5-1) | |
| California bulrush[***] | <i>Schoenoplectus californicus</i> | 64 to 128 (0.5-1) | |
| Camphor tree[***] | <i>Cinnamomum camphora</i> | 2% to 5% v/v (0.02-0.05 lb. ae/gal; maximum 1 lb.ae/A) | |
| Canola, volunteer (non-Clearfield®) | <i>Brassica campestris</i> <i>Brassica napus</i> | 64 to 128 (0.5-1) | |
| Cattail*** | <i>Typha spp.</i> | 32 to 64 (0.25-0.5) | |

| Common Name | Scientific Name | Rate Foliar fl. oz./A (lb. ae/A) | Comments |
|--------------------------------------|---|--|--|
| Chickweed, common | <i>Stellaria media</i> | 64 to 128 (0.5-1) | |
| Chinese tallowtree Popcorn tree | <i>Sapium sebiferum</i> | 64 to 128 (0.5-1) | See Special Weed Control section. |
| Cocklebur, common | <i>Xanthium strumarium</i> | 64 to 128 (0.5-1) | |
| Filaree, redstem whitestem | <i>Erodium cicutarium</i> <i>Erodium moschatum</i> | 64 to 128 (0.5-1) | |
| Flixweed | <i>Descurainia sophia</i> | 64 to 128 (0.5-1) | |
| Frog's bit, Sponge plant | <i>Lymnobia</i> spp. | 16 to 32 (0.125-0.25) | |
| Giant ragweed* | <i>Ambrosia trifida</i> | 32 to 64 (0.25-0.5) | |
| Henbit | <i>Lamium amplexicaule</i> | 64 to 128 (0.5-1) | |
| Jamaican nightshade[***] | <i>Solanum jamaicense</i> | 2% to 5% v/v (0.02-0.05 lb. ae/gal; maximum 1 lb.ae/A) | |
| Japanese stiltgrass | <i>Microstegium vimineum</i> | 32 to 64 (0.25-0.5) | Use MSO at 1 % by spray volume. AG36424 C 1SL Herbicide will provide some residual control of subsequent seedling emergence. |
| Jimsonweed | <i>Datura stramonium</i> | 64 to 128 (0.5-1) | |
| Johnsongrass, rhizome seedling | <i>Sorghum halepense</i> | 32 to 64 (0.25-0.5) 16 to 32 (0.125-0.25) | |
| Knotweed, prostrate | <i>Polygonum aviculare</i> | 64 to 128 (0.5-1) | |
| Kochia | <i>Kochia scoparia</i> | 64 to 128 (0.5-1) | |
| Lambsquarters, common | <i>Chenopodium album</i> | 64 to 128 (0.5-1) | |
| Lettuce, miner's | <i>Montia perfoliata</i> | 64 to 128 (0.5-1) | |
| Mallow, common Venice | <i>Malva neglecta</i> <i>Hibiscus trionum</i> | 64 to 128 (0.5-1) | |
| Mustard spp. | <i>Brassica</i> spp. | 64 to 128 (0.5-1) | |
| Nettle, burning | <i>Urtica urens</i> | 64 to 128 (0.5-1) | |
| Nettleleaf goosefoot | <i>Chenopodium murale</i> | 64 to 128 (0.5-1) | |

| Common Name | Scientific Name | Rate Foliar fl. oz./A (lb. ae/A) | Comments |
|---|---|--|---|
| Nightshade, black Eastern black hairy | <i>Solanum nigrum ptycanthum sarrachoides</i> | 64 to 128 (0.5-1) | |
| Old World climbing fern[***] | <i>Lygodium microphyllum</i> | <u>5% v/v</u> | |
| Pennycress, field | <i>Thlaspi arvense</i> | 64 to 128 (0.5-1) | |
| Phragmites[***] | <i>Phragmites australis</i> | 64 to 128 (0.5-1) | Use 1 qt/A methylated seed oil (MSO); apply in late vegetative stage up to killing frost. Also apply as a spot treatment using 1 % to 2% (0.01-0.02 lb. ae/gal; maximum 1 lb.ae/A) AG36424 C 1SL Herbicide per spray volume. Older stands of phragmites and stands growing in water may be more difficult to control and will require follow-up applications. |
| Pigweed, prostrate redroot smooth spiny | <i>Amaranthus blitoides retroflexus hybridus spinosus</i> | 64 to 128 (0.5-1) | |
| Puncturevine | <i>Tribulus terrestris</i> | 64 to 128 (0.5-1) | |
| Purple loosestrife[***] | <i>Lythrum salicaria</i> | <u>32 to 64</u> <u>(0.25-0.5)</u> | |
| Purslane, common | <i>Portulaca oleracea</i> | 64 to 128 (0.5-1) | |
| Radish, wild | <i>Raphanus raphanistrum</i> | 64 to 128 (0.5-1) | |
| Ragweed, Common giant | <i>Ambrosia artemisiifolia trifida</i> | 64 to 128 (0.5-1) | |
| Rocket, London yellow | <i>Sisymbrium irio Barbarea vulgaris</i> | 64 to 128 (0.5-1) | |
| Saltcedar[***] | <i>Tamarix</i> spp. | 64 to 128 (0.5-1) | Also apply using 2% to 5% (0.02-0.05 lb. ae/gal; maximum 1 lb.ae/A) AG36424 C 1SL Herbicide and 1%MSO per spray volume. |
| Sedge[***], purple yellow | <i>Cyperus rotundus esculentus</i> | 32 to 64 (0.25-0.5) | Also apply using 2% to 5% (0.02-0.05 lb. ae/gal; maximum 1 lb.ae/A) AG36424 C 1SL Herbicide per spray volume. |
| Shepherd's-purse | <i>Capsella bursa- pastoris</i> | 64 to 128 (0.5-1) | |
| Smartweed, ladythumb Pennsylvania swamp | <i>Polygonum persicaria pensylvanicum coccineum</i> | 64 to 128 (0.5-1) | |

| Common Name | Scientific Name | Rate Foliar fl. oz./A (lb. ae/A) | Comments |
|-----------------------------------|-----------------------------------|---|---|
| Spike rush[***] | <i>Eleocharis</i> spp. | 64 to 128 (0.5-1) | |
| Spurge, prostrate | <i>Euphorbia maculata</i> | 64 to 128 (0.5-1) | |
| Sunflower, common | <i>Helianthus annuus</i> | 64 to 128 (0.5-1) | |
| Swinecress | <i>Coronopus didymus</i> | 64 to 128 (0.5-1) | |
| Tansymustard, green | <i>Descurainia pinnata</i> | 64 to 128 (0.5-1) | |
| Taro | <i>Taro</i> spp. | 64 to 128 (0.5-1) 5% v/v (0.05 lb. ae/A; maximum 1 lb.ae/A) | |
| Thistle, Russian | <i>Salsola iberica</i> | 64 to 128 (0.5-1) | |
| Tropical soda-apple[***] | <i>Solanum viarum</i> | <u>2% to-5%-v/v</u> <u>(0.02-0.05 lb. ae/gal;</u> <u>maximum 1 lb.ae/A)</u> | |
| Umbrella plant[***] | <i>Cyperus involucratus</i> | 64 (0.5-1) | Apply with MSO or COC. Also apply as a spot treatment using 5% (0.05 lb. ae/gal; maximum 1 lb.ae/A) AG36424 C 1SL Herbicide per spray volume. |
| Water primrose | <i>Ludwigia</i> spp. | 32 to 64 (0.25-0.5) | Add an aquatic glyphosate herbicide for quicker brownout. See tank mix partner label for rates.** |
| Wetland nightshade[***] | <i>Solanum tampicense</i> | 2% to 5% v/v(0.02- 0.05 lb. ae/gal; maximum 1 lb.ae/A) | |
| Whitetop[***] Hoary cress[***] | <i>Cardaria draba</i> | 8 to 16 (0.063-0.125) | |
| Willoweed panicle | <i>Epilobium brachycarpum</i> | 64 to 128 (0.5-1) | |
| Velvetleaf | <i>Abutilon theophrasti</i> | 64 to 128 (0.5-1) | |

* Suppression of larger, well-established plants

** It is the pesticide user's responsibility to ensure that all products are registered for the intended use.

[*** Not registered for use by California.]

Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

In general, the use of methylated seed oil (MSO) at 1 % v/v will provide the best control with foliar applications.

Special Weed Control- Chinese Tallowtree

AG36424 C 1SL Herbicide at 64 to 128 fl. oz./A (0.5-1 lb.ae/A) or 0.5 to 2.0% v/v (0.005-0.02 lb. ae/gal; maximum 1 lb.ae/A) may be applied as a foliar application for selective control of Chinese tallowtree in and around non-sensitive tree species.

Control Chinese tallowtree with foliar applications using aerial, handgun, or backpack application methods. When treating Chinese tallowtree, ensure that application method and spray volume provide adequate coverage of targeted Chinese tallowtree plants. Add methylated seed oil at 32 fl. oz./A (0.25lb.ae/A) for broadcast applications, or at 1% v/v (0.01 lb. ae/gal; maximum 1 lb.ae/a) for spot backpack and handgun applications. Non-sensitive hardwood species may exhibit varying degrees of leaf discoloration and temporary injury.

Areas that may be Grazed or Cut for Hay

Apply AG36424 C 1SL Herbicide to listed aquatic and terrestrial non-crop sites that may be grazed or cut for hay at a maximum use rate of 1 gallon (1 lb.ae/A) per acre of AG36424 C 1SL Herbicide or 5% (v/v) (0.05 lb. ae/gal; maximum 1 lb.ae/A) spray solution for spot treatments. There are no grazing or haying restrictions.

Storage and Disposal for nonrefillable rigid plastic containers (5 gal or less)

Storage and Disposal

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. Avoid freezing. Store above 40°F. If frozen, poor weed control may result. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(Storage and Disposal for refillable rigid plastic containers (larger than 5 gal)

Storage and Disposal

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. Avoid freezing. Store above 40°F. If frozen, poor weed control may result. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Offer for recycling, if available or recondition if appropriate.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

(Storage and Disposal for nonrefillable rigid plastic containers (larger than 5 gal)

Storage and Disposal

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. Avoid freezing. Store above 40°F. If frozen, poor weed control may result. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times.

Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

WARRANTY DISCLAIMER AND NOTICE

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranty and Limitations Liability before using this product. If the terms are not acceptable, return the unopened product at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use, all of which are beyond the control of Aceto Life Sciences, L.L.C.. d/b/a Actylis. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Aceto Life Sciences, L.L.C., d/b/a Actylis MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aceto Life Sciences, L.L.C., d/b/a Actylis is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Aceto Life Sciences, L.L.C.. d/b/a Actylis DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ACETO LIFE SCIENCES, L.L.C., D/B/A ACTYLIS'S ELECTION, THE REPLACEMENT OF PRODUCT.

All trademarks are the property of their respective owners.

[Made in [China][India][Malaysia]] [,] [formulated in [USA][China][India][Malaysia]] [&] [packaged in [USA][China][India][Malaysia]]

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

IMAZAMOX

GROUP

2

HERBICIDE

AG36424 C 1SL Herbicide

[For The Control of Vegetation in and Around Aquatic and Noncropland Sites] [Including Areas That May Be Grazed or Cut for Hay]

ACTIVE INGREDIENT:

(% by weight)

Ammonium salt of imazamox: 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid*12.1%

OTHER INGREDIENTS:87.9%

TOTAL.....100.0%

*Equivalent to 11.4% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-methoxymethyl)-3- pyridinecarboxylic acid
1 gallon contains 1.0 pound of active ingredient as the free acid.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

| FIRST AID | |
|---|---|
| If on skin or clothing: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice. |
| If in eyes: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after first 5 minutes; then continue rinsing.• Call a poison control center or doctor for treatment advice. |
| If inhaled: | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth to mouth if possible.• Call a poison control center or doctor for further treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or physician or going for treatment. FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887 (24 Hours per Day, 7 Days per Week). | |

In case of emergency endangering health or the environment involving this product, CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887 (24 Hours per Day, 7 Days per Week).

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide may be hazardous to plants outside the treated area. DO NOT apply to water except as specified in this label. DO NOT contaminate water when disposing of equipment washwater and rinsate.

Non-Target Organism Advisory Statement

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Groundwater Advisory

Imazamox has properties and characteristics associated with chemicals detected in groundwater. Imazamox may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of [name of chemical] from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT mix or allow contact with oxidizing agents, as hazardous chemical reaction may occur.

Storage and Disposal for nonrefillable rigid plastic containers (5 gal or less)

Storage and Disposal

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. Avoid freezing. Store above 40°F. If frozen, poor weed control may result. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(Storage and Disposal for refillable rigid plastic containers (larger than 5 gal))

Storage and Disposal

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. Avoid freezing. Store above 40°F. If frozen, poor weed control may result. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Offer for recycling, if available or recondition if appropriate.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides

while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

(Storage and Disposal for nonrefillable rigid plastic containers (larger than 5 gal))

Storage and Disposal

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. Avoid freezing. Store above 40°F. If frozen, poor weed control may result. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

See label booklet for additional Precautionary Statements and Directions for Use.

EPA Reg. No.: 2749-AUI

EPA Est. No.:

Net Contents: [1 qt., 1, & 2.5 gal. and bulk]

**Manufactured [for][by]
Aceto Life Sciences, L.L.C., d/b/a Actylis
4 Tri Harbor Court
Port Washington, NY 11050-4661**