



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

2749-654

Date of Issuance:

11/10/25

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

AG37593 CS EC Fungicide

Name and Address of Registrant (include ZIP Code):

Aceto Life Sciences, L.L.C. d/b/a Actylis
4 Tri Harbor Court
Port Washington, NY 11050

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Stephanie Suarez, Ph.D., Acting Product Manager 22
Fungicide Branch, Registration Division (7505T)

Date:

11/10/25

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 2749-654."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 10/11/2024
- Alternate CSFs 1-3 dated 10/11/2024

If you have any questions, please contact Yasmin Bowers at 202-566-2507 or Bowers.Yasmin@epa.gov.

Enclosure

ACCEPTED

11/10/2025

Under the Federal Insecticide,
Fungicide and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 2749-654

DIFENOCONAZOLE	GROUP	3	FUNGICIDE
PROPICONAZOLE	GROUP	3	FUNGICIDE

AG37593 CS EC

Fungicide

A broad-spectrum fungicide with systemic and curative properties registered for the control of key diseases in almonds; brassica, leafy greens; bulb vegetables; carrots; hazelnuts; pecans; pistachios; soybeans; stone fruits; sugar beets; and tree nuts.

ACTIVE INGREDIENTS:	(% by weight)
Difenoconazole (CAS No. 119446-68-3).....	22.8%
Propiconazole (CAS No. 60207-90-1).....	22.8%
OTHER INGREDIENTS*.....	54.4%
TOTAL.....	100.0%

* Contains petroleum distillates

AG37593 CS EC is formulated as an emulsifiable concentrate (EC) containing 2.08 lb. ai/gal each of difenoconazole and propiconazole

EPA Reg. No. 2749-ALU

EPA Est.

NET CONTENTS:

KEEP OUT OF REACH OF CHILDREN

DAÑERO/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand the label, find someone to explain it to you in detail)

[See] [inside] [label] [booklet] [for] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][.]

FIRST AID	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Immediately call a poison control center or doctor.• DO NOT induce vomiting unless told to do so by the poison control center or doctor.• DO NOT give any liquid to the person.• DO NOT give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage. Contains petroleum distillate. Vomiting may cause aspiration pneumonia.	
Have the product container or label with you when calling a poison control center or physician or going for treatment. FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887 (24 Hours per Day, 7 Days per Week).	

Manufactured [for][by]
Aceto Life Sciences, L.L.C. d/b/a Actylis
4 Tri Harbor Court
Port Washington, NY 11050-4661

PRECAUTIONARY STATEMENTS
Hazards to Human and Domestic Animals
DANGER/PELIGRO

Corrosive. Cause irreversible eye damage. Causes skin irritation. Harmful if swallowed. Harmful if absorbed through skin. **DO NOT** get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Mixers, loaders, applicators and all other handlers must wear:

- Protective eyewear (goggles, face shield, or safety glasses)
- Chemical resistant headgear for overhead exposure
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils or Viton[®] ≥ 14 mils.
- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant apron when cleaning equipment, mixing or loading
- Chemical-resistant footwear plus socks

USER SAFETY REQUIREMENTS

Discard clothing and other absorbed materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide get inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, shrimp, mammals, and aquatic invertebrates. Drift and runoff may be hazardous to aquatic estuarine/marine organisms in water adjacent to treated areas. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

SURFACE WATER ADVISORY

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product may contaminate water through drift or spray in wind. This product has a potential for runoff for several months or more after application. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water feature including ponds, streams, and springs will reduce the potential for contamination of water from runoff water and

sediment. Runoff of this product will be reduced by avoiding applications when rainfall is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming into contact with oxidizing agents, hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water is:

- Protective eyewear (goggles, face shield, or safety glasses)
- Chemical resistant headgear for overhead exposure
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils or Viton[®] ≥ 14 mils
- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant apron when cleaning equipment, mixing, or loading
- Chemical-resistant footwear plus socks

FAILURE TO FOLLOW DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

PRODUCT INFORMATION

AG37593 CS EC is a broad-spectrum fungicide with systemic and curative properties registered for the control of key diseases in almonds; berries; bushberry and low growing; bulb vegetables; carrots; hazelnuts; pecans; pistachios; soybeans; stone fruits; sugar beets; and tree nuts. **AG37593 CS EC** may be applied as a foliar spray in alternating spray programs or in tank mixes with other crop protection products. All applications must be made according to the use directions that follow.

PRODUCT USE INSTRUCTIONS

Application: Thorough coverage is necessary to provide good disease control. Make up no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

Efficacy: Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of AG37593 CS EC has been used. If resistant isolates to Group 3 fungicides are present, efficacy can be reduced. Under high disease pressure, use the highest rate and shortest interval.

Rotational Crops: Please see the table below for the crop rotational restrictions.

Rotational Crop	Planting Time From Last AG37593 CS EC Application
Almond Bean, Dried except cowpeas Berry, Bushberry Subgroup 13-07B Berry, Low Growing Subgroup 13-07G Brassica leafy greens, Subgroup 5B Bulb vegetables, bulb onion Subgroup 3-07A and green onion Subgroup 3-07B Carrot Filbert Pecan Pistachio Rice Soybean Stone fruit Crop Group 12-12 Strawberry Sugar beet Tree nuts Crop Group 14-12 Watercress Wild rice	0 days
Cereals (wheat, barley, triticale, oats, rye)	30 days
Cilantro Corn Corn, Sweet Garden Beet Parsley Peanut	60 days
All other crops intended for food and feed	105 days

Crop Tolerance: Plant tolerance has been found acceptable for all crops on the label; however, not all possible tank-mix combinations have been tested under all conditions. When possible, test the combinations on a small portion of the crop to ensure a phytotoxic response will not occur as a result of application.

MANDATORY SPRAY DRIFT

Aerial Applications

- **DO NOT** release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a medium to ultra coarse spray droplet size (ASABE S572.1).
- The boom length must not exceed 65% of wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.

- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Ground Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium to ultra coarse spray droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENT

- **IMPORTANCE OF DROPLET SIZE**
An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

- **BOOM HEIGHT – Ground Boom**
Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.
- **RELEASE HEIGHT – Aircraft**
Higher release heights increase the potential for spray drift. When applying aerially to crops, **DO NOT** release spray at a height greater than 10 feet above the crop canopy, unless a greater application height is necessary for pilot safety.
- **SHIELDED SPAYERS**
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- **TEMPERATURE AND HUMIDITY**
When making application in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- **TEMPERATURE INVERSIONS**
Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog

or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

- **WIND**
Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

IPM: AG37593 CS EC should be integrated into an overall disease and pest management strategy (IPM) whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. Consult your local agricultural authorities for additional IPM strategies established for your area. **AG37593 CS EC** may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

DIFENOCONAZOLE	GROUP	3	FUNGICIDE
PROPICONAZOLE	GROUP	3	FUNGICIDE

For resistance management, **AG37593 CS EC** contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to **AG37593 CS EC** and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps.

- Rotate the use **AG37593 CS EC** or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, use historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact [Aceto Life Science, L.L.C. d/b/a Actylis] at 516-627-6000. You can also contact your pesticide distributor or university extension specialist to report resistance.

SPRAY EQUIPMENT

Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Use the same size nozzles uniformly spaced across the boom.
- Calibrate sprayer before use.
- Use screens to protect the pump and to prevent nozzles from clogging.
- On suction side of pump use screens that are *16-mesh or coarser*.
- **DO NOT** place a screen in the recirculation line.

- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's directions.

Pump

- Use a pump with capacity to:
 - Maintain sufficient pressure at the nozzle tip to give the required flow rate and droplet size to provide acceptable coverage of the target crop.
 - Provide sufficient agitation in tank to keep mixture in suspension-this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- **DO NOT** air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and state specifications. For specific local directions and spray schedules, consult the current state agricultural directions.

MIXING INSTRUCTIONS

- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Alone (no tank mix):

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add **AG37593 CS EC** to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray after **AG37593 CS EC** has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

AG37593 CS EC + Tank Mixtures: **AG37593 CS EC** is usually compatible with tank-mix partners. To determine the physical compatibility of with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1qt. of water. Add wettable powders and water dispersible granular products first, then liquid flowables and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Adjuvants: When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Council of Producers and Distributors Agrotechnology (CPDA) adjuvant certification program is advised.

Mixing in the Spray Tank

- Add ½-2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and **AG37593 CS EC** to the spray tank.
- Allow **AG37593 CS EC** to completely disperse.
- Spray the mixture with the agitator running
- Observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix product label.
- Label dosage rate must not be exceeded, and the most restrictive label precautions and limitations must be followed.
- This product must not be mixed with any product which prohibits such mixing.

APPLICATION PROCEDURES

Ground Application:

- Apply in a minimum of 15 gal of water per acre, unless specified otherwise.

Aerial Application:

- Use only on crops where aerial application are indicated.
- Apply in a minimum of 5 gal of water per acre, unless specified otherwise.
- **DO NOT** apply under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur.
- **DO NOT** apply directly to humans or animals.

Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation system. **DO NOT** apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.125-0.25 inches/A of water. Excessive water may reduce efficacy.
- If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

RESTRICTIONS: DO NOT inject **AG37593 CS EC** at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1-part **AG37593 CS EC**. **AG37593 CS EC** is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used, but must be replaced once a year. **DO NOT** use Viton, Buna-N, Neoprene, or PVC seals.

Operating Instructions

1. The system must contain a functional check-valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick closing check-valve to prevent the flow of fluid back toward the injection pump.

3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended.

Center Pivot Irrigation Equipment

Notes: (1) Use only with drive systems which provide uniform water distribution. (2) **DO NOT** use end guns when chemigating **AG37593 CS EC** through center pivot systems because of non-uniform application.

- Determine the size of the area to be treated.
- Determine the time required to apply $\frac{1}{8}$ - $\frac{1}{2}$ inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as directed by the equipment manufacturer. When apply **AG37593 CS EC** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of **AG37593 CS EC** required to treat the area covered by the irrigation system.
- Add the required amount of **AG37593 CS EC** and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the **AG37593 CS EC** solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the **AG37593 CS EC** solution has cleared the sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30-minute interval. When applying **AG37593 CS EC** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of **AG37593 CS EC** required to treat the area covered by the irrigation system.
- Add the required amount of **AG37593 CS EC** into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the **AG37593 CS EC** solution has cleared the last sprinkler head.

SPECIFIC INSTRUCTIONS FOR PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-

pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system must be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

3. The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

SPECIFIC USE DIRECTIONS

Almonds[*]

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Alternaria leafspot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum. acutatum</i>) Blossom Blight Brown rot (<i>Monilinia</i> spp.) Brown rot/hull rot (<i>Monilinia</i> spp.) Powdery Mildew (<i>Podosphaera tridactyla</i> , <i>Sphaerotheca pannosa</i>) Scab (<i>Venturia carpophila</i>) Shot hole (<i>Wilsonmyces carpophilus</i>)	5-7 (0.08-0.11 lb. ai difenoconazole and propiconazole)	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply AG37593 CS EC on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p>If monitoring or history indicates the presence of Alternaria, apply 7 fl. oz./A of AG37593 CS EC in the late spring (mid-April to beginning of May) and then repeat the treatment 2-3 weeks later.</p> <p>For blossom blight, begin applications at early bloom and continue through petal fall. Minimum retreatment interval is 14 days.</p> <p>For anthracnose, begin applications at bud break on a 14-day interval.</p> <p>[Optional language if label has a rate range: If disease pressure is high, use the highest rate.] [Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</p>
Application: For best results, sufficient water volume must be used to provide thorough coverage. AG37593 CS EC can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For		

aerial applications, use a minimum of 10 gal/A of water.

RESTRICTIONS:

- **DO NOT** apply more than 28 fl. oz./A/year of **AG37593 CS EC** (which delivers 0.46 lb. ai/A/year of difenoconazole and 0.45 lb. ai/A/year of propiconazole).
- **DO NOT** exceed 4 applications per year at the highest rate (7 fl. oz./A) or 5 applications per year at the lowest rate (5 fl. oz./A).
- **DO NOT** apply more than 0.46 lb. ai/A/year of difenoconazole-containing products.
- **DO NOT** apply more than 0.9 lb. ai/A/year of propiconazole-containing products.
- **DO NOT** use an adjuvant.
- **DO NOT** graze livestock in treated area or cut treated cover crop for feed.
- **DO NOT** apply within 60 days of harvest (60-day PHI).
- Retreatment Interval (RTI) is 14 days.

[* Not Registered for use by California]

Brassica, Leafy Greens subgroup 4-16B[*]: Arugula; broccoli, Chinese; broccoli raab; cabbage, abyssinian; cabbage, Chinese, bok choy; cabbage, seakale; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; rocket, wild; shepherd's purse; turnip greens; watercress; cultivars, varieties, and hybrids of these commodities.

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Alternaria leafspot (<i>Alternaria spp.</i>)	6-7 (0.10-0.11 lb. ai difenoconazole and propiconazole)	Begin applications at first sign of disease. Repeat on a 7-10-day interval. Apply no more than two applications before switching to another fungicide with a different mode of action.
Anthrachnose (<i>Colletotrichum acutatum</i>)		If disease levels continue to increase, immediately switch to a fungicide with a different mode of action.
Black spot (<i>Alternaria spp.</i>)		
Cercospora leaf spot (<i>Cercospora spp.</i>)		[Optional language if label has a rate range: If disease pressure is high, use the highest rate.]
Powdery mildew (<i>Erysiphe polygoni</i>)		[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]
Ring spot (<i>Mycosphaella brassicicola</i>)		[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]

Application: **AG37593 CS EC** is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. **AG37593 CS EC** may be applied by ground, air, or chemigation.

RESTRICTIONS:

- **DO NOT** apply more than 21 fl. oz./A/year of **AG37593 CS EC** (which delivers 0.34 lb. ai/A/year of both difenoconazole and propiconazole).
- **DO NOT** exceed 3 applications per year.
- **DO NOT** apply more than 0.34 lb. ai propiconazole-containing product/A/year.
- **DO NOT** apply more than 0.46 lb. ai difenoconazole-containing product/A/year.
- **DO NOT** apply within 7 days of harvest (7-day PHI).
- Retreatment Interval (RTI) is 7 days.

[* Not Registered for use by California]

Garlic, Shallot, Bulb Vegetables (Onion bulb subgroup 3-07A)[*]: Daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these.

Leek, Welch Onion tops, Green Onion subgroup 3-07B[*]: Chive, fresh leaves; chive, Chinese, fresh leaves; elegans hosta; fritillaria, leaves; kurrat; lady's leek; leek; leek, wild; onion, Beltsville bunching; onion, fresh; onion, green; onion, macrostem; onion, tree, tops; onion, Welsh, tops; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Cercospora leafspot (<i>C. duddiae</i>) Cladosporium leaf blotch (<i>C. allii</i>) Leaf blotch (<i>Cladosporium alliicepae</i>) Powdery Mildew (<i>Leveillula taurica</i>) Purple blotch (<i>Alternaria porri</i>) Stemphylium leaf blight (<i>S. vesicarium</i>) Rust (<i>Puccinia allii</i>) Suppression of: Botrytis leaf blight (<i>B. squamosa</i>)	5-7 (0.08-0.11 lb. ai difenoconazole and propiconazole)	Begin applications prior to disease onset when conditions are conducive for disease. Apply AG37593 CS EC on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. [Optional language if label has a rate range: If disease pressure is high, use the highest rate.] [Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]
Application: For best results, sufficient water volume must be used to provide thorough coverage. AG37593 CS EC can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.		
RESTRICTIONS: <ul style="list-style-type: none"> For green onions, DO NOT apply more than 21 fl. oz./A/year of AG37593 CS EC (which delivers 0.34 lb. ai/A/year of both difenoconazole and propiconazole). For green onions, DO NOT exceed 3 applications per year at the highest rate (7 fl. oz./A) or 4 applications per year at the lowest rate (5 fl. oz./A). For green onions, DO NOT apply more than 0.34 lb. ai/A/year of difenoconazole-containing products. For dry bulb onions, DO NOT apply more than 28 fl. oz./A/year of AG37593 CS EC (which delivers 0.46 lb. ai/A/year of difenoconazole and 0.45 lb. ai/A/year of propiconazole). For dry bulb onions, DO NOT exceed 4 applications per year at the highest rate (7 fl. oz./A) or 5 applications per year at the lowest rate (5 fl. oz./A). For dry bulb onions, DO NOT apply more than 0.46 lb. ai/A/year of difenoconazole-containing products. For all Bulb crop group, DO NOT apply more than 0.45 lb. ai/A/year of propiconazole-containing products. DO NOT apply within 14 days of harvest for dry bulb type (14-day PHI). DO NOT apply within 7 days of harvest for green onions (7-day PHI). Retreatment Interval (RTI) is 7 days. <p>[* Not Registered for use by California]</p>		

Carrots[*]

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
<p>Alternaria leaf blight (<i>Alternaria dauci</i>)</p> <p>Cercospora leaf spot (<i>Cercospora carotae</i>)</p> <p>Powdery mildew (<i>Erysiphe spp.</i>)</p>	<p>5-7 (0.08-0.11 lb. ai difenoconazole and propiconazole)</p>	<p>Begin applications prior to disease onset when conditions are conducive for disease. Apply AG37593 CS EC on a 7- to 10-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p>[Optional language if label has a rate range: If disease pressure is high, use the highest rate.]</p> <p>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</p> <p>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. AG37593 CS EC can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>		
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> DO NOT apply more than 28 fl. oz./A/year of AG37593 CS EC (which delivers 0.46 lb. ai/A/year of difenoconazole and 0.45 lb. ai/A/year of propiconazole). DO NOT exceed 4 applications per year at the highest rate (7 fl. oz./A) or 5 applications per year at the lowest rate (5 fl. oz./A). DO NOT apply more than 0.46 lb. ai/A/year of difenoconazole-containing products. DO NOT apply more than 0.45 lb. ai/A/year of propiconazole-containing products. DO NOT apply within 14 days of harvest (14-day PHI). Retreatment Interval (RTI) is 14 days. 		
[* Not Registered for use by California]		

Cranberry[*]

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
<p>Bitter rot (<i>Colletotrichum gloeosporioides</i>)</p> <p>Blotch rot (<i>Physalospora vacciniae</i>)</p> <p>Cottonball (<i>Monilinia oxycocci</i>)</p> <p>Leaf rust (<i>Pucciniastrum vaccinii</i>)</p> <p>Ripe rot (<i>Coleophoma empetri</i>)</p>	<p>6-7 (0.10-0.11 lb. ai difenoconazole and propiconazole)</p>	<p>For best activity, apply AG37593 CS EC prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 14 day interval.</p> <p>DO NOT apply more than two sequential applications before alternating to a fungicide with a different mode of action.</p> <p>[Optional language if label has a rate range: If disease pressure is high, use the highest rate.]</p> <p>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</p> <p>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</p>

Application: For best results, sufficient water volume must be used to provide thorough coverage. **AG37593 CS EC** can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy. Applicators should use care in making applications near non-target aquatic habitats.

RESTRICTIONS:

- **DO NOT** apply more than 21 fl. oz./A/year of **AG37593 CS EC** (which delivers 0.34 lb. ai/A/year of both difenoconazole and propiconazole).
- **DO NOT** exceed 3 applications per year.
- **DO NOT** apply more than 0.34 lb. ai/A/year of difenoconazole-containing products.
- **DO NOT** apply more than 0.67 lb. ai/A/year of propiconazole-containing products.
- **DO NOT** allow release of irrigation or flood water to non-target aquatic habitat for at least 14 days after the last application.
- **DO NOT** apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- **DO NOT** treat fields used for aquaculture of fish or crustacean.
- **DO NOT** drain water from treated fields into ponds used for aquaculture of fish or crustacean.
- **DO NOT** use water drained from treated field to irrigate other crops.
- **DO NOT** apply to flooded crop.
- Use is limited to Oregon, Washington, and Wisconsin only.
- **DO NOT** apply **AG37593 CS EC** within 45 days of harvest (45-day PHI).
- Retreatment Interval (RTI) is 14 days.

[* Not Registered for use by California]

Filberts[*]: Hazelnuts

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Eastern Filbert Blight (<i>Anisogramma anomala</i>)	5-7 (0.08-0.11 lb. ai difenoconazole and propiconazole)	<p>Begin applications when green leaf tissue becomes visible and continue on a 14- to 21-day interval. Make no more than 2 sequential applications before alternating to another fungicide with a different mode of action.</p> <p>[Optional language if label has a rate range: If disease pressure is high, use the highest rate.]</p> <p>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.]</p> <p>[Optional language if label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</p> <p>NOTE: Under certain conditions, AG37593 CS EC may cause smaller and/or greener leaves. Yields of filberts displaying these characteristics have not been reduced due to AG37593 CS EC treatments.</p>

Application: For best results, sufficient water volume must be used to provide thorough coverage. **AG37593 CS EC** can be applied by either ground, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.

RESTRICTIONS:

- **DO NOT** apply more than 28 fl. oz./A/year of **AG37593 CS EC** (which delivers 0.46 lb. ai/A/year of difenoconazole and 0.45 lb. ai/A/year of propiconazole).
- **DO NOT** exceed 4 applications per year at the highest rate (7 fl. oz./A) or 5 applications per year at the lowest rate (5 fl. oz./A).
- **DO NOT** apply more than 0.46 lb. ai/A/year of difenoconazole-containing products.

- **DO NOT** apply more than 0.9 lb. ai/A/year of propiconazole-containing products.
- **DO NOT** use an adjuvant.
- **DO NOT** graze livestock in treated area or cut treated cover crop for feed.
- **DO NOT** apply within 60 days of harvest (60-day PHI).
- Retreatment Interval (RTI) is 14 days.

[* Not Registered for use by California]

Pecan[*]

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Downy Spot (<i>Mycosphaerella caryigena</i>) Liver Spot (<i>Gnomonia caryae</i>) Pecan Scab (<i>Cladosporium caryigenum</i>) Powdery Mildew (<i>Microsphaera penicillata</i>) Vein Spot (<i>Gnomonia nerviseda</i>) Zonate Leaf Spot (<i>Grovesinia pyramidalis</i>)	5-7 (0.08-0.11 lb. ai difenoconazole and propiconazole)	Begin applications prior to disease onset when conditions are conducive for disease. Apply AG37593 CS EC on a 14-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. [Optional language if label has a rate range: If disease pressure is high, use the highest rate.]
Application: For best results, sufficient water volume must be used to provide thorough coverage. AG37593 CS EC can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 28 fl. oz./A/year of AG37593 CS EC (which delivers 0.46 lb. ai/A/year of difenoconazole and 0.45 lb. ai/A/year of propiconazole). • DO NOT exceed 4 applications per year at the highest rate (7 fl. oz./A) or 5 applications per year at the lowest rate (5 fl. oz./A). • DO NOT apply more than 0.46 lb. ai/A/year of difenoconazole-containing products. • DO NOT apply more than 0.9 lb. ai/A/year of propiconazole-containing products. • DO NOT use an adjuvant. • DO NOT graze livestock in treated area or cut treated cover crop for feed. • DO NOT apply after shuck split. • Retreatment Interval (RTI) is 14 days. 		
[* Not Registered for use by California]		

Pistachios[*]

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Alternaria late blight (<i>Alternaria spp.</i>) Panicle and Shoot Blight (<i>Botryosphaeria dothidea</i>) Septoria leaf spot (<i>S. pistaciarum</i>)	5-7 (0.08-0.11 lb. ai difenoconazole and propiconazole)	Begin applications prior to disease onset when conditions are conducive for disease. Apply AG37593 CS EC on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. For <i>Botryosphaeria</i> , begin applications when green

		<p>leaf tissue becomes visible and continue on a 14-21 day interval. Make no more than 2 sequential applications before alternating to another fungicide with a different mode of action</p> <p>[Optional language if label has a rate range: If disease pressure is high, use the highest rate.]</p> <p>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language is label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</p> <p>NOTE: Under certain conditions, AG37593 CS EC may cause smaller and/or greener leaves. Yields of pistachios displaying these characteristics have not been reduced due to AG37593 CS EC treatments.</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. AG37593 CS EC can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p>		
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 28 fl. oz./A/year of AG37593 CS EC (which delivers 0.46 lb. ai/A/year of difenoconazole and 0.45 lb. ai/A/year of propiconazole). • DO NOT exceed 4 applications per year at the highest rate (7 fl. oz./A) or 5 applications per year at the lowest rate (5 fl. oz./A). • DO NOT apply more than 0.46 lb. ai/A/year of difenoconazole-containing products. • DO NOT apply more than 0.9 lb. ai/A/year of propiconazole-containing products. • DO NOT use an adjuvant. • DO NOT graze livestock in treated area or cut treated cover crop for feed. • DO NOT apply within 60 days of harvest (60-day PHI). • Retreatment Interval (RTI) is 14 days. <p>[* Not Registered for use by California]</p>		

Rice (including wild)[*]

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Aggregate Sheath Spot (<i>Rhizoctonia oryzae-sativae</i>) Black Sheath Rot (<i>Gaeumannomyces graminis var. graminis</i>) Brown Leaf spot (<i>Cochliobolus miyabeanus</i>) Kernel Smut (<i>Neovossia barclayana</i>) Leaf Smut (<i>Entyloma oryzae</i>) Narrow Brown Leaf Spot (<i>Cercospora oryzae</i>)	7.5 (0.12 lb. ai difenoconazole and propiconazole)	<p>Begin applications prior to disease onset when Apply 7.5 fl. oz./A when disease is less than 4 inches above water line usually between panicle differentiation (PD) +5 days to PD +10 days or at initial sign of disease. Under heavy disease pressure and conditions favorable for disease development, a second application may be applied (minimum of 14 day re-treatment interval).</p> <p>For increased control of false smut or stem rot, tank mix AG37593 CS EC with a triazole-containing fungicide.</p>

Suppression of: False smut (<i>Ustilaginoidea virens</i>) Sheath Blight (<i>Rhizoctonia solani</i>) Sheath Spot (<i>Rhizoctonia oryzae</i>) Stem Rot (<i>Sclerotium oryzae</i>)		
Application: For best results, sufficient water volume must be used to provide thorough coverage AG37593 CS EC can be applied by ground or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15 fl. oz./A/year of AG37593 CS EC (which delivers 0.24 lb. ai/A/year of both difenoconazole and propiconazole). • DO NOT exceed 2 applications per year. • DO NOT apply more than 0.34 lb. ai/A/year of propiconazole-containing products. • DO NOT apply more than 0.24 lb. ai/A/year of difenoconazole-containing products. • DO NOT treat rice fields used for aquaculture of fish or crustacean. • DO NOT drain water from treated rice fields into ponds used for commercial fish farming. • DO NOT apply when weather conditions favor drift from treated areas to non-target aquatic habitat. • DO NOT allow release of irrigation or flood water for at least 7 days after the last application. • DO NOT apply AG37593 CS EC within 35 days of harvest (35-day PHI). • DO NOT apply to stubble or ratoon crop rice. • DO NOT use water drained from treated fields to irrigate other crops. • DO NOT drain water from treated rice fields into ponds used for aquaculture of fish or crustacean. • For wild rice: Use is limited to MN only. • Retreatment Interval (RTI) is 14 days. <p>[* Not Registered for use by California]</p>		

Soybean[*]

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Alternaria leaf spot (<i>Alternaria spp.</i>) Anthracnose (<i>Colletotrichum truncatum</i>) Brown spot (<i>Septoria glycines</i>) Cercospora blight and leaf spot (<i>C. kikuchii</i>) Frogeye leaf spot (<i>Cercospora sojina</i>) Pod and stem blight (<i>Diaporthe phaseolorum</i>) Powdery mildew (<i>Microspheera diffusa</i>)	5-7 (0.08-0.11 lb. ai difenoconazole and propiconazole)	Begin applications prior to disease onset when conditions are conducive for disease. Apply AG37593 CS EC on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. [Optional language if label has a rate range: If disease pressure is high, use the highest rate.] [Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language is label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.] Soybean Rust: Apply 7 fl. oz. at first indication that disease is in the area. For best control,

Soybean rust (<i>Phakopsora pachyrhizi</i>)		<p>preventive applications work best. Repeat on a 14-day interval. Use higher rate when disease is present in field and incidence is less than 2% (2 plants in 100 are infected). If incidence is greater than this or if disease is in the mid-canopy, control will not be acceptable. Scouting for the disease and/or being aware of the proximity of the disease via monitoring systems will aid in the proper timing to maximize the effectiveness of the fungicide applications.</p> <p>On certain varieties, AG37593 CS EC may cause crinkled, smaller and/or greener leaves. Yields of beans displaying these symptoms have not been reduced.</p>
Application: For best results, sufficient water volume must be used to provide thorough coverage. AG37593 CS EC can be applied by either ground or aerial application. For aerial applications, use a minimum of 2 gal/A of water. Addition of an oil-based additive is recommended for improved coverage and penetration when applying by air.		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 13.5 fl. oz./A/year of AG37593 CS EC (which delivers 0.22 lb. ai/A/year of both difenoconazole and propiconazole). • DO NOT exceed 2 applications per year. • DO NOT apply more than 0.22 lb. ai/A/year of difenoconazole-containing products. • DO NOT apply more than 0.34 lb. ai/A/year of propiconazole-containing products. • AG37593 CS EC may be applied up to Stage R6 or within 14 days of harvest, whichever is longer. • DO NOT feed soybean hay, forage and silage. • Retreatment Interval (RTI) is 14 days. <p>[* Not Registered for use by California]</p>		

Stone Fruit Crop Group 12-12[*]: Apricots, Cherries, Sweet Cherries, Tart Nectarines, Peaches, Plums, Plumcot, Prunes, and cultivars and/or hybrids of these.

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Alternaria spot and Fruit rot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum spp.</i>) Brown rot blossom blight and Fruit rot (<i>Monilinia fructicola</i> , <i>M. laxa</i>) Leaf rust (<i>Tranzschelia discolor</i>) Powdery mildew (<i>Sphaerotheca pannosa</i> , <i>Podosphaera clandestina</i>) Scab (<i>Cladosporium carpophilum</i>) Shot hole (<i>Wilsonomyces carpophilus</i>)	5-7 (0.08-0.11 lb. ai difenoconazole and propiconazole)	<p>For brown rot blossom blight, begin applications at early bloom and continue through petal fall. For brown rot on fruit, apply as needed a maximum of two sprays during the pre-harvest period up to the day of harvest (minimum of a 7-day retreatment interval). If high inoculum and severe disease conditions persist, apply a registered non-Group 3 fungicide.</p> <p>For all other diseases, follow the Brown Rot Blossom Blight schedule. Make additional applications on a 10- to 14-day interval from the end of petal fall to harvest.</p> <p>[Optional language if label has a rate range: If disease pressure is high, use the highest rate.]</p> <p>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language is label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</p>
Application: For best results, sufficient water volume must be used to provide thorough coverage. AG37593 CS EC		

can be applied by either ground or aerial application. Stone fruit diseases are most effectively controlled by ground applications. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.

PRECAUTIONS: Applications of propiconazole during bloom to Stanley plums have occasionally caused fruit to be less oval in shape and smaller in size at harvest. To avoid this, **DO NOT** apply A253.02 to Stanley plums earlier than 21 days prior to harvest.

RESTRICTIONS:

- **DO NOT** apply more than 28 fl. oz./A/year of A253.02 (which delivers 0.46 lb. ai/A/year of difenoconazole and 0.45 lb. ai/A/year of propiconazole).
- **DO NOT** exceed 4 applications per year at the highest rate (7 fl. oz./A) or 5 applications per year at the lowest rate (5 fl. oz./A).
- **DO NOT** apply more than 0.46 lb. ai/A/year of difenoconazole-containing products.
- **DO NOT** apply more than 0.56 lb. ai/A/year of propiconazole-containing products.
- **AG37593 CS EC** may be applied on the day of harvest (0-day PHI).
- Retreatment Interval (RTI) is 7 days.

[* Not Registered for use by California]

Sugar Beets

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
<p>Cercospora leafspot (C. beticola)</p> <p>Powdery mildew (Erysiphe polygoni)</p>	<p>5-7 (0.08-0.11 lb. ai difenoconazole and propiconazole)</p>	<p>Begin applications preventively or on a forecast system. For powdery mildew, apply at first sign of disease</p> <p>Apply AG37593 CS EC on a 10- to 21-day schedule. Make only one AG37593 CS EC spray then alternate to a nontriazole fungicide (non-Group 3) that is registered on sugar beets for these diseases.</p> <p>[Optional language if label has a rate range: If disease pressure is high, use the highest rate.]</p> <p>[Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language is label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]</p>
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. AG37593 CS EC can be applied by either ground, chemigation, or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 5 gal/A of water. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>		
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 21 fl. oz./A/year of AG37593 CS EC (which delivers 0.34 lb. ai/A/year of both difenoconazole and propiconazole). • DO NOT exceed 3 applications per year. • DO NOT apply more than 0.46 lb. ai/A/year of difenoconazole-containing products. • DO NOT apply more than 0.34 lb. ai/A/year of propiconazole-containing products. • DO NOT apply within 21 days of harvest (21-day PHI). • Retreatment Interval (RTI) is 10 days. 		

Tree Nuts Crop Group 14-12[*]: Beechnut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Hickory, Macadamia; Walnut, black; Walnut English. (See **Specific Direction for Almonds, Filberts, Pecans, and Pistachios**)

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Foliar Diseases	5-7 (0.08-0.11 lb. ai difenoconazole and propiconazole)	Begin applications prior to disease onset when conditions are conducive for disease. Apply AG37593 CS EC on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action. [Optional language if label has a rate range: If disease pressure is high, use the highest rate.] [Optional language if label has a single rate and interval range: If disease pressure is high, use the shortest interval.] [Optional language is label has a rate range and interval range: If disease pressure is high, use the shortest interval and highest rate.]
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. AG37593 CS EC can be applied by either ground or aerial application. Use a minimum of 15 gal/A of water for ground applications. For aerial applications, use a minimum of 10 gal/A of water.</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 28 fl. oz./A/year of AG37593 CS EC (which delivers 0.46 lb. ai/A/year of difenoconazole and 0.45 lb. ai/A/year of propiconazole). • DO NOT exceed 4 applications per year at the highest rate (7 fl. oz./A) or 5 applications per year at the lowest rate (5 fl. oz./A). • DO NOT apply more than 0.46 lb. ai/A/year of difenoconazole-containing products. • DO NOT apply more than 0.9 lb. ai/A/year of propiconazole-containing products. • DO NOT use an adjuvant. • DO NOT graze livestock in treated areas or cut treated cover crop for feed. • DO NOT apply within 60 days of harvest (60-day PHI). • Retreatment Interval (RTI) is 14 days. <p>[* Not Registered for use by California]</p>		

Watercress[*]

DISEASES CONTROLLED	Use Rate fl. oz. product/A	APPLICATION INSTRUCTIONS
Cercospora leafspot (<i>Cercospora spp.</i>)	7 (0.11 lb. . ai difenoconazole and propiconazole)	For best activity, apply AG37593 CS EC prior to or early in the disease development. An adjuvant may be added at specified rates. Apply on a 7-14 day interval. For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the application. DO NOT apply more than two sequential applications before alternating to a fungicide with a different mode of action.
<p>Application: For best results, sufficient water volume must be used to provide thorough coverage. AG37593 CS EC can be applied by ground, chemigation, or aerial application. For aerial applications, use a minimum of 5 gal/A of water. Applicators should use care in making applications near non-target aquatic habitats.</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 21 fl. oz./A/year of AG37593 CS EC (which delivers 0.34 lb. . ai/A/year of both difenoconazole and propiconazole). 		

- **DO NOT** exceed 3 applications per year.
- **DO NOT** apply more than 0.46 lb. . ai/A/year of difenoconazole-containing products.
- **DO NOT** apply more than 0.45 lb. . ai/A/year of propiconazole-containing products.
- Make no more than 2 applications before harvesting leaves.
- **DO NOT** apply directly to water and **DO NOT** allow water in a treated field for at least 24 hours.
- **DO NOT** apply **AG37593 CS EC** within 30 days of harvest (30-day PHI).
- Retreatment Interval (RTI) is 7 days.

[* Not Registered for use by California]

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used must be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[For Residential uses]

[Nonrefillable container. DO NOT reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.]

[For Commercial Uses]

[For plastic containers ≤ 5 gallons: Nonrefillable Container: DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

[For plastic containers > 5 gallons: Nonrefillable container. DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.]

WARRANTY DISCLAIMER AND NOTICE

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranty and Limitations Liability before using this product. If the terms are not acceptable, return the unopen product at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this

product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aceto Life Sciences, L.L.C. d/b/a Actylis. To the extent consistent with applicable law all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ACETO LIFE SCIENCES, L.L.C. D/B/A ACTYLIS MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aceto Life Sciences, L.L.C. d/b/a Actylis is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, ACETO LIFE SCIENCES, L.L.C. D/B/A ACTYLIS DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ACETO LIFE SCIENCES, L.L.C. D/B/A ACTYLIS'S ELECTION, THE REPLACEMENT OF PRODUCT.

[Option marketing and non-FIFRA label statements]

[All trademarks are the property of their respective owners.]

[Active ingredient made [in] [Country] , [formulated] [in Country] [and] [package] [in Country].]

[Shake Well Before Using]

[*Not Registered for Use in [State]]

[Peel Down for Directions]