



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (7505T)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

EPA Reg. Number:

2749-655

Date of Issuance:

3/31/26

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

AG37589 CS SC Fungicide

Name and Address of Registrant (include ZIP Code):

Aceto Life Sciences, L.L.C. d/b/a Actylis  
 4 Tri Harbor Court  
 Port Washington, NY 11050

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

*Continues page 2*

Signature of Approving Official:

Kristy Crews, Ph.D., Product Manager 22  
 Fungicide Branch, Registration Division (7505P)

Date:

3/31/26

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 2749-655."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):


- Basic CSF dated 07/25/2024
- Alternate CSF 1 dated 07/25/2024

If you have any questions, please contact James Orrock by phone at 202-566-2862 or by email at [orrock.james@epa.gov](mailto:orrock.james@epa.gov).

Enclosure- Stamped Label

[Note to reviewer: [Text] in brackets denotes optional or explanatory language]  
[Master Label]

[Alternate Brand Names: ]



AZOXYSTROBIN	GROUP	11	FUNGICIDE
--------------	-------	----	-----------

## AG37589 CS SC Fungicide

<b>ACTIVE INGREDIENT:</b>	(% by weight)
Azoxystrobin: methyl ( <i>E</i> )-2-[2-[6-(2-cyanophenoxy) pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate*.....	22.9%
<b>OTHER INGREDIENTS**</b> .....	77.1%
<b>TOTAL</b> .....	100.0%

\*IUPAC

Contains 2.08 lb of active ingredient per gallon

### KEEP OUT OF REACH OF CHILDREN. CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 2749-X0XX

EPA Est. No.:

Net Contents:

FIRST AID	
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>● Take off contaminated clothing.</li><li>● Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>● Call a poison control center or doctor for treatment advice.</li></ul>
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or physician or going for treatment. FOR MEDICAL EMERGENCIES INVOLVING THIS PRODUCT, CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887 (24 Hours per Day, 7 Days per Week).	

FOR CHEMICAL SPILL, LEAK, FIRE, EXPOSURE OR MEDICAL EMERGENCY INVOLVING THIS PRODUCT  
CALL CHEMTREC® TOLL FREE 1-800-424-9300 or 1-703-527-3887 (24 Hours per Day, 7 Days per Week).

[See] [inside] [label] [booklet] [for] [additional] [Precautionary Statements][,] [and] [Directions for Use]  
[including] [Storage and Disposal] [instructions][.]

Manufactured [for][by]  
Aceto Life Sciences, L.L.C., d/b/a Actylis  
4 Tri Harbor Court  
Port Washington, NY 11050-4661

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans and Domestic Animals**  
**CAUTION**

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**Personal Protective Equipment (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, polyvinyl chloride (PVC)  $\geq 14$  mils, or Viton  $\geq 14$  mils
- Shoes plus socks

In addition, **mixer/loaders/applicators** using mechanically pressurized handwands except when applying to Christmas tree farms, must wear a minimum of NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Human flagging is prohibited.

Respirator fit testing, medical qualification, and training

Using a program that conforms to OSHA's requirements (see 29 CFR Part 1910.134), employers must verify that any handler who uses a respirator is:

- Fit-tested and fit-checked,
- Trained, and
- Examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn. A qualified medical practitioner is a physician or other licensed health care professional who will evaluate the ability of a worker to wear a respirator. The initial evaluation consists of a questionnaire that asks about medical conditions (such as a heart condition) that would be problematic for respirator use. If concerns are identified, then additional evaluations, such as a physical exam, might be necessary. The initial evaluation must be done before respirator use begins. Handlers must be reexamined by a qualified medical practitioner if their health status or respirator style or use-conditions change.

Upon request by local/state/federal/tribal enforcement personnel, employers must provide documentation demonstrating how they have complied with these requirements.

**USER SAFETY REQUIREMENTS**

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **User Safety Recommendations**

#### **Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **Engineering Controls**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for “applicators and other handlers” and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

### **ENVIRONMENTAL HAZARDS**

Azoxystrobin is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Azoxystrobin can be persistent for several months or longer.

**DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

#### **Ground Water Advisory**

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

#### **Surface Water Advisory**

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Aceto Life Sciences, L.L.C., d/b/a Actylis immediately if you observe any adverse environmental effects due to use of this product.

## PHYSICAL AND CHEMICAL HAZARDS

**DO NOT** mix or allow coming in contact with oxidizing and reducing agent hazardous chemical reaction may occur.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, polyvinyl chloride (PVC)  $\geq 14$  mils, or Viton  $\geq 14$  mils
- Shoes plus socks

## USE PRECAUTIONS

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

## POLLINATOR ADVISORY STATEMENT

This product may adversely impact the forage and habitat of local pollinators, including the monarch butterfly (and its larvae), birds, or bats if it reaches non-target areas. Protect pollinators by following label directions to minimize spray drift.

## PRODUCT INFORMATION

AG37589 CS SC Fungicide is a broad spectrum, preventative fungicide with systemic and curative properties for the control of many important plant diseases. AG37589 CS SC Fungicide Flowable Fungicide may also improve the yield and/or quality of the crop. These additional benefits are due to positive effects on plant physiology. The effects may vary according to factors including the crop, crop hybrid, or environment. AG37589 CS SC Fungicide may be applied as a foliar spray in alternating spray programs or in tank mixes with other registered crop protection products. All applications must be made according to the use directions that follow.

## PRODUCT USE INSTRUCTIONS

**Application:** Thorough coverage is necessary to provide good disease control. Make no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

**Adjuvants:** When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification is advised.

**Efficacy:** Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of AG37589 CS SC Fungicide has been used. If resistant isolates to Group 11 fungicides are present, efficacy can be reduced for certain diseases. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, with highly susceptible varieties, or when environmental conditions are conducive to disease.

## INTEGRATED PEST (DISEASE) MANAGEMENT

Integrate AG37589 CS SC Fungicide into an overall disease and pest management strategy whenever the use of a fungicide is required. Follow cultural practices known to reduce disease development, including selection of varieties with disease tolerance, removal of plant debris in which inoculum overwinters, and proper timing and placement of irrigation.

Consult your local agricultural authorities for additional IPM strategies established for your area. AG37589 CS SC Fungicide may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

**Crop Tolerance:** Plant tolerance has been found to be acceptable for all crops on the label, however, not all possible tank-mix combinations have been tested under all conditions. When possible, test the combinations on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of application. See Product Use Precautions for apple phytotoxicity information.

## RESISTANCE MANAGEMENT

For resistant management, AG37589 CS SC Fungicide contains a Group 11 fungicide. Any fungal population may contain individuals naturally resistant to AG37589 CS SC Fungicide and other Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed. Conform to resistance management strategies established for the crop and use area when using this product. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label.

Aceto Life Sciences, L.L.C., d/b/a Actylis encourages responsible resistance management to ensure effective long- term control of the fungal diseases on this label. Follow the crop specific resistance management specifications in the directions for use.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of AG37589 CS SC Fungicide or other Group 11 fungicides (strobilurins, including pyraclostrobin and trifloxystrobin) within a growing season sequence with different fungicide groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the

target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.

- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Aceto Life Sciences, L.L.C., d/b/a Actylis representatives at 1-516-627-6000 or visit the Fungicide Resistance Action Committee (FRAC) on the web at [www.frac.info](http://www.frac.info). You can also contact your pesticide distributor or university extension specialist to report resistance.

If there are no resistance management directions on the number of applications in the directions for use, then follow the directions in the table below.

<b>If planned total number of fungicide applications per crop is:</b>	1	2	3	4	5	6	7	8	9	10	11	12
<b>Specified Solo QoI fungicide sprays</b>	1	1	2	2	2	2	2	3	3	3	3	4
<b>Specified QoI fungicide sprays in mixture (tank-mix or formulated)</b>	1	2	2	2	2	3	3	4	4	5	5	6

In situations requiring multiple sprays, develop season long spray programs for Group 11 (QoI) fungicides. In crops where two sequential Group 11 fungicide applications are made, alternate with two or more applications of a fungicide that is not in Group 11. If more than 12 applications are made, observe the following guidelines:

- When using a QoI fungicide as a solo product, the number of applications must be no more than ½ (33%) of the total number of fungicide applications per season.
- For QoI mixes in programs in which tank mixes or pre-mixes of QoI with mixing partners of a different mode of action are utilized, the number of QoI containing applications must be no more than ½ (50%) of the total number of fungicide applications per season.
- In programs in which applications of QoI are made with both solo products and mixtures, the number of QoI containing applications must be no more than ½ (50%) of the total number of fungicide applications per season.

If a Group 11 fungicide is applied to the seed or soil, **DO NOT** make another application with a Group 11 fungicide for at least 3 weeks.

### Rotational Crop Restrictions

The following crops may be planted at the specified interval following application of AG37589 CS SC Fungicide.

## Crop Rotational Interval

	<b>Plant back interval</b>
Buckwheat and millet	12 months
All other crops with Azoxystrobin registered uses	0 days

## SOILBORNE/SEEDLING DISEASE CONTROL

For those crops that have specific use directions for soilborne disease control: AG37589 CS SC Fungicide can provide control of many soilborne diseases if applied early in the growing season. Specific applications for soilborne diseases include in-furrow applications and banded applications applied over the row, either shortly after plant emergence or during herbicide applications or cultivation. These applications will provide control of pre- or postemergence damping off and diseases that infect plants at the soil-plant interface.

The use of either type of application depends on the cultural practices in the region. In some locations, one type of application may provide better disease control than the other, depending on the timing of the disease epidemic. Seedling diseases are generally controlled by in-furrow applications while banded applications are more effective against soilborne diseases that develop later in the season. Consult your local expert to get some guidance regarding application type.

Under cool, wet conditions, crop injury from soil directed applications can occur.

### BANDED

- Apply AG37589 CS SC Fungicide prior to infection as a directed spray to the soil, using single or multiple nozzles, adjusted to provide thorough coverage of the lower stems and the soil surface surrounding the plants.
- Limit band width to 7 inches or less.
- Apply AG37589 CS SC Fungicide at a rate of 0.40-0.80 fl oz product/1000 row feet (0.15-0.30 lb ai/A). For banded applications on 22-inch rows, the maximum application rate is 0.70 fl oz/1000 row feet (0.26 lb ai/A).
- These applications come into contact with the foliage and are counted as foliar applications when considering resistance management.
- They may be applied during cultivation or hilling operations to provide soil incorporation.

### IN-FURROW

- Apply AG37589 CS SC Fungicide as an in-furrow spray in 3-15 gallons of water at planting.
- Mount the spray nozzle so the spray is directed into the furrow just before the seeds are covered.
- Use the higher rate when the weather conditions are expected to be conducive for disease development, if the field has a history of Pythium problems, or if minimum/low till programs are in place.

## IN-FURROW APPLICATION RATES

RATE per 1,000 Row Feet	RATE OF AG37589 CL 4SC FUNGICIDE FOR IN-FURROW AND BANDED APPLICATIONS
----------------------------	---

fl oz product	lb ai/A	22	30	32	34	36	38	40	48	60	72	80
		Product per Acre (fl oz)										
0.40	0.15	9.5	7.0	6.5	6.1	5.8	5.5	5.2	4.4	3.5	2.9	2.6
0.60	0.23	14.3	10.5	9.8	9.2	8.7	8.3	7.8	6.5	5.2	4.4	3.9
0.80	0.30		13.9	13.1	12.3	11.6	11.0	10.5	8.7	7.0	5.8	5.2
1.00	0.38					14.5	13.8	13.1	10.9	8.7	7.3	6.5
1.20	0.45								13.1	10.5	8.7	7.8
1.38	0.54								15.0	12.0	10.0	9.0
1.50	0.60									13.1	10.9	9.8
1.72	0.68									15.0	12.5	11.2
2.00	0.75										14.5	13.1
2.07	0.81										15.0	13.5
2.30	0.90											15.0

Do not apply more than 15 fl oz/A.

Row spacing (inches)	Row-Foot Per Acre
22	23,760
30	17,424
32	16,335
34	15,374
36	14,520
38	13,756
40	13,068
48	10,890
60	8,712
72	7,260
80	6,534

### DRIP

Refer to the **Application Instructions Through Irrigation System** section.

### USE RESTRICTIONS

- **DO NOT** use AG37589 CS SC Fungicide through airblast application equipment on grapes in the following townships and boroughs of Erie County, Pennsylvania: North East, Harborcreek, Lawrence Park, Erie, Presque Isle, Millcreek, Fairview, Girard and Springfield. This prohibition is intended to help eliminate phytotoxicity problems with apples observed in this geographic location.
- To help manage fungicide resistance, **DO NOT** use for commercial transplant production in the greenhouse except where specified on the label.

## PHYTOTOXICITY

AG37589 CS SC Fungicide is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

**DO NOT** spray AG37589 CS SC Fungicide where spray drift may reach apple trees.

**DO NOT** use spray equipment which has been previously used to apply AG37589 CS SC Fungicide to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

**DO NOT** apply when weather conditions favor drift from treated areas to non-target aquatic habitat.

## SPRAY DRIFT MANAGEMENT

To avoid spray drift, **DO NOT** apply when conditions favor drift beyond the target area. The interaction of many equipment and weather related factors determine the potential for spray drift. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

### MANDATORY SPRAY DRIFT

#### Aerial Applications

- **DO NOT** release spray at a height greater than 10 ft. above the crop canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzles that deliver Medium to coarse spray droplets in accordance with ASABE Standard S-572.1.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

#### Groundboom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

## **SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.  
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

#### **Controlling Droplet Size-Ground Boom**

- Volume—Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure—Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle—Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

#### **Controlling Droplet Size-Aircraft**

- Adjust Nozzles—Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### **BOOM HEIGHT—Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **RELEASE HEIGHT—Aircraft**

Higher release heights increase the potential for spray drift.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### **WIND**

Drift potential generally increase with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## MIXING AND APPLICATION METHODS

### Spray Equipment

AG37589 CS SC Fungicide may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

#### Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Ensure that nozzles are the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Ensure that screens placed on the suction side of the pump are *16-mesh or coarser*.
- **DO NOT** place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's directions.

#### Pump

- Use a pump with capacity to:
  - (1) Maintain 35-40 psi at nozzles.
  - (2) Provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- **DO NOT** air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and state guidance. For specific local directions and spray schedules, consult the current state agricultural specifications.

#### Mixing Instructions

- AG37589 CS SC Fungicide is a suspension concentrate (SC) formulation.
- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

#### AG37589 CS SC Fungicide Alone (No Tank Mix)

- Add ½-½ of the required amount of water to the spray or mixing tank.
- With the agitator running, add AG37589 CS SC Fungicide to the tank.

- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after AG37589 CS SC Fungicide has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

**AG37589 CS SC Fungicide + Tank Mixtures:** AG37589 CS SC Fungicide is usually compatible with all tank-mix partners listed on this label. To determine the physical compatibility of AG37589 CS SC Fungicide with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

AG37589 CS SC Fungicide has demonstrated some phytotoxic effects when mixed with products that are formulated as emulsifiable concentrates (EC). These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.

### **Mixing in the Spray Tank**

- Add ½-½ of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and AG37589 CS SC Fungicide to the spray tank.
- Allow AG37589 CS SC Fungicide to completely disperse.
- Spray the mixture with the agitator running.

## **APPLICATION INSTRUCTIONS THROUGH IRRIGATION SYSTEMS (CHEMIGATION)**

### **Application Through Irrigation Systems (Chemigation)**

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. **DO NOT** apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.1-0.25 inches/acre. Excessive water may reduce efficacy.
- If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments when the need arises.

**Spray Preparation:** Thoroughly clean the chemical tank and injector system. Flush system with clean water.

**Drip Irrigation:** AG37589 CS SC Fungicide may be applied through drip irrigation systems for soilborne disease control. Ensure that the soil has adequate moisture capacity prior to drip application.

Terminate drip irrigation at fungicide depletion from the main feed supply tank or after 6 hours from start, whichever is shorter. For maximum efficacy, delay subsequent irrigation (water only) for at least 24 hours following drip application.

### **Sprinkler Irrigation**

- Apply this product through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move irrigation systems.
- **DO NOT** apply this product through any other type of irrigation system except as specified on this label.
- Apply with center pivot or continuous-move equipment distributing ½ acre-inch or less during treatment.
- In general, use the least amount of water required for proper distribution and coverage.
- If stationary systems (solid set, handlines or wheel lines other than continuous-move) are used, inject this product into no more than the last 20-30 minutes of the set.
- **DO NOT** apply when winds are greater than 10-15 mph to avoid drift or wind skips.
- **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.
- Plant injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform treated water.
- Thorough coverage of foliage is required for good control.
- Maintain good agitation during the entire application period.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.

### **Operating Instructions**

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water- source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick- closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments when the need arises.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label- prescribed safety devices for public water systems are in place.

### **Center Pivot Irrigation Equipment**

**Notes:** (1) Use only with drive systems which provide uniform water distribution. (2) **DO NOT** use end guns when chemigating AG37589 CS SC Fungicide through center pivot systems because of non-uniform application.

- Determine the size of the area to be treated.
- Determine the time required to apply ½-½ inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as specified by the equipment manufacturer. When applying AG37589 CS SC Fungicide through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of AG37589 CS SC Fungicide required to treat the area covered by the irrigation system.
- Add the required amount of AG37589 CS SC Fungicide and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the AG37589 CS SC Fungicide solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the AG37589 CS SC Fungicide solution has cleared the sprinkler head.

### **Solid Set, Hand Move, and Moving Wheel Irrigation Equipment**

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30-minute interval. When applying AG37589 CS SC Fungicide through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of AG37589 CS SC Fungicide required to treat the area covered by

the irrigation system.

- Add the required amount of AG37589 CS SC Fungicide into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the AG37589 CS SC Fungicide solution has cleared the last sprinkler head.

### **Specific Instructions for Public Water Systems**

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

## USE DIRECTIONS FOR SPECIFIC CROPS

**ALFALFA** (See Nongrass Animal Feeds Forage, Fodder, Straw and Hay)

### ALMONDS

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Alternaria Leaf and Fruit spot ( <i>Alternaria alternata</i> ) Anthracnose ( <i>Colletotrichm acutatum</i> ) Leaf Blight ( <i>Seimatosporium lichenicola</i> ) Leaf Rust ( <i>Tranzshcelia discolor</i> ) Scab ( <i>Cladosporium carpophilum</i> ) Shot hole ( <i>Wilsonomyces carpophilus</i> )	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Begin applications prior to disease development and continue throughout the season following the resistance management guidelines. Applications may be made by ground, air or chemigation. For aerial applications apply in a minimum of 15 GPA. Thorough and uniform coverage is essential for disease control. Reduced efficacy has been observed when uniform coverage cannot be obtained.  AG37589 CS SC Fungicide may be applied by air growth stages prior to and including 5 weeks after petal fall. An adjuvant may be added at specified rates.  Anthracnose, scab and shot hole: Begin applications prior to disease development and continue at 7- to 14-day intervals throughout the season.
Brown Rot Blossom Blight ( <i>Monilinia laxa</i> , <i>M. fructicola</i> )	12.0-15.5 fl. oz.. (0.20-0.25 lb. ai/A)	Blossom blights: Begin applications at early bloom and continue through petal fall.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A). When applying at 12.0 fl oz/A, <b>DO NOT</b> apply more than 7 applications per year.</li> <li>● <b>DO NOT</b> apply within 28 days of harvest (28-day PHI)</li> </ul>		

## ARTICHOKE, GLOBE

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Ramularia Leaf Spot <i>(Ramularia cynarae)</i>	11.0-15.5 fl. oz. (0.18-0.25 lb. ai/A)	Begin applications prior to disease onset when conditions are conducive for disease, and continue as needed throughout the season at a 2-3 week interval, up to and including the day of harvest. <b>DO NOT</b> apply at less than 7-day intervals.
<p><b>Application:</b> For best results, sufficient water volume must be used to provide thorough coverage. AG37589 CS SC Fungicide can be applied by ground, chemigation, or aerial application. For ground applications, apply in 50-200 gallons of water per acre to obtain coverage without excessive runoff. For aerial applications, use a minimum of 5 gal/A of water. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>		
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 88 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 8 applications per year at the low rate (11.0 fl oz/A).</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

## ASPARAGUS

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Stemphyllium Purple Spot <i>(Stemphyllium vesicarium)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	<p>Begin applications prior to disease development and continue throughout the season on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. Use a minimum of 10 gallons of water per acre by ground, and minimum of 3 gallons per acre by air. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
<p><b>RESTRICTIONS</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>DO NOT</b> apply within 100 days of harvest (100-day PHI).</li> </ul>		

## BANANAS PLANTAINS

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Black Sigatoka <i>(Mycosphaerella fijiensis)</i> Yellow Sigatoka <i>(Mycosphaerella musicola)</i>	5.5-8.5 fl. oz. (0.09- 0.135 lb. ai/A)	Begin applications prior to disease development and continue throughout the season every 12-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>RESTRICTIONS</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>• <b>DO NOT</b> apply at less than 12-day interval.</li> <li>• <b>DO NOT</b> apply more than 66 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.08 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>• <b>DO NOT</b> apply more than 7 applications per year at the high rate (8.5 fl oz/A) or 12 applications per year at the low rate (5.5 fl oz/A).</li> <li>• AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

## CEREALS - Barley Oats, Rye

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Kernel Blight or Black Point <i>(Alternaria spp.)</i> <i>(Cochliobolus sativus)</i> Leaf Rust ( <i>Puccinia hordei</i> ) ( <i>P. recondita</i> )	6.0-12.0 fl. oz. (0.10-0.20 lb. ai/A)	Apply prior to disease development. Protecting the flag leaf is important for maximizing disease control. For best results, sufficient water volume must be used to provide thorough coverage.
Barley Stripe <i>(Drechslera graminea = Pyrenophora graminea)</i> Net Blotch ( <i>Pyrenophora teres</i> ) Scald ( <i>Rhynchosporium secalis</i> ) Septoria Leaf and Glume Blotch <i>(Septoria spp., Stagonospora spp.)</i> Spot Blotch <i>(Cochliobolus sativus)</i> Stem Rust <i>(Puccinia graminis f.sp. tritici)</i> Stripe Rust <i>(Puccinia striiformis)</i> Tan Spot <i>(Pyrenophora trichostroma)</i>	9.0-12.0 fl. oz. (0.15-0.20 lb. ai/A)	AG37589 CS SC Fungicide can be applied by ground, air or chemigation. A crop oil concentrate adjuvant may be added at 1.0% v/v to optimize efficacy. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy. DO NOT apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11. <b>DO NOT</b> make more than two (2) applications of AG37589 CS SC Fungicide or other Group 11 fungicide per season.
Powdery Mildew ( <i>Erysiphe graminis f. sp. hordei</i> ) Stagonospora Blotch <i>(Stagonospora nodorum)</i>	12.0 fl. oz. (0.20 lb. ai/A)	

**RESTRICTIONS:**

- **DO NOT** exceed the maximum rate listed in this crop table.
- Do not apply after Feekes 10.54.
- **DO NOT** apply at less than 14-day interval.
- **DO NOT** apply more than 24 fl oz product/A/year.
  - **DO NOT** apply more than 0.40 lb ai/A/year of azoxystrobin-containing products.
- **DO NOT** apply more than 2 applications per year at the high rate (12.0 fl oz/A) or 4 applications per year at the low rate (6.0 fl oz/A). When applying at 9.0 fl oz/A, **DO NOT** apply more than 2 applications per year.
- **DO NOT** apply within 7 days of grazing or harvest (7-day PHI) for forage and hay.

**BERRIES, BUSHBERRY, CROP SUBGROUP 13-07B** Including: Aronia berry; blueberry, highbush; blueberry, lowbush; buffalo currant; Chilean guava; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; Jostaberry; Juneberry (Saskatoon berry); Lingonberry; native currant; Salal; sea buckthorn; cultivars, varieties, and/or hybrids of these

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Alternaria Fruit Rot <i>(Alternaria spp.)</i> Anthracnose Fruit Rot <i>(Colletotrichum gloeosporoides)</i> Botryosphaeria Canker <i>(Botryosphaeria spp.)</i> Leaf Spot and Blotch <i>(Mycosphaerella spp., Septoria spp.)</i> Mummyberry <i>(Monilinia vaccinii-corymbosi)</i> Phomopsis Leaf Spot, Twig Blight and Stem Canker <i>(Phomopsis vaccinii)</i> Powdery Mildew <i>(Sphaerotheca spp.)</i> Septoria Blight <i>(Septoria spp.)</i> Spur Blight <i>(Didymella spp., Phoma spp.)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Begin applications prior to disease development and continue throughout the season on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.

**RESTRICTIONS**

- **DO NOT** exceed the maximum rate listed in this crop table.
- **DO NOT** apply at less than 7-day interval.
- **DO NOT** apply more than 42 fl oz of product/A/year.
  - **DO NOT** apply more than 0.75 lb ai/A/year of azoxystrobin-containing products.
- **DO NOT** apply more than 2 applications per year at the high rate (15.5 fl oz/A) or 7 applications per year at the low rate (6.0 fl oz/A).
- AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).

**BERRIES, CANEBERRY, CROP SUBGROUP 13-07A** Including: Blackberry, Bingleberry, Boysenberry, Dewberry, Lowberry, Marionberry, Olallieberry, Youngberry, Loganberry, Red and Black Raspberry, Wild Raspberry; cultivars, varieties, and/or hybrids of these

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Anthracnose <i>(Spaceloma necator)</i> <i>(Elsinoe veneta)</i> Botryosphaeria Canker <i>(Botryosphaeria dothidea)</i> Colletotrichum Rot <i>(Colletotrichum gloeosporioides)</i> Leaf Spot and Blotch <i>(Mycosphaerella spp.)</i> <i>(Septoria rubi)</i> <i>(Sphaerulina rubi)</i> Powdery Mildew <i>(Sphaerotheca macularis)</i> <i>(Microphaera spp.)</i> <i>(Oidium spp.)</i>  Rosette or Double Blossom of Blackberries <i>(Cercospora rubi)</i> Spur Blight <i>(Didymella applanata)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Begin applications at onset of disease and continue as required until harvest. Make applications on a 7- to 14-day schedule. Use a minimum water volume of 10 gallons per acre by ground and a minimum of 3 gallons by air.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
Blackberry Rust <i>(Phragmidium spp.)</i>	10-15.5 (0.16-0.25 lb. ai/A)	
<b>RESTRICTIONS</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year.               <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A). When applying at 10 fl oz/A, <b>DO NOT</b> apply more than 9 applications per year.</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**BERRIES, LOW GROWING, CROP SUBGROUP 13-07G (except Cranberry)** Including: Strawberry, Bearberry, Bilberry, Cloudberry, Muntries, Partridgeberry including all cultivars and/or hybrids of these.

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Anthracnose <i>(Colletotrichum fragariae)</i> Leather Rot <i>(Phytophthora cactorum)</i> Powdery Mildew <i>(Sphaerotheca macularis)</i>  Suppression of Botrytis on the Foliage <i>(Botrytis cinerea)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	<p>Begin applications prior to disease development and continue throughout the season on a 7- to 10-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p>For leather rot control apply 2 applications on a 7-day schedule from late bloom through harvest.</p> <p>Field Nurseries: Apply to young plants in field nurseries by ground, drip, or overhead chemigations.</p> <p>If applying through drip irrigation, calculate the rate as a band application with a band width equal to the root zone width. Inject AG37589 CS SC Fungicide into the irrigation water.</p> <p><b>For dip applications at transplanting for commercial berry production:</b> For suppression of root and crown rot caused by <i>Colletotrichum</i> spp., mix 5-8 fl oz of AG37589 CS SC Fungicide per 100 gallons of water. Dip plants for 2-5 minutes. Plant treated plants as quickly as possible. It is advised that transplants be washed to remove excess soil prior to dipping. For continued anthracnose control, follow with foliar applications beginning 2-3 weeks after transplant.</p> <p><b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
<b>Soilborne Diseases</b> Seedling Root Rot, Basal Stem Rot <i>(Rhizoctonia solani)</i>	0.40-0.80 fl oz/1000 row feet (0.0065- 0.013 lb. ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<p><b>RESTRICTIONS</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 60 fl oz of product/A/year.               <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.0 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 3 applications per year at the high rate (15.5 fl oz/A) or 10 applications per year at the low rate (6.0 fl oz/A).</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**BERRIES, LOW GROWING, CROP SUBGROUP 13-07H(except Strawberry)**

Including: Cranberry, Bearberry, Bilberry, Cloudberry, Lingonberry, Muntries, and Partridgeberry including all cultivars and/or hybrids of these.

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Cottonball ( <i>Monilinia oxycocci</i> ) Fruit Rots ( <i>Physalospora vaccinii</i> ) ( <i>Glomerella cingulata</i> ) ( <i>Coleophoma empetri</i> ) Lophodermium Twig Blight ( <i>Lophodermium</i> spp.)	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Begin applications at 5-10% bloom for fruit rot, cottonball, and twig blight. Continue applications on a 7- to 14-day schedule if conditions are favorable for disease development. Applications may be made by ground, chemigation or air.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
Fairy Ring (suppression) ( <i>Psilocybe</i> spp.)	15.5 fl. oz. (0.25 lb. ai/A)	Make the first application at bud break. Measure the ring diameter and add 10 feet to that diameter. Apply AG37589 CS SC Fungicide at a rate equivalent to 15.5 fl oz/A in 30 – 100 gallons of water to the affected area.  Irrigation (1 – 2 hours) following application is advisable to ensure penetration to the base of the plant. If necessary make another application 2 – 4 weeks later. For ground application ensure adequate water volume for thorough canopy penetration.
<p><b>RESTRICTIONS</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>DO NOT</b> treat cranberry fields used for aquaculture of fish and crustacea.</li> <li>● <b>DO NOT</b> apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Use care in making applications near non-target aquatic habitats.</li> <li>● <b>DO NOT</b> apply to flooded crop.</li> <li>● <b>DO NOT</b> allow release of irrigation or flood water to non-target aquatic habitat for at least 14 days after the last application.</li> <li>● <b>DO NOT</b> apply within 3 days of harvest (3-day PHI).</li> </ul>		

**BRASSICA, HEAD and STEM SUBGROUP 5A including:** Broccoli Chinese, Broccoli (gai lon), Brussels Sprouts, Cabbage Chinese, Cabbage (napa), Chinese Mustard Cabbage (gai choy), Cauliflower Cavalo, Broccolo Kohlrabi; Including all cultivars and/or hybrids of these

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Alternaria Leaf Spot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum spp.)</i> Cercospora Leaf Spot <i>(Cercospora brassicicola)</i> Downy Mildew <i>(Peronospora parasitica)</i> Pin Rot <i>(Alternaria spp.)</i> Powdery Mildew <i>(Erysiphe polygoni)</i> Rhizoctonia Blight <i>(Rhizoctonia solani)</i> Ring Spot <i>(Mycosphaerella brassicicola)</i> White Leaf Spot <i>(Pseudocercospora capsellae)</i> White Rust <i>(Albugo candida)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Begin applications prior to disease development and continue throughout the season on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. Use a minimum of 10 gallons of water per acre by ground, and minimum of 3 gallons per acre by air.  <b>DO NOT</b> apply more than two applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<p><b>RESTRICTIONS</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A).</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**BRASSICA, LEAFY GREENS SUBGROUP 5B including:** Broccoli Raab, Cabbage, Chinese Collards, Kale, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens, and including all cultivars and/or hybrids of these

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Alternaria Leaf Spot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum spp.)</i> Black Spot <i>(Alternaria spp.)</i> Cercospora Leaf Spot <i>(Cercospora spp.)</i> Downy Mildew <i>(Peronospora parasitica)</i> Powdery Mildew <i>(Erysiphe polygoni)</i> Ring Spot <i>(Mycosphaerella brassicicola)</i> White Rust ( <i>Albugo candida</i> )	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Begin applications prior to disease development and continue throughout the season on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>Soilborne Diseases</b> Seedling Root Rot, Basal Stem Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet (0.0065-0.013 lb ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<b>RESTRICTIONS</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 42 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.75 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 2 applications per year at the high rate (15.5 fl oz/A) or 7 applications per year at the low rate (6.0 fl oz/A).</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**BULB VEGETABLES: GARLIC, LEEK,**

**BULB ONION SUBGROUP 3-07A:** Daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, bulb; onion, Chinese, bulb; onion, pearl; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these.

**GREEN ONION SUBGROUP 3-07B:** Chive, fresh leaves; chive, Chinese, fresh leaves; elegans hosta; fritillaria, leaves; kurrat; lady's leek; leek; leek, wild; onion, Beltsville bunching; onion, fresh; onion, green; onion, macrostem; onion, tree, tops; onion, Welsh, tops; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
<b>Foliar Diseases</b> Cladosporium Leaf Blotch <i>(Cladosporium allii)</i> Powdery Mildew <i>(Leveillula taurica)</i> Purple Blotch and Leaf Blight <i>(Alternaria porri)</i> <i>(Stemphylium vesicarium)</i> Rust <i>(Puccinia allii)</i>	6.0-12.0 fl. oz. (0.10-0.20 lb. ai/A)	For downy mildew, make preventative applications on a 5- to 7-day schedule.  For all other diseases, begin applications prior to disease development and continue throughout the season every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. If applications are made by air, use the higher rates for adequate control. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
Botrytis Leaf Blight <i>(Botrytis aclada)</i> Downy Mildew <i>(Peronospora destructor)</i>	9.0-15.5 fl. oz. (0.15-0.25 lb. ai/A)	Mixtures of AG37589 CS SC Fungicide with insecticides and silicone adjuvants must be tested for crop safety before application to the crop.
<b>Soilborne Diseases</b> Rhizoctonia Damping-Off <i>(Rhizoctonia solani)</i>	0.40-0.80 fl oz/1000 row feet (0.0065- 0.013 lb ai/1000 row feet)	For soilborne/seedling disease control, see directions under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section. If the application is an in-furrow application, spray just prior to seed placement so that the majority of the chemical is under the seed. This will reduce the potential for phytotoxicity, especially if fertilizer is added to the application.
<b>RESTRICTIONS</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 5-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A). When applying at 9.0 fl oz/A, <b>DO NOT</b> apply more than 10 applications per year. When applying at 12.0 fl oz/A, <b>DO NOT</b> apply more than 7 applications per year.</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**CANOLA (see Oilseed Crops for additional information)**

<b>DISEASE CONTROLLED</b>	<b>RATE AG37589 CS SC Fungicide/ACRE</b>	<b>APPLICATION INSTRUCTIONS</b>
Alternaria Blackspot <i>(Alternaria spp.)</i> Blackleg ( <i>Leptosphaeria maculans</i> ) Sclerotinia Stem Rot <i>(Sclerotinia sclerotiorum)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	<p>In general, apply 7.0 fl oz of AG37589 CS SC Fungicide at early bud followed by 14.0 fl oz at about 45 days before harvest. A third application of 7.0 fl oz may be made 30 days before harvest.</p> <p>Specifically for blackleg, make applications at the 2- to 4-leaf stage. For Alternaria or Sclerotinia, apply 9.0-15.5 fl oz product/A at 10-25% flowering (3-7 days following first flower). Use the higher rate under heavy disease pressure or when conditions are favorable for disease. For control of Alternaria alone, 8.0 fl oz product/A may be applied at pod stage (approximately 95% petal fall).</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Applications may be made by ground, air or chemigation. Use a minimum of 10 gallons of water per acre for ground applications.</p>

**RESTRICTIONS**

- **DO NOT** exceed the maximum rate listed in this crop table.
- **DO NOT** apply at less than 14-day interval.
- **DO NOT** apply more than 24 fl oz of product/A/year.
  - **DO NOT** apply more than 0.45 lb ai/A/year of azoxystrobin-containing products.
- **DO NOT** apply more than 1 application per year at the high rate (15.5 fl oz/A) or 4 applications per year at the low rate (6.0 fl oz/A).
- **DO NOT** apply within 30 days of harvest (30-day PHI).

**CARROTS**

<b>DISEASE CONTROLLED</b>	<b>RATE AG37589 CS SC Fungicide/ACRE</b>	<b>APPLICATION INSTRUCTIONS</b>
Cercospora Leaf Spot <i>(Cercospora spp.)</i> Early Blight ( <i>Cercospora carotae</i> ) Late Blight <i>(Alternaria dauci)</i> Powdery Mildew <i>(Erysiphe spp.)</i> White Mold <i>(Sclerotium rolfsii)</i>  For additional diseases, see <b>Vegetables, Root, Subgroup.</b>	9.0-20.0 fl. oz. (0.15-0.33 lb. ai/A)	<p>Begin applications prior to disease development and continue throughout the season every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>

<b>Soilborne Diseases</b> Rhizoctonia Root Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet (0.0065-0.013 lb. ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
---	---	---

**RESTRICTIONS**

- **DO NOT** exceed the maximum rate listed in this crop table.
- **DO NOT** apply at less than 7-day interval.
- **DO NOT** apply more than 120 fl oz of product/A/year.
  - **DO NOT** apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.
- **DO NOT** apply more than 6 applications per year at the high rate (20.0 fl oz/A) or 13 applications per year at the low rate (9.0 fl oz/A).
- AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).

**CELERY**

<b>DISEASE CONTROLLED</b>	<b>RATE AG37589 CS SC Fungicide/ACRE</b>	<b>APPLICATION INSTRUCTIONS</b>
Early Blight ( <i>Cercospora apii</i> ) Late Blight ( <i>Septoria apicola</i> )  For additional diseases, see <b>Leafy Vegetables</b> .	9.0-15.5 fl. oz. (0.15-0.25 lb. ai/A)	Begin applications prior to disease development and continue throughout the season every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>Soilborne Diseases</b> Rhizoctonia Root Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet (0.0065-0.013 lb ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**RESTRICTIONS**

- **DO NOT** exceed the maximum rate listed in this crop table.
- **DO NOT** apply at less than 7-day interval.
- **DO NOT** apply more than 90 fl oz of product/A/year.
  - **DO NOT** apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- **DO NOT** apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 10 applications per year at the low rate (9.0 fl oz/A).
- AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).

## CHRISTMAS TREES

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Diplodia Tip Blight <i>(Diplodia pinea)</i> Lophodermium Needlecast <i>(Lophodermium pinastri)</i> Swiss Needlecast <i>(Phaeocryptopus gaumannii)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Begin applications prior to disease development and continue throughout the season at 7- to 21-day intervals following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>RESTRICTIONS</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>• <b>DO NOT</b> apply at less than 7-day interval.</li> <li>• <b>DO NOT</b> apply more than 120 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>• <b>DO NOT</b> apply more than 7 applications per year at the high rate (15.5 fl oz/A) or 20 applications per year at the low rate (6.0 fl oz/A).</li> <li>• <b>DO NOT</b> apply with handheld equipment.</li> </ul>		

**CITRUS FRUIT CROP 10-10:** Australian Desert Lime (*Eremocitrus glauca*); Australian Finger Lime (*Microcitrus australasica*); Australian Round Lime (*Microcitrus australis*); Brown River Finger Lime (*Microcitrus papuana*); Calamondin (*Citrofortunella microcarpa*); Citron (*Citrus medica*); Citrus Hybrids, *Citrus* spp., *Eremocitrus* spp., *Fortunella* spp., *Microcitrus* spp., and *Poncirus* spp.; Grapefruit (*Citrus paradise*); Japanese Summer Grapefruit (*Citrus natsudaidai*); Kumquat (*Fortunella* spp.); Lemon (*Citrus limon*); Lime (*Citrus aurantiifolia*); Mediterranean Mandarin (*Citrus deliciosa*); Mount White Lime (*Microcitrus garrowayae*); New Guinea Wild Lime (*Microcitrus warburgiana*); Orange, Sour (*Citrus aurantium*); Orange, Sweet (*Citrus sinensis*); Pummelo (*Citrus maxima*); Russell River Lime (*Microcitrus inodora*); Satsuma Mandarin (*Citrus unshiu*); Sweet Lime (*Citrus limetta*); Tachibana Orange (*Citrus tachibana*); Tahiti Lime (*Citrus latifolia*); Tangelo (*Citrus x tangelo*); Tangerine (Mandarin) (*Citrus reticulata*); Tangor (*Citrus nobilis*); Trifoliolate Orange (*Poncirus trifoliata*); Uniq Fruit (*Citrus aurantium* Tangelo group); cultivars, varieties and/or hybrids of these.

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Albinism <i>(Alternaria alternata pv citri)</i> Alternaria Leaf and Fruit Spot <i>(Alternaria citri)</i> Anthracnose <i>(Colletotrichum acutatum,</i> <i>C. gloeosporioides)</i> Cercospora Leaf Spot <i>(Cercospora spp.)</i> Diplodia Stem-End Rot <i>(Diplodia natalensis)</i> Greasy Spot <i>(Mycosphaerella citri)</i> Melanose <i>(Diaporthe citri)</i> Penicillium Decays Green Mold, Whisker Mold,	12.0-15.5 fl. oz. (0.20-0.25 lb. ai/A)	Begin applications prior to disease development and continue throughout the season on 7- to 21- day intervals following the resistance management guidelines. Under conditions that favor severe disease epidemics, use the higher application rates. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. Use a horticultural spray oil to improve control of greasy spot.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11. <b>DO NOT</b> make more than four (4) applications of AG37589 CS SC Fungicide or other Group 11 fungicide per season.

<p>Suppression of Blue Mold (<i>Penicillium</i> spp.)  Phomopsis Stem-End Rot (<i>Phomopsis citrii</i>)  Post Bloom Fruit Drop (PFD) (<i>Colletotrichum acutatum</i>)  Powdery Mildew (<i>Erysiphe</i> spp.)  Scab (<i>Elsinoe fawcettii</i>)  Sweet Orange Scab (<i>Elsinoe australis</i>)</p>		
<p>Black Spot (<i>Guidnardia citricarpa</i>)</p>	<p>9.0-15.5 fl. oz. (0.15-0.25 lb. ai/A)</p>	
<p>Pummelo  Citrus Hybrid (Uniq fruit only)</p> <p><b>Soilborne Diseases</b>  Seedling Root Rot, Basal Stem Rot  (<i>Rhizoctonia solani</i>)</p>	<p>0.40-0.80 fl oz/1000 row feet (0.0065-0.013 lb ai/1000 row feet)</p>	<p>For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.</p>
<p><b>RESTRICTIONS</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 10 applications per year at the low rate (9.0 fl oz/A). When applying at 12.0 fl oz/A, <b>DO NOT</b> apply more than 7 applications per year.</li> <li>● <b>DO NOT</b> use AG37589 CS SC Fungicide in citrus plant propagation nurseries.</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**CLOVER (AND STANDS CONTAINING CLOVER) (See Nongrass Animal Feeds Forage, Fodder, Straw and Hay)**

**CORN Field Pop Sweet (Includes Seed Production)**

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Rust ( <i>Puccinia sorghi</i> )	6.0-9.0 fl. oz. (0.10-0.15 lb. ai/A)	For gray leaf spot, apply AG37589 CS SC Fungicide at the onset of disease. A second application may be required 14 days later if disease pressure persists.
Anthracnose Leaf Blight ( <i>Colletotrichum graminicola</i> ) Eye Spot ( <i>Aureobasidium zeae</i> ) Gray Leaf Spot ( <i>Cercospora sorghi</i> ) Northern Corn Leaf Blight ( <i>Setosphaeria turcica</i> ) Northern Corn Leaf Spot ( <i>Cochliobolus carbonum</i> ) Physoderma Brown Spot ( <i>Physoderma maydis</i> ) Southern Corn Leaf Blight ( <i>Cochliobolus heterostrophus</i> ) Southern Rust ( <i>Puccinia polyspora</i> )	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	For all other diseases, begin applications prior to disease development and may continue throughout the season every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11. For field corn and field corn grown for seed, <b>DO NOT</b> make more than two (2) applications per season.
Early Application (V4 – V8)	6.0 fl. oz. (0.10 lb. ai/A)	AG37589 CS SC Fungicide may be applied early (V4 – V8) for early season disease control and beneficial physiological benefits. If mixing with herbicides, other than solo glyphosate products, or mesotrione products, consult your local Aceto Life Sciences, L/L/C d/b/a Actylis representative.
<b>Soilborne Diseases</b> Rhizoctonia Root and Stalk Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet (0.0065-0.013 lb ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section
<p><b>RESTRICTIONS</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 120 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 7 applications per year at the high rate (15.5 fl oz/A) or 20 applications per year at the low rate (6.0 fl oz/A). When applying at 9.0 fl oz/A, <b>DO NOT</b> apply more than 13 applications per year.</li> <li>● <b>DO NOT</b> apply within 7 days of harvest (7-day PHI).</li> </ul>		

**COTTON**

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
<p>Alternaria Leaf Spot (<i>Alternaria</i> spp.) Anthracnose (<i>Glomerella gossypii</i>) Areolate Mildew (<i>Ramularia gossypii</i>) Ascochyta Blight (<i>A. gossypii</i>) Boll Rots (<i>Ascochyta gossypii</i>, <i>Alternaria</i> spp., <i>Diplodia</i> spp., <i>Phoma</i> spp.) Cotton Rust (<i>Puccinia schedonnardi</i>) Diplodia Boll Rot (<i>Diplodia</i> spp.) Hardlock (<i>Fusarium verticillioides</i>) Leaf Spots and Blights (<i>Alternaria</i> spp., <i>Ascochyta gossypii</i>, <i>Cercospora</i> spp., <i>Stemphyllium</i> spp.) Southwestern Cotton Rust (<i>Puccinia cacabata</i>) (<i>Puccinia</i> spp.) Stemphyllium Leaf Spot (<i>Stemphyllium</i> spp.) Target spot (<i>Corynespora</i> <i>cassicola</i>)</p>	<p>6.0-9.0 fl. oz. (0.1-0.15 lb. ai/A)</p>	<p>For optimum disease control, begin applications prior to or in the early stages of disease development. Applications may be made by ground, air, or chemigation. An adjuvant may be added at specified rates. Minimum application volumes for air and ground are 5 and 10 gallons per acre, respectively.</p> <p>Target the first AG37589 CS SC Fungicide application at approximately pinhead square to first bloom to protect the plant from diseases. Subsequent application(s) are specified on a 14- to 21-day schedule. An additional application may be made depending on environmental conditions and the health of the cotton plant.</p> <p>Under poor environmental conditions conducive to seedling disease and poor cotton growth, AG37589 CS SC Fungicide may be applied to early season cotton to suppress damping off and other diseases which result in plant stand loss.</p> <p><b>DO NOT</b> apply more than two foliar applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternating with a fungicide that has a different mode of action. <b>DO NOT</b> make more than three (3) foliar applications of AG37589 CS SC Fungicide or other Group 11 fungicides per crop per acre per year.</p>
<p>Pythium Seedling Blight (<i>Pythium aphanidermatum</i>) Rhizoctonia Seedling Blight (<i>Rhizoctonia solani</i>)</p>	<p><b>In-Furrow</b> 0.40-0.80 fl oz product per 1000 row feet (0.0065- 0.013 lb. ai/1000 row feet)</p>	<p><b>AG37589 CS SC Fungicide Application Directions:</b> Apply AG37589 CS SC Fungicide as an in-furrow spray in 3-7 gallons of water at planting. Mount the spray nozzle so the spray is directed into the furrow just before the seed are covered. Use the higher rate when the weather conditions are expected to be conducive for disease development, if the field has a history of Pythium problems, or if minimum/low till programs are in place.</p> <p>See the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section for table illustrating total fluid ounces per acre with various row spacings.</p>
<p><b>RESTRICTIONS</b></p> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>• <b>DO NOT</b> apply at less than 14-day interval.</li> <li>• <b>DO NOT</b> apply more than 27 fl oz of product/crop/year as a foliar spray.</li> <li>• <b>DO NOT</b> apply more than 3 applications per year at the high rate (9.0 fl oz/A) or 4 applications per year at the low rate (6.0 fl oz/A).</li> <li>• AG37589 CS SC Fungicide may be applied up to 45 days before harvest (45-day PHI).</li> </ul>		

**CUCURBITS, CROP GROUP 9** Cantaloupe, Chayote, Chinese-Waxgourd, Cucumber, Gourds, Honeydew, Melons, *Momordica* spp.(bitter melon, balsam apple), Muskmelon, Watermelon, Pumpkin,Squash, Zucchini, cultivars and.or hybrids of these.

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Alternaria Blight <i>(Alternaria cucumerina)</i> Anthracnose <i>(Colletotrichum lagenarium)</i> Belly Rot ( <i>Rhizoctonia solani</i> ) Cercospora Leaf Spot <i>(Cercospora citrulina)</i> Downy Mildew <i>(Pseudoperonospora cubensis)</i> Gummy Stem Blight <i>(Didymella bryoniae)</i> Leaf Spots ( <i>Alternaria</i> spp., <i>Cercospora</i> spp.) Myrothecium Canker <i>(Myrothecium roridum)</i> Plectosporium Blight <i>(Plectosporium tabacinum)</i> Powdery Mildew <i>(Sphaerotheca fuliginea, Erysiphe cichoracearum)</i> Target Leaf Spot <i>(Corynespora cassicola)</i> Ulocladium Leaf Spot <i>(Ulocladium cucurbitae)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	<p>For both downy and powdery mildew, make preventative applications on a 5- to 7-day schedule. For belly rot control, make the first application at the 1-3 leaf crop stage with a second application just prior to vine tip over or 10-14 days later whichever occurs first.</p> <p>For all other diseases, begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season every 7-14 days following the resistance management guidelines.</p> <p>Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> tank mix AG37589 CS SC Fungicide with crop oil concentrates (COC), methylated spray oil (MSO) or silicon adjuvants.</p> <p><b>DO NOT</b> tank mix AG37589 CS SC Fungicide with malathion, dicofol, endosulfan, methomyl, potassium salt of fatty acids, &amp; 2,6-dichloro-4-nitroaniline.</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p><b>DO NOT</b> make more than four (4) foliar applications of AG37589 CS SC Fungicide or other Group 11 fungicides per crop per acre per year.</p>
<b>Soilborne Diseases</b> Rhizoctonia Root Rot <i>(Rhizoctonia solani)</i>	0.40-0.80 fl oz/1000 row feet (0.0065- 0.013 lb ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<b>RESTRICTIONS</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 5-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year.               <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>DO NOT</b> apply within 1 day of harvest (1-day PHI).</li> </ul>		

**FRUITING VEGETABLES CROP GROUP 8-10**

**PEPPER: Bell Pepper, Non-Bell Pepper, Sweet Non-Bell, Pepper  
EGGPLANT, ORKA, PEPINO**

**Including all cultivars and/or hybrids of these**

**See specific directions for use for Tomatoes.**

**Complete List of Fruiting Vegetables:** African Eggplant; Bell Pepper; Eggplant; Martynia; Nonbell Pepper; Okra; Pea Eggplant; Pepino; Roselle; Scarlet Eggplant; cultivars, varieties; and/or hybrids of these.

<b>DISEASE CONTROLLED</b>	<b>RATE AG37589 CS SC Fungicide/ACRE</b>	<b>APPLICATION INSTRUCTIONS</b>
Anthracnose ( <i>Colletotrichum</i> spp.) Powdery Mildew ( <i>Sphaerotheca</i> spp.)	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Begin applications prior to disease development and continue throughout the season on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>Soilborne Diseases</b> Rhizoctonia Seedling Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet (0.0065- 0.013 lb ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 60 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.0 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 3 applications per year at the high rate (15.5 fl oz/A) or 10 applications per year at the low rate (6.0 fl oz/A).</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**GRAPES and OTHER SMALL FRUIT VINE CLIMBING SUBGROUP 13-07F (Except fuzzy kiwifruit)** Amur River, Grape, Kiwifruit, Hardy, Maypop, Muscadines, Schisandra Berry, Including all cultivars and/or hybrids of these

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
<p>Black Rot (<i>Guignardia bidwellii</i>) Downy Mildew (<i>Plasmopara viticola</i>) Phomopsis Cane and Leaf Spot (<i>Phomopsis viticola</i>) Powdery Mildew (<i>Uncinula necator</i>)</p> <p><b>Suppression Only:</b> Botrytis Bunch Rot (<i>Botrytis cinerea</i>)</p>	<p>10.0-15.5 fl. oz. (0.16-0.25 lb. ai/A)</p>	<p>Begin applications prior to disease development and continue throughout the season every 10-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> apply more than two sequential foliar applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternating with a fungicide that is not in Group 11.</p> <p style="text-align: center;"><b>ATTENTION</b></p> <p>AG37589 CS SC Fungicide is extremely phytotoxic to certain apple varieties.</p> <p>AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit)</p> <p><b>DO NOT</b> spray AG37589 CS SC Fungicide where spray drift may reach apple trees.</p> <p><b>DO NOT</b> use spray equipment which has been previously used to apply AG37589 CS SC Fungicide to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.</p> <p>AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 10-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 9 applications per year at the low rate (10.0 fl oz/A).</li> <li>● <b>DO NOT</b> apply within 14 days of harvest (14-day PHI).</li> </ul>		

## GRASSES (GROWN FOR SEED)

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Ergot Stem Diseases Powdery Mildew ( <i>Erysiphe graminis</i> ) Rust ( <i>Puccinia</i> spp.)	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	<p>Begin applications prior to disease development and continue throughout the season on a 10- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 10-day interval.</li> <li>● <b>DO NOT</b> apply more than 48 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.8 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 3 applications per year at the high rate (15.5 fl oz/A) or 8 applications per year at the low rate (6.0 fl oz/A).</li> <li>● AG37589 CS SC Fungicide may be applied up to 8 days prior to harvest (swathing) (8- day PHI).</li> </ul>		

**HERBS & SPICES (EXCEPT BLACK PEPPER) CROP GROUP 19** Allspice; Angelica; Anise (seed); Anise, star; Annatto; Balm; Basil; Borage; Burnet; Camomile; Caper (buds); Caraway; Caraway, Black; Cardamon; Cassia (buds); Catnip; Celery Seed; Chervil (dried); Chive; Chive, Chinese; Cinnamon; Clary; Clove (buds); Coriander (cilantro or Chinese parsley) (leaf); Coriander (seed); Costmary; Culantro (leaf and seed); Cumin; Curry (leaf); Dill (seed); Dillweed; Fennel, Common; Fennel, Florence (seed); Fenugreek; Grains of Paradise; Horehound; Hyssop; Juniper (berry); Lavender; Lemongrass; Lovage (leaf and seed); Mace; Marigold; Marjoram; Mustard (seed), Nasturtium; Nutmeg; Parsley (dried); Pennyroyal; Pepper, White; Poppy Seed; Rosemary; Rue; Saffron; Sage; Savory, Summer and Winter Sweet Bay; Tansy; Tarragon; Thyme; Vanilla; Wintergreen; Woodruff; Wormwood

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Corynespora Blight ( <i>Corynespora cassicola</i> ) Dill Blight ( <i>Cercosporidium punctum</i> ) Phoma Blight ( <i>Passalora puncta</i> )	6.0-15.5 fl. oz. (0.10- 0.25 lb. ai/A)	Begin AG37589 CS SC Fungicide applications at the onset of disease development and continue throughout the season on a 7-day schedule, following the resistance management guidelines. Applications may be made by ground only. An adjuvant may be added at specified rates. Use a minimum of 30 gallons of water per acre.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>WASABI</b> Fusarium Rhizome and Root Rot ( <i>Pythium</i> spp.)	6.2-15.4 fl. oz. (0.10- 0.25 lb. ai/A)	Begin AG37589 CS SC Fungicide applications at the onset of disease development and continue throughout the season on a 7-day schedule, following the resistance management guidelines. Applications may be made by ground or through the irrigation system (chemigation). An adjuvant may be added at specified rates. Use a minimum of 30 gallons of water per acre.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A).</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**Leafy Vegetables (except Brassica), Crop Group 4** Amaranth, Arugula, Cardoon, Celery, Celtuce, Chervil, Chrysanthemum, Edible, Corn Salad, Cress, Dandelion, Dock, Endive, Fennel, Lettuce, Head and Leaf, Orach, Parsley, Purslane, Radicchio, Rhubarb, Spinach, Swiss, Chard, Including cultivars and/or hybrids of these

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
<p><b>Foliar Diseases</b>            Alternaria Leaf Spot  <i>(Alternaria sonchi, A. spp.)</i>            Anthracnose  <i>(Microdochium panattonianum, Colletotrichum dematium)</i>            Ascochyta Leaf Spot  <i>(Ascochyta spp.)</i>            Cercospora Leaf Spot  <i>(Cercospora spp.)</i>            Rust  <i>(Puccinia spp., Uromyces spp.)</i>            Septoria Leaf Spot  <i>(Septoria petroselini)</i>            White Rust <i>(Albugo occidentalis)</i></p>	<p>6.0-15.5 fl. oz.            (0.10-0.25 lb.            ai/A)</p>	<p>For both downy and powdery mildew, make preventative applications on a 5- to 7-day schedule.</p> <p>For all other diseases, begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p><b>ATTENTION:</b> Applications of AG37589 CS SC Fungicide to leafy vegetable foliage have contributed to phytotoxicity under certain circumstances. Proceed with caution with regard to tank mixes and adjuvants when treating all leafy vegetables with AG37589 CS SC Fungicide. AG37589 CS SC Fungicide must not be tank mixed on leaf lettuce with permethrin, fosetyl-AL, lambda-cyhalothrin products, or another product that may increase the penetration of AG37589 CS SC Fungicide into the leaf surface, including, but not limited to, silicone wetters.</p>
<p>Downy Mildew <i>(Bremia lactucae)</i>            Powdery Mildew  <i>(Erysiphe cichoracearum)</i></p>	<p>12.0-15.5 fl. oz.            (0.20-0.25 lb.            ai/A)</p>	
<p><b>Soilborne Diseases</b>            Webb Blight,            Bottom Rot,            Crater Rot, Root Rot  <i>(Rhizoctonia solani)</i></p>	<p>0.40-0.80 fl            oz/1000 row            feet (0.0065-            0.013 lb            ai/1000 row            feet)</p>	<p>For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 5-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year.               <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A). When applying at 12.0 fl oz/A, <b>DO NOT</b> apply more than 7 applications per year.</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**Legume Vegetables, Dry and Succulent, Crop Group 6 and Legume Vegetables, Foliage of any Cultivar of Bean (*Phaseolus* spp.) and Field Pea (*Pisum* spp.), Crop Group 7**

Bean (*Lupinus* spp.) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin)  
 Bean (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)  
 Bean (*Vigna* spp.) (includes adzuki bean, asparagus bean, blackeyed pea, cowpea, catjang, Chinese longbean, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean)  
 Bean (Glycine max) Soybean, Immature Seed (edamame)  
 Broad bean (fava bean) (*Vicia faba*)  
 Chickpea (garbanzo bean) (*Cicer arietinum*)  
 Guar (*Cyamopsis tetragonoloba*)  
 Jackbean (*Canavalia ensiformis*)  
 Lablab Bean (hyacinth bean) (*Lablab purpureus*)  
 Lentil (*Lens esculenta*)  
 Pea (*Pisum* spp.) (includes dwarf pea, edible- pod pea, English pea, garden pea, green pea, field pea, snow pea, sugar snap pea)  
 Pigeon Pea (*Cajanus cajan*)  
 Sword Bean (*Canavalia gladiata*)

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Bean Rust ( <i>Uromyces appendiculatus</i> )	6.0 fl. oz. (0.10 lb. ai/A)	Begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season every 7-14 days following the resistance management guidelines. Use the higher rates under severe disease pressure.
Alternaria Blight ( <i>Alternaria</i> spp.) Alternaria Leaf Spot ( <i>Alternaria alternata</i> ) Anthracnose ( <i>Colletotrichum lindemuthianum</i> ) Ascochyta Blight ( <i>Mycosphaerella pinodes</i> ) Ascochyta Leaf and Pod Spot ( <i>Ascochyta</i> spp.) Ascochyta Leaf Spot ( <i>Ascochyta phaseolorum</i> ) Rust ( <i>Phakopsora</i> spp.) Southern Blight ( <i>Sclerotium rolfsii</i> ) Web Blight ( <i>Rhizoctonia solani</i> )	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. For rust, use of a non-ionic surfactant is advised.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.

<p><b>Soilborne Diseases</b> Rhizoctonia Root Rot (<i>Rhizoctonia solani</i>)</p>	<p>0.40-0.80 fl oz/1000 row feet (0.0065- 0.013 lb ai/1000 row feet)</p>	<p>For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.</p> <p>AG37589 CS SC Fungicide can be applied to the furrow and covering soil at planting time in a 7-inch band. Avoid a concentrated stream directly on the seed or delayed emergence may occur.</p> <p>If using a narrow spray as an in-furrow spray, adjust the spray stream to hit the soil next to the seed but not hit the seed.</p> <p>NOTE: Conduct a seed safety test with your crop before making in-furrow applications.</p>
---	--	--

**RESTRICTIONS:**

- **DO NOT** exceed the maximum rate listed in this crop table.
- **DO NOT** apply at less than 7-day interval.
- **DO NOT** apply more than 90 fl oz of product/A/year.
  - **DO NOT** apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- **DO NOT** apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A).
- **Pre-Harvest Interval (PHI):**
  - **DO NOT** apply within 14 days of harvest (14-day PHI) of dry legume vegetables (dry bean and dry pea seeds).
  - AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI) for succulent beans and peas.
- For use on soybeans, please refer to the soybean crop directions for use.

**MINT (FRESH OR FOR PROCESSING INTO MINT OIL)**

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Leaf Spot <i>(Ramularia spp.)</i> <i>(Alternaria spp.)</i> <i>(Phoma, spp.)</i> Powdery mildew <i>(Erysiphe spp.)</i> Rust <i>(Puccinia menthae)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season on a 7- to 10-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>Soilborne Diseases</b> Seedling Root Rot, Basal Stem Rot <i>(Rhizoctonia solani)</i>	0.40-0.80 fl oz/1000 row feet (0.0065- 0.013 lb ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 42 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.75 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 2 applications per year at the high rate (15.5 fl oz/A) or 7 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>Pre-Harvest Interval (PHI):</b> <ul style="list-style-type: none"> <li>○ For processed mint, <b>DO NOT</b> apply within 7 days of harvest (7-day PHI).</li> <li>○ For fresh mint, AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul> </li> </ul>		

**Nongrass Animal Feeds Forage, Fodder, Straw and Hay, Crop Group 18** For pure/mixed stands of the following or stands mixed with grasses:

Alfalfa (*Medicago sativa* subsp. *sativa*), Bean, Velvet (*Mucuna pruriens* var. *utilis*), Clover (*Trifolium* spp., *Melilotus* spp.), Kudzu (*Pueraria lobata*), Lespedeza (*Lespedeza* spp.), Lupin (*Lupinus* spp.), Sainfoin (*Onobrychis viciifolia*), Trefoil (*Lotus* spp.), Vetch (*Vicia* spp.), Vetch, Crown (*Coronilla varia*), Vetch, Milk (*Astragalus* spp.)

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Alternaria Leaf Spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum trifolii</i> ) Black Patch ( <i>Rhizoctonia leguminicola</i> ) Cercospora Leaf Spot ( <i>Cercospora</i> spp.) Common Leaf Spot ( <i>Pseudopezizza solani</i> ) Downy Mildew ( <i>Peronospora</i> spp.) Leaf Spot ( <i>Leptosphaerulina briosiai</i> ) Powdery Mildew ( <i>Oidium</i> spp., <i>Erysiphe</i> spp.) Rhizoctonia and Stem Blight ( <i>Rhizoctonia solani</i> )  Rust ( <i>Phakopsora</i> spp.) ( <i>Uromyces</i> spp.) Spring Black Stem and Leaf Spot ( <i>Phoma medicaginis</i> ) Stagonospora Leaf Spot ( <i>Stagonospora meliloti</i> ) Stemphyllium Leaf Spot ( <i>Stemphyllium</i> spp.) Summer Black Stem and Leaf Spot ( <i>Cercospora medicaginis</i> ) Yellow Leaf Blotch ( <i>Leptotrichilia medicaginis</i> )	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	<p>Begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season. Use the higher rates under severe disease pressure. Applications may be made by ground, air or chemigation. Use of an additive including crop oil concentrate or non- ionic surfactant is advised.</p> <p>For management of outbreaks of Asian soybean rust and other Puccinia species on alternate host species including kudzu, lespedeza, trefoil and vetch, apply AG37589 CS SC Fungicide to forages grown in the vicinity of soybeans and other legume crops (beans and peas) as a part of an Asian rust disease management strategy. Consult with local experts and university extension agents for the latest advice.</p> <p><b>DO NOT</b> apply more than three sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>

Sclerotinia Crown Rot and Wilt on Clover <i>(Sclerotinia trifoliorum)</i>	10.0 fl. oz. (0.17 lb. ai/A)	
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply more than 0.25 lb ai/A per cutting.</li> <li>● <b>DO NOT</b> apply at less than 14-day interval.</li> <li>● <b>DO NOT</b> apply more than 42 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.75 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 2 applications per year at the high rate (15.5 fl oz/A) or 7 applications per year at the low rate (6.0 fl oz/A). When applying at 10.0 fl oz/A, <b>DO NOT</b> apply more than 4 applications per year.</li> <li>● <b>DO NOT</b> apply within 14 days of grazing or harvest (14-day PHI) for forage and hay.</li> <li>● Not for use on rangeland.</li> </ul>		

**Oilseed Crops Crop Group 20** Crambe Flax

Mustard, Indian Mustard, Field Mustard, Black Rapeseed Rapeseed, Indian Safflower Sunflower

Including all cultivars and/or hybrids of these.

**Complete List of Oilseed Crops:** Borage; Calendula; Castor Oil Plant; Chinese Tallowtree; Cottonseed; Crambe; Cuphea; Echimium; Euphorbia; Evening Primrose; Flax Seed; Gold of Pleasure; Hare’s Ear Mustard; Jojoba; Lesquerella; Lunaria; Meadowfoam; Milkweed; Mustard Seed; Niger Seed; Oil Radish; Poppy Seed; Rapeseed; Rose Hip; Safflower; Sesame; Stokes Aster; Sunflower; Sweet Rocket; Tallowwood; Tea Oil Plant; Vernonia; cultivars, varieties, and/or hybrids of these

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Alternaria Leaf Spot <i>(Alternaria spp.)</i> Downy Mildew <i>(Plasmopora halstedii,</i> <i>Plasmopora helianthi)</i> Pasmio <i>(Septoria linicola grass)</i> Sunflower Rust <i>(Puccinia helianthi)</i>	6.0-15.5 fl. oz. (0.1-0.25 lb. ai/A)	Apply 6.0 fl oz of AG37589 CS SC Fungicide at early bud followed by 14.0 fl oz at about 45 days before harvest. A third application of 7.0 fl oz may be made 30 days before harvest. Applications may be made by ground, air or chemigation. Use a minimum of 10 gallons of water per acre for ground applications.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 14-day interval.</li> <li>● <b>DO NOT</b> apply more than 24 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.45 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 1 applications per year at the high rate (15.5 fl oz/A) or 4 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>DO NOT</b> apply within 30 days of harvest (30-day PHI).</li> </ul>		

PEANUTS

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
<p><b>Soilborne Diseases – early season (in-furrow application)</b> Aspergillus Crown Rot (<i>Aspergillus niger</i>) Pythium Damping Off (<i>Pythium</i> spp.) Stem Rot/White Mold Suppression (<i>Sclerotium rolfsii</i>)</p>	<p>0.40-0.80 fl oz/1000 row feet (0.0065-0.013 lb ai/1000 row feet)</p>	<p>Apply AG37589 CS SC Fungicide in-furrow at planting for control of various seed/seedling diseases including early season suppression of stem rot. See directions and rates under <b>PRODUCT INFORMATION</b> section.</p>
<p><b>Soilborne Diseases – mid-late season</b> Rhizoctonia Peg and Pod Rot (<i>Rhizoctonia solani</i>) Stem Rot/White Mold (<i>Sclerotium rolfsii</i>)</p> <p><b>Suppression Only:</b> Cylindrocladium Black Rot (<i>Cylindrocladium crotalariae</i>) Pythium Pod Rot (<i>Pythium myriotylum</i>)</p>	<p>12.0-24.5 fl. oz. (0.20-0.40 lb. ai/A)</p>	<p>Apply AG37589 CS SC Fungicide at approximately 60 and 90 days after planting as a foliar application. This application regime may be applied earlier in the season if environmental conditions favor disease development.</p> <p>These two applications of AG37589 CS SC Fungicide will provide protection against the soilborne diseases and will also provide control of the foliar diseases listed for a 10- to 14-day period after each spray. Under heavy disease pressure and/or where there is high rainfall and/or irrigation, use 18.5-24.5 fl oz/A. For light disease pressure and dry environmental conditions (non-irrigated, low rainfall), use 12.0-24.5 fl oz/A. For control of Pythium, a rate of 24.5 fl oz/A is required. Additional applications of other fungicides on a leaf spot application schedule will be required to provide season-long disease control of the leaf spot diseases.</p> <p>Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p>
<p><b>Foliar Diseases</b> Early Leaf Spot (<i>Cercospora arachidicola</i>) Late Leaf Spot (<i>Cercosporidium personatum</i>) Rust (<i>Puccinia arachidis</i>) Web Blotch (<i>Phoma arachidicola</i>)</p>	<p>6.0-18.5 fl. oz. (0.10-0.30 lb. ai/A)</p>	<p>For foliar disease control only, a lower rate of AG37589 CS SC Fungicide may be applied on a 10- to 14- day interval.</p> <p><b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>

**RESTRICTIONS:**

- **DO NOT** exceed the maximum rate listed in this crop table.
- **DO NOT** apply at less than 10-day interval.
- **DO NOT** apply more than 49 fl oz of product/A/year.
  - **DO NOT** apply more than 0.8 lb ai/A/year of azoxystrobin-containing products.
- **DO NOT** apply more than 2 applications per year at the high rate (24.5 fl oz/A) or 8 applications per year at the low rate (6.0 fl oz/A). When applying at 12.0 fl oz/A, **DO NOT** apply more than 4 applications per year. When applying at 18.5 fl oz/A, **DO NOT** apply more than 2 applications per year.
- **DO NOT** apply within 14 days of harvest (14-day PHI).

## PISTACHIOS

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Alternaria Late Blight <i>(Alternaria alternata)</i> Botryosphaeria Panicle and Shoot Blight <i>(Botryosphaeria dothidea)</i> Septoria Leaf Spot <i>(Septoria pistaciarum)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	Begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season on 7- to 21-day intervals following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year.               <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>DO NOT</b> apply within 7 days of harvest (7-day PHI).</li> </ul>		

## POTATOES

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Black Dot ( <i>Colletotrichum coccodes</i> ) Early Blight ( <i>Alternaria solani</i> ) Late Blight ( <i>Phytophthora infestans</i> ) Powdery Mildew ( <i>Erysiphe cichoracearum</i> )	6.0-20.0 fl. oz. (0.10- 0.33 lb. ai/A)	<p><b>Early blight</b> - For a 7-day application schedule, use AG37589 CS SC Fungicide 6.2 fl oz product/A. For a 14-day application schedule, use the 12.0 fl oz product/A rate.</p> <p><b>Late blight</b> - Apply AG37589 CS SC Fungicide at 12.0 fl oz product/A on a 7-day schedule. Initiate late blight applications in a preventative schedule prior to disease development according to local practices. If late blight symptoms develop or conditions favor disease, switch immediately to a non-Group 11 fungicide, using a 5-day schedule. Addition of a spreader/sticker may improve coverage.</p> <p>For all other diseases, begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season every 7-14 days following the resistance management guidelines. Use the high rate and the shorter interval if disease epidemics are severe. Applications may be made by ground, air or chemigation.</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
<p><b>Soilborne Diseases</b></p> Black Dot ( <i>Colletotrichum coccodes</i> ) Black Scurf ( <i>Rhizoctonia solani</i> ) Silver Scurf ( <i>Helminthosporium solani</i> )	0.40-0.80 fl oz/1000 row feet (0.0065- 0.013 lb ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 120 fl oz of product/A/year.               <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 6 applications per year at the high rate (20.0 fl oz/A) or 20 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>DO NOT</b> apply within 14 days of harvest (14-day PHI).</li> </ul>		

## QUINOA

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Leaf Spot ( <i>Ascochyta hyalospora</i> ) Stalk Rot ( <i>Phoma exigua</i> )	12 fl. oz. (0.20 lb. ai/A)	Apply prior to disease development.  An adjuvant may be added at specified rates.
<b>Application:</b> AG37589 CS SC Fungicide can be applied by either ground, chemigation, or aerial application.		
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 14-day intervals.</li> <li>● <b>DO NOT</b> apply more than 24 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.40 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● When applying at 12 fl oz/A, <b>DO NOT</b> apply more than 2 applications per year.</li> <li>● <b>Pre-Harvest Interval (PHI):</b> <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply within 7 days (7-day PHI) for forage and hay.</li> <li>○ <b>DO NOT</b> apply within 14 days of grazing (14-day PHI).</li> <li>○ <b>DO NOT</b> apply within 30 days of harvest (30-day PHI).</li> </ul> </li> </ul>		

# RICE

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
<b>Sheath/Stem Diseases</b> Sheath Blight <i>(Rhizoctonia solani)</i>	6.0-18.5 fl. oz. (0.10-0.30 lb. ai/A)	Apply AG37589 CS SC Fungicide prior to disease development. Applications may be made by ground, air or chemigation. For aerial application, use volumes of 5-10 GPA. An adjuvant may be added at specified rates.
Aggregate Sheath Spot <i>(Ceratobasidium oryzae-sativae = Rhizoctonia oryzae-sativae)</i> Black Sheath Rot <i>(Gaeumannomyces graminis var. graminis)</i> Sheath Spot <i>(Rhizoctonia oryzae)</i> Stem Rot <i>(Magnaporthe salvinii = Sclerotium oryzae = Nakateae sigmoidea)</i>	9.0-18.5 fl. oz. (0.15 - 0.30 lb. ai/A)	<p>For sheath blight control, application rates may vary from 9.0 to 12.0 fl oz/A depending on the growth stage of the rice and the severity of the disease. Consult with your local extension personnel on sheath blight control.</p> <p>For other stem/sheath diseases including stem rot, black sheath rot, aggregate sheath spot and sheath spot, apply when disease is less than 4 inches above water line usually between panicle differentiation (PD) +5 days to PD +10 days or at initial sign of disease. Under heavy disease pressure and conditions favorable for disease development, a second application may be applied.</p> <p>For foliar and panicle diseases, apply AG37589 CS SC Fungicide prior to disease development. AG37589 CS SC Fungicide must be applied as a preventative treatment for blast control and applied prior to favorable conditions for blast development.</p> <p>For panicle blast, make an application at mid-boot to boot-split but prior to full head emergence. Apply a second application when panicles are approximately 60-90% emerged from the boot (7-14 days later).</p> <p>When AG37589 CS SC Fungicide is being applied for panicle blast on continuous rice acreage (no rotation to other crops), apply no more than two sequential foliar applications of AG37589 CS SC Fungicide or other Group 11 fungicides over multiple years before alternating with a fungicide with a different mode of action. <b>DO NOT</b> make more than two foliar applications of AG37589 CS SC Fungicide or other Group 11 fungicides per acre per season.</p>
<b>Foliar Diseases</b> Brown Leaf Spot <i>(Cochliobolus miyabeanus)</i> Leaf Smut <i>(Entyloma oryzae)</i> Narrow Brown Leaf Spot <i>(Cercospora janseana = Cercospora oryzae)</i>		
<b>Panicle Diseases</b> Kernel Smut <i>(Tilletia barclayana = Neovossia barclayana)</i> Panicle Blast <i>(Pyricularia grisea)</i>		

**RESTRICTIONS:**

- **DO NOT** exceed the maximum rate listed in this crop table.
- **DO NOT** treat rice fields used for aquaculture of fish and crustaceans.
- **DO NOT** apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Use care in making applications near non-target aquatic habitats.
- **DO NOT** apply at less than 7-day interval.
- **DO NOT** apply more than 42 fl oz of product/A/year.
  - **DO NOT** apply more than 0.70 lb ai/A/year of azoxystrobin-containing products.
- **DO NOT** apply more than 2 applications per year at the high rate (18.5 fl oz/A) or 7 applications per year at the low rate (6.0 fl oz/A). When applying at 9.0 fl oz/A, **DO NOT** apply more than 4 applications per year.
- **DO NOT** allow release of irrigation or flood water for at least 14 days after the last application.
- **DO NOT** apply within 28 days of harvest (28-day PHI).

## SORGHUM

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Anthracnose <i>(Colletotrichum graminicola)</i> Gray Leaf Spot <i>(Cercospora sorghi)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	<p>Begin AG37589 CS SC Fungicide applications prior to disease development. Use the high rates under conditions favorable for severe disease pressure, dense plant canopies, or when susceptible varieties are planted. Contact extension personnel for local economic thresholds and timings for specific diseases in your area. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
<b>Soilborne Diseases</b> Damping-Off <i>(Rhizoctonia solani, Pythium aphanadermatum)</i>	0.40-0.80 fl oz/1000 row feet (0.0065- 0.013 lb ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● For grain and stover, <b>DO NOT</b> apply more than 42 fl oz of product/A/year (0.75 lb ai/A/year of azoxystrobin-containing products).</li> <li>● For forage, <b>DO NOT</b> apply more than 30 fl oz of product/A/year (0.5 lb ai/A/year of azoxystrobin-containing products).</li> <li>● For grain and stover, <b>DO NOT</b> apply more than 2 applications per year at the high rate (15.5 fl oz/A) or 7 applications per year at the low rate (6.0 fl oz/A).</li> <li>● For forage, <b>DO NOT</b> apply more than 1 application per year at the high rate (15.5 fl oz/A) or 5 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>DO NOT</b> apply within 14 days of harvest (14-day PHI).</li> </ul>		

**SOYBEANS, SOYBEAN, IMMATURE SEED (Edamame)**

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/A CRE	APPLICATION INSTRUCTIONS
Aerial Blight <i>(Rhizoctonia solani)</i> Alternaria Leaf Spot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum truncatum)</i> Brown Spot <i>(Septoria glycines)</i> Cercospora Blight and Leaf Spot <i>(Cercospora kikuchii)</i> Frogeye Leaf Spot <i>(Cercospora sojina)</i> Pod and Stem Blight <i>(Diaporthe phaseolorum)</i> Rust <i>(Phakopsora spp.)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	<p>Begin AG37589 CS SC Fungicide applications prior to disease development. Use the high rates under conditions favorable for severe disease pressure, dense plant canopies, or when susceptible varieties are planted. Contact Extension personnel for local economic thresholds and timings for specific diseases in your area. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. Use of a crop oil concentrate or non-ionic surfactant with the lower use rate is advised.</p> <p>Soybean rust: AG37589 CS SC Fungicide may be used at 4 fl oz/A when tank mixed with a triazole registered for use on soybean rust.</p> <p><b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
<b>Soilborne Diseases</b> Rhizoctonia solani <i>(Rhizoctonia solani)</i> Southern blight <i>(Sclerotium rolfsii)</i>	0.40-0.80 fl oz/1000 row feet (0.0065-0.013 lb. ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 14-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> make more than one application at 15.5 fl oz product/acre or 0.25 lb ai/A to soybean forage and hay.</li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>Pre-Harvest Interval (PHI):</b> <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply within 14 days of harvest (14-day PHI) of soybeans (beans).</li> <li>○ AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI) to soybean forage and hay.</li> </ul> </li> </ul>		

**Stone Fruits, Crop Group 12-12**

Apricot Cherry, Sweet Cherry, Tart Nectarine Peach  
 Plum Plumcot Prune

**Complete List of Stone Fruit Crops:** Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these.

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Brown Rot Blossom Blight and Fruit Rot ( <i>Monilinia fructicola</i> , <i>M. laxa</i> )	12.0-15.5 fl. oz. (0.20-0.25 lb. ai/A)	For brown rot blossom blight, begin applications at early bloom and continue through petal fall. For brown rot on fruit, AG37589 CS SC Fungicide may be applied to fruit up to the day of harvest.  For scab, begin applications at petal fall and continue at 7- to 14-day intervals.
Scab ( <i>Cladosporium carpophilum</i> ) Alternaria Spot and Fruit Rot ( <i>Alternaria alternata</i> ) Anthracnose ( <i>Colletotrichum prunicola</i> , <i>C. gloeosporioides</i> ) Leaf Rust ( <i>Tranzschelia discolor</i> ) Powdery Mildew ( <i>Sphaerotheca pannosa</i> , <i>Podosphaera clandestina</i> ) Shot Hole ( <i>Wilsonomyces carpophilus</i> )	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	For all other diseases, begin application at the onset of disease as a protectant fungicide and continue on a 7- to 14-day schedule.  For peaches only, 9.0-15.5 fl oz of AG37589 CS SC Fungicide may be used for scab control.  Applications may be made by ground, air or chemigation.  DO NOT apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 90 fl oz of product/A/year.             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A). When applying at 12.0 fl oz/A, <b>DO NOT</b> apply more than 7 applications per year.</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**SUGAR BEETS**

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/AC RE	APPLICATION INSTRUCTIONS
<p><b>Foliar Diseases</b>                      Alternaria Leaf Spot                      (<i>Alternaria</i> spp.,  <i>A. alternata</i>)                      Ascochyta Leaf Spot                      (<i>Ascochyta cynarae</i>)                      Rust                      (<i>Uromyces betae</i>, <i>Puccinia helianthi</i>)                      White Rust                      (<i>Albugo tragopogonis</i>)</p>	<p>6.0-20.0 fl. oz.                      (0.10-0.33 lb. ai/A)</p>	<p>For powdery mildew, make preventative applications on a 5- to 7-day schedule. For all other diseases, begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
<p>Cercospora Leaf Spot                      (<i>Cercospora betae</i>,  <i>C. pastinaceae</i>)                      Powdery Mildew                      (<i>Erysiphe polygoni</i>,  <i>Leveillula taurica</i>)</p>	<p>9.0-15.5 fl. oz.                      (0.15-0.25 lb. ai/A)</p>	
<p><b>Soilborne Diseases</b>                      Circular Spot, Southern Blight                      (<i>Sclerotium rolfsii</i>)                      Pythium Root Rot                      (<i>Pythium aphanidermatum</i>)                      Rhizoctonia Stem Canker, Crown Rot                      (<i>Rhizoctonia solani</i>)</p>	<p>0.40-0.80 fl oz/1000 row feet                      (0.0065-0.013 lb ai/1000 row feet)</p>	<p>For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.</p> <p><b>For sugar beets</b> apply 3-7 inch banded applications in a minimum of 10 gallons per acre at the 2- to 8-leaf stage. <b>DO NOT</b> apply as a dribble application over the seed row. Tank mixtures of AG37589 CS SC Fungicide with crop oil concentrates (COC) or methylated spray oil (MSO) may result in crop injury. If cool soil conditions are expected after planting which could result in an extended period of plant emergence, <b>DO NOT</b> apply AG37589 CS SC Fungicide in-furrow. If using AG37589 CS SC Fungicide at the time of planting, <b>DO NOT</b> use a starter fertilizer with it.</p>

**RESTRICTIONS:**

- **DO NOT** exceed the maximum rate listed in this crop table.
- **DO NOT** apply at less than 5-day interval.
- **DO NOT** apply more than 120 fl oz of product/A/year.
  - **DO NOT** apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.
- **DO NOT** apply more than 6 applications per year at the high rate (20.0 fl oz/A) or 20 applications per year at the low rate (6.0 fl oz/A). When applying at 9.0 fl oz/A, **DO NOT** apply more than 13 applications per year. When applying at 15.5 fl oz/A, **DO NOT** apply more than 7 applications per year.
- Apply as an in-furrow spray in a minimum of 10 gallons per acre.
- AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).

# SUGARCANE

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Brown Rust <i>(Puccinia melanocephala)</i> Orange Rust <i>(Puccinia kuehnii)</i>	9.0-12.0 fl. oz. (0.15-0.20 lb. ai/A)	<p>Begin AG37589 CS SC Fungicide applications prior to rust development, and continue throughout the season every 14-28 days following resistance management guidelines.</p> <p>Scout fields and begin applications at the earliest sign of rust. An adjuvant may be used at specified rates. For ground applications, apply AG37589 CS SC Fungicide in sufficient water volume for adequate coverage and canopy penetration. Applications may be made by ground, air or chemigation.</p> <p><b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicide, before alternation with a fungicide that is not in Group 11. <b>DO NOT</b> make more than four foliar applications of AG37589 CS SC Fungicide or other Group 11 fungicide per acre per year.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 14-day interval.</li> <li>● <b>DO NOT</b> apply more than 48 fl oz of product/A/year.               <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.80 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 4 applications per year at the high rate (12.0 fl oz/A) or 5 applications per year at the low rate (9.0 fl oz/A).</li> <li>● <b>DO NOT</b> apply within 30 days of harvest (30-day PHI).</li> <li>● When applying by air, use no less than 5 gallons spray solution per acre.</li> </ul>		

**TI PALM, LEAVES, and ROOTS**

<b>DISEASE CONTROLLED</b>	<b>RATE AG37589 CS SC Fungicide/ACRE</b>	<b>APPLICATION INSTRUCTIONS</b>
<p><b>Foliar Diseases</b>                      Alternaria Leaf Spot  <i>(Alternaria spp., A. alternata)</i>                      Ascochyta Leaf Spot  <i>(Ascochyta cynarae)</i>                      Phyllostica Leaf Spot  <i>(Phyllostica spp.)</i>                      Rust  <i>(Uromyces betae, Puccinia helianthi)</i>                      White Rust  <i>(Albugo tragopogonis)</i></p>	<p>6.0-20.0 fl. oz.                      (0.10-0.33 lb. ai/A)</p>	<p>For powdery mildew, make preventative applications on a 5- to 7-day schedule. For all other diseases, begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p><b>DO NOT</b> apply more than six applications of AG37589 CS SC Fungicide per year for <i>Phyllostica</i> spp.</p>
<p>Cercospora Leaf Spot  <i>(Cercospora betae, C. pastinaceae)</i>                      Powdery Mildew  <i>(Erysiphe polygoni, Leveillula taurica)</i></p>	<p>9.0-15.5 fl. oz.                      (0.15-0.25 lb. ai/A)</p>	<p><b>DO NOT</b> apply more than eight applications of AG37589 CS SC Fungicide per year for <i>Cercospora</i> spp.</p>
<p><b>Soilborne Diseases</b>                      Circular Spot, Southern Blight  <i>(Sclerotium rolfsii)</i>                      Pythium Root Rot  <i>(Pythium aphanidermatum)</i>                      Rhizoctonia Stem Canker, Crown Rot  <i>(Rhizoctonia solani)</i></p>	<p>0.40-0.80 fl oz/1000 row feet (0.0065-0.013 lb ai/1000 row feet)</p>	<p>For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 5-day interval.</li> <li>● <b>DO NOT</b> apply more than 120 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 6 applications per year at the high rate (20.0 fl oz/A) or 20 applications per year at the low rate (6.0 fl oz/A). When applying at 9.0 fl oz/A, <b>DO NOT</b> apply more than 13 applications per year. When applying at 15.5 fl oz/A, <b>DO NOT</b> apply more than 7 applications per year.</li> <li>● Apply as an in-furrow spray in a minimum of 10 gallons per acre.</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

**TOBACCO**

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
<p>Blue Mold <i>(Peronospora tabacina)</i> Frogeye Leaf Spot <i>(Cercospora nicotianae)</i> Target Spot <i>(Rhizoctonia solani)</i></p>	<p>6.0-12.0 fl. oz. (0.1-0.2 lb. ai/A)</p>	<p>Begin AG37589 CS SC Fungicide applications prior to disease development or at first indication that blue mold is in the area. <b>DO NOT</b> apply AG37589 CS SC Fungicide as a curative application. If blue mold is present in the field, initiate applications with dimethomorph prior to an AG37589 CS SC Fungicide application. Apply on a 7- to 14-day interval with shorter intervals under conditions conducive to disease development. For ground applications, apply AG37589 CS SC Fungicide in sufficient water volume for adequate coverage and canopy penetration. For aerial application, use volumes of 10-15 GPA. Applications may be made by ground, air or chemigation. <b>DO NOT</b> apply AG37589 CS SC Fungicide on greenhouse seedlings. <b>DO NOT</b> tank mix with endosulfan. Tank mixing AG37589 CS SC Fungicide with insecticides formulated as emulsifiable concentrates (EC) or containing high amounts of solvents, may cause some crop injury.</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>NOTE: AG37589 CS SC Fungicide may enhance weather flecking on the leaves of certain tobacco types. This does not affect yield and quality.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 30 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.52 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 2 applications per year at the high rate (12.0 fl oz/A) or 5 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>DO NOT</b> apply within 21 days of harvest (21-day PHI).</li> </ul>		

**TOBACCO TRANSPLANTS IN GREENHOUSE GA, KY, IN, MD, MO, NC, OH, PA, SC, TN and VA only**

<b>DISEASE CONTROLLED</b>	<b>RATE AG37589 CS SC Fungicide/ACRE</b>	<b>APPLICATION INSTRUCTIONS</b>
Target Spot ( <i>Rhizoctonia solani</i> )	6.0 fl. oz. (0.1 lb. ai/A)	Apply 6 oz/A or 0.14 oz (4ml)/1000 sq ft in enough water for thorough coverage (5 gal/1000 sq ft advised). Make only one application prior to transplanting.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply more than 6 fl oz of product/A/year in the greenhouse. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.52 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● Make only one application in the greenhouse prior to transplanting.</li> </ul>		

**Tomatoes Tomatillos Subgroup 8-10A** Including all cultivars and/or hybrids of these. **Complete List of Tomato Crops:** Bush Tomato; Cocona; Currant Tomato; Garden Huckleberry; Goji Berry; Groundcherry; Naranjilla; Sunberry; Tomatillo; Tomato; Tree Tomato; cultivars, varieties, and/or hybrids of these.

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Anthracnose <i>(Colletotrichum coccodes)</i> Black Mold <i>(Alternaria alternata)</i> Buckeye Rot <i>(Phytophthora spp.)</i> Early Blight <i>(Alternaria solani)</i> Powdery Mildew <i>(Oidiopsis sicula)</i> Septoria Leaf Spot <i>(Septoria lycopersici)</i> Target Spot <i>(Corynespora cassiicola)</i>	5.0-6.2 fl. oz. (0.08-0.10 lb. ai/A)	<p>Begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season following the resistance management guidelines. For late blight, apply AG37589 CS SC Fungicide at 5- to 7-day intervals. For all other tomato diseases, apply AG37589 CS SC Fungicide on 7- to 21-day intervals. Applications may be made by ground, air or chemigation.</p> <p><b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Under certain weather conditions (particularly high temperatures) AG37589 CS SC Fungicide in combination with high rates of silicone-based or oil containing (petroleum or crop) additives or adjuvants may cause injury. <b>DO NOT</b> exceed 0.125% adjuvant (v/v). Consult an Aceto Life Sciences, L.L.C. d/b/a Actylis representative for more information concerning additives or adjuvants.</p>
Late Blight <i>(Phytophthora infestans)</i>	6.2 fl. oz. (0.10 lb. ai/A)	<p>A tank mixture with Dimethoate may cause crop injury.</p> <p>On fresh market tomatoes <b>DO NOT</b> use adjuvants or tank mix AG37589 CS SC Fungicide with any emulsifiable concentrate (EC) product.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 5-day interval.</li> <li>● <b>DO NOT</b> apply more than 35 fl oz of product/A/year.               <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.6 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 5 applications per year at the high rate (6.2 fl oz/A) or 7 applications per year at the low rate (5.0 fl oz/A).</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		



**TROPICAL FRUIT** Acerola Atemoya Avocado Biriba Canistel Cherimoya Custard Apple Dragon Fruit Feijoa Guava Ilima Jaboticaba Jackfruit Longan Loquat Lychee Mango Papaya Passionfruit Pawpaw Persimmon Pulasan Rambutan Sapodilla Sapote, Black Sapote, Mamey Sapote, White Soursop Star Apple Starfruit Sugar Apple Spanish Lime Tamarind

<b>DISEASE CONTROLLED</b>	<b>RATE AG37589 CS SC Fungicide/ACRE</b>	<b>APPLICATION INSTRUCTIONS</b>
Anthracnose <i>(Colletotrichum spp.)</i> Cercospora Leaf Spot <i>(Cercospora spp.)</i> Powdery Mildew <i>(Erysiphe spp.)</i> Rust <i>(Puccinia spp.)</i>	6.0-15.5 fl. oz. (0.10-0.25 lb. ai/A)	<p>Begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season on a 10- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p>Follow the resistance management guidelines in the Resistance Management Section. <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
<b>Soilborne Diseases</b> Seedling Root Rot, Basal Stem Rot <i>(Rhizoctonia solani)</i>	0.40-0.80 fl oz/1000 row feet (0.0065- 0.013 lb. ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<b>RESTRICTIONS:</b> <ol style="list-style-type: none"> <li>1) <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>2) <b>DO NOT</b> apply at less than 10-day interval.</li> <li>3) <b>DO NOT</b> apply more than 90 fl oz of product/A/year.               <ol style="list-style-type: none"> <li>a. <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ol> </li> <li>4) <b>DO NOT</b> apply more than 5 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A).</li> <li>5) AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ol>		

**VEGETABLES, LEAVES OF ROOT AND TUBER GROUP AND ROOT SUBGROUP** Beet, Garden<sup>1,2</sup> Burdock<sup>1,2</sup> Carrot<sup>1,2</sup> Cassava, Bitter and Sweet<sup>1</sup> Celeriac (celery root)<sup>1,2</sup> Chervil, Turnip-Rooted<sup>1,2</sup> Chicory<sup>1,2</sup> Dasheen (taro)<sup>1</sup> Ginseng<sup>2</sup> Horseradish<sup>2</sup> Parsley, Turnip-Rooted<sup>2</sup> Parsnip<sup>1,2</sup> Radish<sup>1,2</sup> Radish, Oriental(daikon)<sup>1,2</sup> Rutabaga<sup>1,2</sup> Salsify<sup>2</sup> Salsify, Black<sup>1,2</sup> Salsify, Spanish<sup>2</sup> Skirret<sup>2</sup> Sweet Potato<sup>1</sup> Tanier<sup>1</sup> Turnip<sup>1,2</sup> Yam, True<sup>1</sup>

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
<b>Foliar Diseases</b> Alternaria Leaf Spot <i>(Alternaria spp., A. alternata)</i> Ascochyta Leaf Spot <i>(Ascochyta cynarae)</i> Rust <i>(Uromyces betae, Puccinia helianthi)</i> White Rust <i>(Albugo tragopogonis)</i>	6.0-20.0 fl. oz. (0.10-0.33 lb. ai/A)	For powdery mildew, make preventative applications on a 5- to 7-day schedule. For all other diseases, begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
Cercospora Leaf Spot <i>(Cercospora betae, C. pastinaceae)</i> Powdery Mildew <i>(Erysiphe polygoni, Leveillula taurica)</i>	9.0-15.5 fl. oz. (0.15-0.25 lb. ai/A)	
<b>Soilborne Diseases</b> Circular Spot, Southern Blight <i>(Sclerotium rolfsii)</i> Pythium Root Rot <i>(Pythium aphanidermatum)</i> Rhizoctonia Stem Canker, Crown Rot <i>(Rhizoctonia solani)</i>	0.40-0.80 fl oz/1000 row feet (0.0065-0.013 lb. ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 5-day interval.</li> <li>● <b>DO NOT</b> apply more than 120 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 6 applications per year at the high rate (20.0 fl oz/A) or 20 applications per year at the low rate (6.0 fl oz/A). When applying at 9.0 fl oz/A, <b>DO NOT</b> apply more than 13 applications per year. When applying at 15.5 fl oz/A, <b>DO NOT</b> apply more than 7 applications per year.</li> <li>● Apply as an in-furrow spray in a minimum of 10 gallons per acre.</li> <li>● AG37589 CS SC Fungicide may be applied the day of harvest (0-day PHI).</li> </ul>		

<sup>1</sup> = Leaves of Root and Tuber Vegetables, Crop Group 2

<sup>2</sup> = Root Vegetable, Crop Subgroup 1B

**Vegetables, Tuberos and Corm Subgroup 1C** Arracacha Arrowroot Artichoke, Chinese and Jerusalem, Canna, Edible Cassava, Edible, Bitter and Sweet, Chayote (root) Chufa, Dasheen (Taro) Ginger, Leren Potato, Sweet Potato Tanier Turmeric Yam, Bean Yam, True

<b>DISEASE CONTROLLED</b>	<b>RATE AG37589 CS SC Fungicide/ACRE</b>	<b>APPLICATION INSTRUCTIONS</b>
<b>Foliar Diseases</b> Alternaria Leaf Spot <i>(Alternaria spp., A. Alternata)</i> Ascochyta Leaf Spot <i>(Ascochyta cynarae)</i> Rust <i>(Uromyces betae, Puccinia helianthi)</i> White Rust <i>(Albugo tragopogonis)</i>	6.0-20.0 fl. oz. (0.10-0.33 lb. ai/A)	For powdery mildew, make preventative applications on a 5- to 7-day schedule. For all other diseases, begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season every 7- 14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  <b>DO NOT</b> apply more than one application of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
Cercospora Leaf Spot <i>(Cercospora betae, C. pastinaceae)</i> Powdery Mildew <i>(Erysiphe polygoni, Leveillula taurica)</i>	9.0-15.5 fl. oz. (0.15-0.25 lb. ai/A)	
<b>Soilborne Diseases</b> Circular Spot, Southern Blight <i>(Sclerotium rolfsii)</i> Rhizoctonia Stem Canker, Crown Rot <i>(Rhizoctonia solani)</i> Pythium Root Rot <i>(Pythium aphanidermatum)</i>	0.40-0.80 fl oz/1000 row feet (0.0065-0.013 lb ai/1000 row feet)	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 5-day interval.</li> <li>● <b>DO NOT</b> apply more than 120 fl oz of product/A/year.                             <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 6 applications per year at the high rate (20.0 fl oz/A) or 20 applications per year at the low rate (6.0 fl oz/A). When applying at 9.0 fl oz/A, <b>DO NOT</b> apply more than 13 applications per year. When applying at 15.5 fl oz/A, <b>DO NOT</b> apply more than 7 applications per year.</li> <li>● <b>DO NOT</b> apply within 14 days of harvest (14-day PHI).</li> </ul>		

## WATERCRESS

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Cercospora Leaf Spot ( <i>Cercospora</i> spp.)	6.0-15.5 fl. oz. (0.10- 0.25 lb. ai/A)	<p>Begin AG37589 CS SC Fungicide applications prior to disease development and continue throughout the season on a 7- to 10-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p><b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 93 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 6 applications per year at the high rate (15.5 fl oz/A) or 15 applications per year at the low rate (6.0 fl oz/A).</li> <li>● <b>DO NOT</b> apply within 7 days of harvest (7-day PHI).</li> </ul>		

# CEREALS

## WHEAT TRITICALE

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
Leaf Rust ( <i>Puccinia triticina</i> = <i>Puccinia recondita</i> f.sp. <i>tritici</i> ) Septoria Leaf and Glume Blotch ( <i>Septoria tritici</i> , <i>Septoria nodorum</i> ) Stem Rust ( <i>Puccinia graminis</i> ) Stripe Rust ( <i>Puccinia striiformis</i> ) Tan Spot ( <i>Pyrenophora tritici-repentis</i> )	4.0-12.0 fl. oz. (0.07-0.20 lb. ai/A)	Apply AG37589 CS SC Fungicide prior to disease development. Applications may be made by ground, air or chemigation. A crop oil concentrate adjuvant may be added at 1.0% v/v to optimize efficacy.  <b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicide before alternation with a fungicide that is not in Group 11. <b>DO NOT</b> make more than two applications of AG37589 CS SC Fungicide or other Group 11 fungicide per season.
Powdery Mildew ( <i>Erysiphe graminis</i> )	7.5-11.0 fl. oz. (0.125- 0.175 lb. ai/A)	
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> apply after Feekes 10.54.</li> <li>● <b>DO NOT</b> apply at less than 14-day interval.</li> <li>● <b>DO NOT</b> apply more than 24 fl oz of product/A/year.               <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.40 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 2 applications per year at the high rate (12.0 fl oz/A) or 6 applications per year at the low rate (4.0 fl oz/A). When applying at 7.5 fl oz/A, <b>DO NOT</b> apply more than 3 applications per year. When applying at 11.0 fl oz/A, <b>DO NOT</b> apply more than 2 applications per year.</li> <li>● <b>DO NOT</b> apply within 7 days (7-day PHI) for forage and hay.</li> <li>● <b>DO NOT</b> apply within 14 days of grazing (14-day PHI).</li> </ul>		

## WILD RICE

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
<p>Brown Spot (<i>Bipolaris oryzae</i> or <i>Bipolaris sorokiana</i>)</p> <p>Also known as <i>Helminthosporium oryzae</i> and <i>H. sativum</i></p> <p>Stem Rot (<i>Nakataea sigmoidea</i>)</p>	<p>12.5-15.5 fl. oz. (0.20-0.25 lb. ai/A)</p>	<p>Apply AG37589 CS SC Fungicide prior to disease development. Applications may be made by ground, air, or chemigation. For aerial application, use volumes of 5-10 GPA. An adjuvant may be added at specified rates.</p> <p>For foliar diseases, apply AG37589 CS SC Fungicide prior to disease development. Apply during tillering, boot, early heading, or at initial sign of disease. Under heavy disease pressure and conditions favorable for disease development, a second application may be applied.</p> <p><b>DO NOT</b> apply more than two sequential applications of AG37589 CS SC Fungicide or other Group 11 fungicide before alternation with a fungicide that is not in Group 11. Do not make more than two applications of AG37589 CS SC Fungicide or other Group 11 fungicide per season.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> exceed the maximum rate listed in this crop table.</li> <li>● <b>DO NOT</b> treat wild rice fields used for aquaculture of fish and crustaceans.</li> <li>● <b>DO NOT</b> apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Use care in making applications near non-target aquatic habitats.</li> <li>● <b>DO NOT</b> apply at less than 7-day interval.</li> <li>● <b>DO NOT</b> apply more than 37.5 fl oz of product/A/year. <ul style="list-style-type: none"> <li>○ <b>DO NOT</b> apply more than 0.70 lb ai/A/year of azoxystrobin-containing products.</li> </ul> </li> <li>● <b>DO NOT</b> apply more than 2 applications per year at the high rate (15.5 fl oz/A) or 3 applications per year at the low rate (12.5 fl oz/A).</li> <li>● <b>DO NOT</b> allow release of irrigation or flood water for at least 14 days after the last application.</li> <li>● <b>DO NOT</b> apply within 28 days of harvest (28-day PHI).</li> </ul>		

**AG37589 CS SC Fungicide Rate Conversion Chart**

<b>F1 Ounces Product/A</b>	<b>Lb ai/A</b>	<b>Treated Acres/Gal Product</b>
4.0	0.07	32.0
5.0	0.08	25.6
5.5	0.09	23.2
6.0	0.10	21.3
6.2	0.10	21.3
7.0	0.11	18.3
8.5	0.14	15.4
9.0	0.15	14.2
9.2	0.15	14.2
10.0	0.16	13.0
11.0	0.18	11.6
12.0	0.20	10.4
12.3	0.20	10.4
13.0	0.21	9.8
14.0	0.23	9.1
15.4	0.25	8.3
15.5	0.25	8.3
18.3	0.30	6.9
18.5	0.30	6.9
20.0	0.33	6.4
20.3	0.33	6.4
24.5	0.40	5.2

## POST HARVEST APPLICATIONS

### BANANAS

### PLANTAINS

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS	
Crown Rot/Crown Mold <i>(Colletotrichum musae,</i> <i>Fusarium pallidoroseum,</i> <i>Acremonium spp.,</i> <i>Ceratocystis paradoxa,</i> <i>Glomerella cingulata,</i> <i>Penicillium spp.)</i>	200-400 ppm solution	Apply AG37589 CS SC Fungicide as a single application of a 200-400 ppm solution to achieve good coverage. The application may be made as a spray, dip or may be painted onto the cut ends of the bananas. Application of the 200 ppm rate is appropriate for short distance transportation (e.g., within the USA). When a longer time in transport is expected (export), use the 300-400 ppm rate. If alum (1% w/v) is added to the spray solution, stir the suspension frequently as sedimentation and flocculation may occur. Addition of a non-ionic surfactant (0.10% v/v) may improve the compatibility of this mixture.  <b>Amount of AG37589 CS SC Fungicide to Mix 100 Gallons for Post-Harvest Banana Applications</b>	
		<b>AG37589 CS SC Fungicide Use Rate</b>	<b>100.0 gal Spray Solution</b>
		200 ppm	11 fl oz
		300 ppm	15 fl oz
		400 ppm	21 fl oz
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>• <b>DO NOT</b> make more than one application to bananas as post-harvest treatment.</li> <li>• AG37589 CS SC Fungicide may be degraded by exposure to direct sunlight. <b>DO NOT</b> store treated fruit in direct sunlight.</li> </ul>			

CITRUS FRUIT CROP GROUP 10-10 Calamondin

Citron  
 Citrus Hybrids Grapefruit Kumquat Lemon  
 Lime Mandarin  
 Orange (sour and sweet)  
 Pummelo Satsuma Mandarin Tangerine  
 Uniq Fruit Hybrid

**Complete List of Citrus Fruit Crops:** Australian Desert Lime (*Eremocitrus glauca*); Australian Finger Lime (*Microcitrus australasica*); Australian Round Lime (*Microcitrus australis*); Brown River Finger Lime (*Microcitrus papuana*); Calamondin (*Citrofortunella microcarpa*); Citron (*Citrus medica*); Citrus Hybrids, *Citrus* spp., *Eremocitrus* spp., *Fortunella* spp., *Microcitrus* spp., and *Poncirus* spp.; Grapefruit (*Citrus paradise*); Japanese Summer Grapefruit (*Citrus natsudaidai*); Kumquat (*Fortunella* spp.); Lemon (*Citrus limon*); Lime (*Citrus aurantiifolia*); Mediterranean Mandarin (*Citrus deliciosa*); Mount White Lime (*Microcitrus garrowayae*); New Guinea Wild Lime (*Microcitrus warburgiana*); Orange, Sour (*Citrus aurantium*); Orange, Sweet (*Citrus sinensis*); Pummelo (*Citrus maxima*); Russell River Lime (*Microcitrus inodora*); Satsuma Mandarin (*Citrus unshiu*); Sweet Lime (*Citrus limetta*); Tachibana Orange (*Citrus tachibana*); Tahiti Lime (*Citrus latifolia*); Tangelo (*Citrus x tangelo*); Tangerine (Mandarin) (*Citrus reticulata*); Tangor (*Citrus nobilis*); Trifoliolate Orange (*Poncirus trifoliata*); Uniq Fruit (*Citrus aurantium* Tangelo group); cultivars, varieties and/or hybrids of these.

DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/A CRE	APPLICATION INSTRUCTIONS
Penicillium Decays Green Mold, Whisker Mold, Suppression of Blue Mold ( <i>Penicillium</i> spp.) Diplodia Stem- End Rot ( <i>Diplodia natalensis</i> ) Phomopsis Stem-End Rot ( <i>Phomopsis citrii</i> )	See Application Instructions	Post-harvest treatment of citrus must be conducted with a closed automated system only, and not in an automated system that is not closed. Post-harvest treatment of citrus must not be made using a mechanically-pressurized handgun.
		<p><b>For high volume (dilute) applications:</b> Mix 32-64 fl oz of AG37589 CS SC Fungicide in 25-100 gallons of an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated. Use T-Jet, flooders, or similar application systems.</p> <p><b>For low volume (concentrate) applications:</b> Mix 32-64 fl oz of AG37589 CS SC Fungicide in 7- 25 gallons of water, wax/oil emulsion, or aqueous dilution of wax/oil emulsion for the crop being treated. Apply to 250,000 lb of fruit. Use a controlled-droplet type of applicator or similar system.</p>
		<p><b>For dip applications:</b> Mix 32-64 fl oz of AG37589 CS SC Fungicide in 100 gallons of water, wax/oil emulsion, or aqueous dilution of wax/oil emulsion. Dip for approximately 30 seconds and allow fruit to drain. For maximum decay control, treat citrus fruit once before storage and once after storage, just prior to marketing.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> make more than two applications to citrus fruit as post-harvest treatments.</li> <li>● AG37589 CS SC Fungicide may be degraded by exposure to direct sunlight. <b>DO NOT</b> store treated fruit in direct sunlight.</li> </ul>		

### Tuberous and Corm Vegetable Subgroup 1C - Post harvest

Arracacha; Arrowroot; Artichoke, Chinese; Artichoke, Jerusalem; Canna, Edible; Cassava, Bitter and Sweet; Chayote (root); Chufa; Dasheen; Ginger; Leren; Potato; Sweet Potato; Tanier; Turmeric; Yam Bean; Yam, True.

Use AG37589 CS SC Fungicide as a post-harvest spray for the control of certain post-harvest rots caused by Silver Scurf (*Helminthosporium solani*), *Fusarium* species, Late Blight (*Phytophthora infestans*), and Pink Rot (*Phytophthora erythroseptica*).

Application Method	DISEASE CONTROLLED	RATE AG37589 CS SC Fungicide/ACRE	APPLICATION INSTRUCTIONS
In-Line Aqueous Spray Application	Silver Scurf Fusarium Dry Rot Late Blight Pink Rot	0.6 fl oz/ton of tubers	<ul style="list-style-type: none"> <li>● Ensure proper coverage of the tubers. Ensure tubers are tumbling as they are treated.</li> <li>● Mix the fungicide solution in an appropriate amount of water for the crop being treated.</li> <li>● Use T-jet, CDA, or similar application system.</li> </ul>
<b>DO NOT make more than one post-harvest application to the tubers.</b>			
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>● <b>DO NOT</b> use on seed potatoes or seed pieces.</li> <li>● Ensure the AG37589 CS SC Fungicide solution remains in suspension by using agitation.</li> </ul>			

---

**STORAGE AND DISPOSAL**

---

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage**

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

**Pesticide Disposal**

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

**Container Handling [less than or equal to 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

**Container Handling [greater than 5 gallons]**

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn

the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

#### WARRANTY DISCLAIMER AND NOTICE

---

#### IMPORTANT: READ BEFORE USE

---

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ACETO LIFE SCIENCES, L.L.C., D/B/A ACTYLIS MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aceto Life Sciences, L.L.C., d/b/a Actylis is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Aceto Life Sciences, L.L.C., d/b/a Actylis DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT Aceto Life Sciences, L.L.C., D/B/A ACTYLIS'S ELECTION, THE REPLACEMENT OF PRODUCT

[Option marketing and non-FIFRA label statements]

[All trademarks are the property of their respective owners.]

[Active ingredient made [in] [Country] , [formulated] [in Country] [and] [package] [in Country].]

[Shake Well Before Using]

[\*Not for Use in [State]]

[Peel Down for Directions]