DESCRIPTION OF INDIANHEAD FLY KILLER

When used as directed, it is especially suitable for use in dairies, harse barns, restaurants, flaur mills, family dwellings, food markets, meat packing plants,

INDIANHEAD FLY KILLER is a petroleum base insecticide and depends upon its high content of pyrethrin for its exceptional insect-killing power. All open food containers and exposed foods should be covered during spraying. Use as a space spray in food processing plants only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using.

INDIANHEAD Fly Killer is formulated and produced for satisfactory use in all steam, compressed air, and manual sprayers, capable of producing a fine floating mist (4 mecrons).

KEEP FLUID AT ROOM TEMPERATURE CAUTION: HARMF JL IF SWALLOWED. REMOVE OR COVER FISH BOWLS AND AQUARIUMS WHEN SPRAYING. NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. ILITIES AND EQUIPMENT MUST BE THOROUGHLY WASHED WITH AN AC-CEPTABLE DETERGENT SOLUTION AND THEN RINSED WITH POTABLE WATER TO REMOVE ALL TRACES OF CONTAMINATION. VACATE TREATED AREA AFTER TREATVENT. VENTILATE BEFORE RECOULPYING AREλ.



2777-21

INDIANHEAD

ACTIVE INCREDIENTS	8 100%
*N-octyl bicycloheptene dicarbo Pyrethrins	
**Technical piperenyl butoxide Petroleum distillate	
TOTAL ACTIVE INGREDIEN	
*MKG 284, Insecticide Swersies % 8. **Consists of \$24% (busylcarbity)	Parent No. 2,476,612. M-propylpipprenyi) eti
and 0.06% relates compounts.	COMPANY
P. O. BOX 6:	38
SCOTTSDALE, ARIZO	

ACHTICAN: KEEP OUT OF REACH OF CHILDREN SEE LEFT PANEL FOR ADDITIONAL CAUTIONS

DIRECTIONS

To control FLIES, MOSQUITOES, FLYI POSED MOTHS, fag infested area as fol APPLICATION: Close all outside vents and off all coolers, fans and ventilators. Opi cupboards, storage bins and stock rooms. toward walls, windows and cailing until room fagged from cailing to floor. A minimum ounces of INDIANHEAD Fly Killer per 100 space is usually sufficient to fag encloses above listed FLYING INSECTS.

OUTSIDE APPLICATION: AS AN AID I ING FLIES AND MOSQUITOES: Open all cans. Open all doors and windows of sheds buildings in area. Move dispensing machi windward side of area so that fag may slow! through entire space to be treated. If pr days or hours when drift is from high to lo very low velocity. Systematic and freque old in the control of flies and masquitoes,

To control ROACHES, ANTS, CRICKETS SILVER FISH, FLEAS, TICKS AND GRA exposed stages, fag infested areas as fallow PLICATION: Close all outside vents and off all coolers, fans and ventilators. Or cupboards, storage bins and stock rooms. S rics, boxes and closely piled merchandise. be obtained if the operator will place the chine approximately 6 to 8 feet from th sprayed. Direct the fcg toward walls and a ing attention to cracks, crevices, pipe (boards and other openings where insects Direct the spray so as to contact as many sible. Three to 4 ownces per 1000 cubic fo directions above, is usually sufficient.

BEST AVAILABLE COPY

ACCEPTED

JUL 1 7 1985

Under the Factoral Insertica to Fungleide, and Reden with viel. as amended, for the pesic, de Cobnu beretzipes KPA Reg. No. 3

