



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

June 5, 2025

Katie Woodall
Agent for Decco US Post-Harvest
Decco US Post-Harvest
1713 S. California Ave.
Monrovia, CA 91016-0120

Subject: PRIA Label Amendment – Add uses that are already approved.
Product Name: Decco Fludi SC
EPA Registration Number: 2792-84
Application Date: 08/01/2023
Case Number: 487451

Dear Katie Woodall:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or

claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.


If you have any questions, please contact Senedu Alemu at alemu.senedu@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Manjula Unnikrishnan", written in a cursive style.

Manjula Unnikrishnan, Product Manager
21 Fungicide Branch
Registration Division (7505T)
Office of Pesticide Programs

Enclosure



FLUDIOXONIL	GROUP	12	FUNGICIDE
-------------	-------	----	-----------

Decco Fludi SC

[ABN: DECCO Sustain]

For Control of listed Post-Harvest Diseases.

Active Ingredient:	
Fludioxonil*	20.9%
Other Ingredients	79.1%
Total	100.0%

*CAS No. 131341-86-1

Decco Fludi SC is a flowable suspension concentrate.

Decco Fludi SC contains 1.92 lbs ai per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements and direction for use inside the booklet.

EPA Reg. No. 2792-84

EPA Est. No.: _____

Net Contents: _____

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to by a poison control center or doctor. • DO NOT give anything to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>For chemical emergency assistance (spill, leak, fire, or accident), call CHEMTREC: (800) 424-9300 Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Poison Control Hotline at 1-800-222-1222 for emergency medical treatment information 24 hours per day, 7 days per week.</p>	

PRECAUTIONARY STATEMENTS

Hazards to Human and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contact with skin or clothing. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, and Viton® \geq 14 mils.

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. For terrestrial uses: **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsates.

Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Physical or Chemical Hazards

DO NOT store or use near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

PRODUCT INFORMATION

Resistance Management

Decco Fludi SC is a protective fungicide used to aid in the control of several post-harvest diseases in post-harvest treatment facilities. Decco Fludi SC contains fludioxonil that is in the phenylpyrrole class of chemistry and has a unique mode of action, which leads to increased glycerol synthesis. Fungal isolates with acquired resistance to Group 12 may eventually dominate the fungal population if Group 12 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by fludioxonil or other Group 12 fungicides. A disease management program that includes alternation or tank mixes between Decco Fludi SC and other labeled fungicides that have a different mode of action may prevent pathogen populations from developing resistance. Use sanitation and other cultural practices to minimize disease in order to control disease and prevent or delay disease development. Application Instruction: To avoid product degradation, **DO NOT** store treated fruit in direct sunlight. To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Decco Fludi SC or other Group 12 fungicides within a growing season sequence with different groups that control the same pathogens.

- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM guidances for specific crops and pathogens.
- For information or to report suspected resistance contact Decco US Post-Harvest Inc. at (626) 358 1838. You can also contact university extension specialist to report resistance.

MIXING PROCEDURES

Vigorously shake the product container before mixing. Vigorous agitation is necessary for proper dispersal of the product. Ensure the mix solution remains in suspension by using agitation.

If tank-mixing, add the specified amount of other products specified for tank mixture after Decco Fludi SC has completely and uniformly dispersed into the mix carrier. Add tank mix partners in this order unless label directions or other considerations indicate otherwise: wettable powders, wettable granules (dry flowables), liquid flowables, liquids, and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Continue agitation to maintain a uniform suspension until all of the spray solution has been applied. Maintain agitation until all of the mixture has been applied.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CROP USE DIRECTIONS

Citrus Fruit Group 10-10

Australian desert lime (*Eremocitrus glauca*); Australian finger lime (*Microcitrus australasica*); Australian round lime (*Microcitrus australis*); Brown River finger lime (*Microcitrus papuana*); Calamondin (*Citrofortunella microcarpa*); Citron (*Citrus medica*); Citrus hybrids, *Citrus* spp., *Eremocitrus* spp., *Fortunella* spp., *Microcitrus* spp., and *Poncirus* spp.; Grapefruit (*Citrus paradisi*); Japanese summer grapefruit (*Citrus natsudaikai*); Kumquat (*Fortunella* spp.); Lemon (*Citrus limon*); Lime (*Citrus aurantiifolia*); Mediterranean mandarin (*Citrus deliciosa*); Mount White lime (*Microcitrus garrowayae*); New Guinea wild lime (*Microcitrus warburgiana*); Orange, sour (*Citrus aurantium*); Orange, sweet (*Citrus sinensis*); Pummelo (*Citrus maxima*); Russell River lime (*Microcitrus inodora*); Satsuma mandarin (*Citrus unshiu*); Sweet lime (*Citrus*

limetta); Tachibana orange (*Citrus tachibana*); Tahiti lime (*Citrus latifolia*); Tangelo (*Citrus x tangelo*); Tangerine (Mandarin) (*Citrus reticulata*); Tangor (*Citrus nobilis*); Trifoliate orange (*Poncirus trifoliata*); Uniq fruit (*Citrus aurantium* Tangelo group); cultivars, varieties and/or hybrids of these.

Use Decco Fludi SC as a post-harvest dip, drench, flood, or spray for control of post-harvest disease caused by:

Green or blue mold (*Penicillium* spp.); Diplodia stem-end rot (*Lasiodiplodia theobromae*); Phomopsis stem-end rot (*Diaporthe citri*); and Gray mold (*Botrytis cinerea*).

Application Method	Disease	Rate	Application Directions
In-Line Dip / Drench	Green mold Blue mold Diplodia stem-end rot Gray mold	33-66 fl oz / 100 gallons (0.5 to 1.0 lb a.i. fludioxonil/100 gallons	Mix 33-66 fl oz of Decco Fludi SC (0.5 to 1.0 lb a.i. fludioxonil) in 100 gal of an appropriate water, wax / oil emulsion, or aqueous dilution of wax / oil emulsion. Dip for a minimum of 30 seconds and allow fruit to drain.
In-Line Aqueous or Fruit Coating Spray Application	Green mold Blue mold Diplodia stem-end rot Gray mold	33-66 fl oz / 250,000 lbs of fruit (0.5 to 1.0 lb a.i. fludioxonil / 250,000 lbs of fruit)	Ensure proper coverage of the fruit. Use T-jet, CDA, or similar application system. Mix the fungicide solution in an appropriate water, wax/oil emulsion, or aqueous dilution of wax/oil emulsion for crop being treated.
Restrictions: DO NOT apply more than 66 fl oz of product (1.0 lb a.i. fludioxonil) per application. DO NOT apply more than two applications (2.0 lb a.i. fludioxonil) per year. Retreatment Interval: For maximum decay control, treat fruit once before storage and once after storage, just prior to marketing.			
Ensure the Decco Fludi SC solution remains in suspension by using agitation. Decco Fludi SC is stable at temperatures of 60 °C (or 140 °F) that can be used to disinfect high-volume, recycling tanks.			

Pome Fruit

Apple (*Malus domestica*); Azarole (*Crataegus azarolus*); Crabapple (*Malus* spp.); Loquat (*Eriobotrya japonica*); Mayhaw (*Crataegus aestivalis*, *C. opaca*, and *C. rufula*); Medlar (*Mespilus germanica*); Pear (*Pyrus communis*); Pear, Asian (*Pyrus* spp.); Quince (*Cydonia oblonga*); Quince, Chinese (*Chaenomeles speciosa*); Quince, Japanese (*Chaenomeles japonica*); Tejocote (*Crataegus mexicana*) and cultivars, varieties and / or hybrids of these.

Use Decco Fludi SC as a post-harvest dip, drench, flood, or spray for control post-harvest diseases caused by:

Blue mold (*Penicillium expansum*); Gray mold (*Botrytis cinerea*); Bull's-eye rot (*Neofabraea* spp.); Rhizopus rot (*Rhizopus stolonifera*); Bitter rot (*Colletotrichum gloeosporioides*); Sphaeropsis rot (*Sphaeropsis pyriputrescens*); Phacidiopycnis rot (*Phacidiopycnis piri*); Speck rot (*Phacidiopycnis washingtonensis*); White rot (*Botryosphaeria dothidea*); Alternaria rot (side rot); and Surface mold (*Alternaria alternata*).

Application Method	Disease	Rate	Application Directions
Bin / Truck Drench or In-Line Dip / Drench or Flooder	Blue mold Gray mold Bitter rot Speck rot White rot Phacidiopycnis rot Sphaeropsis rot Alternaria rot and surface mold	10-16 fl oz / 100 gallons (0.15 to 0.24 lb a.i. fludioxonil / 100 gallons)	Ensure proper coverage of the fruit. For re-cycling in-line drench or dip treatments, the fungicide solution may be prepared in water. For in-line drench or dip application, treat fruit for 15-30 seconds and allow fruit to drain. Fruit coatings may be applied separately after aqueous fungicide treatment.
	Rhizopus rot Bull's-eye rot	16 fl oz / 100 gallons (0.24 lb a.i. fludioxonil / 100 gallons)	
Restrictions for Bin / Truck Drench or In-Line Dip / Drench or Flooder Applications: DO NOT apply more than 16 fl oz of product (0.24 lb a.i. fludioxonil) per application. DO NOT apply more than two applications (0.48 lb a.i. fludioxonil) per year. Retreatment Interval: For maximum decay control, treat fruit once before storage and once after storage, just prior to marketing.			
In-Line Aqueous or Fruit Coating Spray Application	Blue mold Gray mold Rhizopus rot Bull's-eye rot Bitter rot Sphaeropsis rot Phacidiopycnis rot White rot	16-32 fl oz / 200,000 lbs of fruit (0.24 to 0.48 lb a.i. fludioxonil/ 200,000 lbs of fruit)	Ensure proper coverage of the fruit. Mix the fungicide solution in an appropriate water, wax/oil emulsion, or aqueous dilution of a

	Alternaria rot and surface mold		wax / oil emulsion for crop being treated. Use T-jet, CDA, or similar application system.
Restrictions for In-Line Aqueous or Fruit Coating Spray Applications: DO NOT apply more than 32 fl oz of product (0.48 lb a.i. fludioxonil) per application. DO NOT apply more than two applications (0.96 lb a.i. fludioxonil) per year. Retreatment Interval: For maximum decay control, treat fruit once before storage and once after storage, just prior to marketing.			
Ensure the Decco Fludi SC solution remains in suspension by using agitation. Decco Fludi SC is stable at temperatures of 60 °F (or 140 °F) that can be used to disinfect high volume, recycling tanks.			

Pomegranates

Use Decco Fludi SC as a post-harvest dip / drench for control of Botrytis fruit rot in pomegranates.

Application Method	Disease	Rate	Application Directions
In-Line Dip / Drench	Botrytis fruit rot	32 fl oz / 100 gallons of water (0.48 lb a.i. fludioxonil/ 100 gallons of water)	Mix 32 fl oz of Decco Fludi SC (0.48 lb a.i. fludioxonil) in 100 gal of an appropriate water, wax / oil emulsion, or aqueous dilution of wax / oil emulsion. Dip for approximately 30 seconds and allow fruit to drain.
Restrictions: DO NOT apply more than 32 fl oz of product (0.48 lb a.i. fludioxonil) per application. DO NOT apply more than one application (0.48 lb a.i. fludioxonil) per year.			
Ensure the Decco Fludi SC solution remains in suspension by using agitation. Decco Fludi SC is stable at temperature of 60 °C (or 140 °F) than can be used to disinfect high-volume, recycling tanks.			

Cherries: Capulin; Cherry, black; Cherry, Nanking; Cherry, sweet (*Prunus avium*); Cherry, tart (*Prunus cerasus*); as well as other cultivars and hybrids of these.

Use Decco Fludi SC as a post-harvest treatment for control of post-harvest diseases caused by: Brown rot (*Monilinia* spp.); Gray mold (*Botrytis cinerea*); Rhizopus rot (*Rhizopus stolonifer*); Gilbertella rot (*Gilbertella persicaria*)

Application Method	Disease	Rate	Application Directions
In-line Aqueous or Flooder Application High-Volume (Diluted-Spray) Application	Brown rot Gray mold Rhizopus rot Gilbertella rot	16-32 fl oz per 50,000 lbs of fruit (0.24 to 0.48 lb a.i. fludioxonil per 50,000 lbs of fruit)	Mix 16 fl oz of Decco Fludi SC (0.24 lb a.i. fludioxonil) in 50-100 gallons or 32 fl oz of Decco Fludi SC (0.48 lb a.i. fludioxonil) in 100 gallons of water, wax/emulsion, or aqueous dilution of a wax/oil emulsion. Use flooders, T-jet, or similar application system.
Restrictions: DO NOT apply more than 32 fl oz of product (0.48 lb a.i. fludioxonil) per application. DO NOT apply more than one application (0.48 lb a.i. fludioxonil) per year.			
Ensure the Decco Fludi SC solution remains in suspension by using agitation.			

Stone Fruit: Apricot (*Prunus armeniaca*); Apricot, Japanese; Jujube, Chinese; Nectarine (*Prunus persica*); Peach (*Prunus persica*); Plum (*Prunus domestica*, *Prunus* spp.); Plum, American; Plum, Beach; Plum, Canada; Plum, cherry; Plum Chickasaw (*Prunus angustifolia*); Plum, Damson (*Prunus domestica* spp. *Insititia*); Plum, Japanese (*Prunus salicina*); Plum, Klamath; Plum, prune; Plumcot (*Prunus armeniaca* x *P. domestica*); Prune (fresh) (*Prunus domestica*, *Prunus* spp.); Sole; as well as other cultivars and hybrids of these.

Use Decco Fludi SC as a post-harvest dip/drench or spray for control of post-harvest diseases caused by:

Brown rot (*Monilinia* spp.); Gray mold (*Botrytis cinerea*); Rhizopus rot (*Rhizopus stolonifer*); Gilbertella rot (*Gilbertella persicaria*)

Application Method	Disease	Rate	Application Directions
In-Line Dip / Drench	Brown rot Gray mold Rhizopus rot Gilbertella rot	16 fl oz / 100 gallons (0.24 lb a.i. fludioxonil/ 100 gallons)	Mix 16 fl oz of Decco Fludi SC (0.24 lb a.i. fludioxonil) in 100 gallons of water, wax/emulsion, or aqueous dilution of wax/oil emulsion. Dip for approximately 30 seconds and allow fruit to drain.
Restrictions for In-Line Dip / Drench Applications: DO NOT apply more than 16 fl oz of product (0.24 lb a.i. fludioxonil) per application.			

DO NOT apply more than one application (0.24 lb a.i. fludioxonil) per year.			
In-Line Aqueous or Fruit Coating Spray Application	Brown rot Gray mold Rhizopus rot Gilbertella rot	16-32 fl oz per 200,000 lbs of fruit (0.24 to 0.48 lb a.i. fludioxonil per 200,000 lbs of fruit)	Ensure proper coverage of the fruit. Mix 16-32 fl oz of Decco Fludi SC (0.24 to 0.48 lb a.i. fludioxonil) in an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for crop being treated. Use T-Jet, CDA, or similar application system. For maximum efficacy, use low volume concentrated application system for treatment of plums.
Restrictions for In-Line Aqueous or Fruit Coating Spray Applications: DO NOT apply more than 32 fl oz of product (0.48 lb a.i. fludioxonil) per application. DO NOT apply more than one application (0.48 lb a.i. fludioxonil) per year.			
Ensure the Decco Fludi SC solution remains in suspension by using agitation. Decco Fludi SC is stable at temperature of 60 °C (or 140 °F) than can be used to disinfect high-volume, recycling tanks.			

Sweet Potato

Use Decco Fludi SC as a post-harvest dip/drench or low volume application for control of post-harvest rots caused by *Rhizopus stolonifer*.

Application Method	Disease	Rate	Application Directions
In-Line Dip / Drench	Rhizopus rot	16-32 fl oz / 100 gallons (0.24 to 0.48 lb a.i. fludioxonil/ 100 gallons)	Mix 16-32 fl oz of Decco Fludi SC (0.24 to 0.48 lb a.i. fludioxonil) in 100 gallons of water, wax/emulsion, or aqueous dilution of wax/oil emulsion. Dip for approximately 30 seconds and allow fruit to drain.
Restrictions for In-Line Dip / Drench Applications: DO NOT apply more than 32 fl oz of product (0.48 lb a.i. fludioxonil) per application.			

DO NOT apply more than one application (0.48 lb a.i. fludioxonil) per year.			
In-Line Aqueous or Fruit Coating Spray Application	Rhizopus rot	16 fl oz / 200,000 lbs of sweet potatoes 16 fl oz (0.24 lb a.i. fludioxonil / 200,000 lbs of sweet potatoes)	Ensure proper coverage of the fruit. Mix 16 fl oz of Decco Fludi SC (0.24 lb a.i. fludioxonil) in an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for crop being treated. Use-T-jet, CDA or similar application system.
Restrictions for In-Line Aqueous or Fruit Coating Spray Applications: DO NOT apply more than 16 fl oz of product (0.24 lb a.i. fludioxonil) per application. DO NOT apply more than one application (0.24 lb a.i. fludioxonil) per year.			
Ensure the Decco Fludi SC solution remains in suspension by using agitation. Decco Fludi SC is stable at temperature of 60 0C (or 140 0F) than can be used to disinfect high-volume, recycling tanks.			

True Yam

Use Decco Fludi SC as a post-harvest dip for control of certain post-harvest rots caused by *Penicillium* and *Fusarium* species.

Application Method	Disease	Rate	Application Directions
Post-Harvest Dip Application	Brown rot Gray mold Rhizopus rot Gilbertella rot	16-32 fl oz per 100 gallons (0.24 to 0.48 lb a.i. fludioxonil per 100 gallons)	Mix 16-32 fl oz of Decco Fludi SC (0.24 to 0.48 lb a.i. fludioxonil) in 100 gallons of an appropriate water, wax/emulsion, or aqueous dilution of wax/oil emulsion. Dip for approximately 30 seconds and allow fruit to drain.
Restrictions: DO NOT apply more than 32 fl oz of product (0.48 lb a.i. fludioxonil) per application. DO NOT apply more than one application (0.48 lb a.i. fludioxonil) per year.			
Ensure the Decco Fludi SC solution remains in suspension by using agitation.			

Tuberous and Corm Vegetable Subgroup 1C[*]

Arracacha; Arrowroot; Artichoke, Chinese; Artichoke, Jerusalem; Canna, Edible; Cassava, Bitter and Sweet; Chayote (root); Chufa; Dasheen; Ginger; Leren; Potato; Sweet Potato; Tanier; Turmeric; Yam Bean; Yam, True.

Use Decco Fludi SC as a post-harvest spray for the control of certain post-harvest rots caused by Black rot[*], *Fusarium* species[*], Late Blight (*Phytophthora infestans*) [*], Pink Rot* (*Phytophthora erythroseptica*)[*], Pythium leak[*], Rhizopus rot[*], and Silver scurf (*Helminthosporium solani*)[*]

Application Method	Disease	Rate	Application Directions
In-Line Aqueous Spray Application	Black Rot[*] Fusarium dry rot[*] Late blight[*] Pink rot[*] Pythium leak[*] Rhizopus rot[*] Silver scurf [*]	0.6 fl oz / ton of tubers (0.009 lb a.i. fludioxonil/ ton of tubers)	Ensure proper coverage of the tubers. Tubers must be tumbling as they are treated. Mix the fungicide solution in an appropriate amount of water for the crop being treated. Use T-jet, CDA, or similar application system.
Restrictions: DO NOT apply more than 0.6 fl oz of product (0.009 lb a.i. fludioxonil) per application. DO NOT apply more than one application (0.009 lb a.i. fludioxonil) per year. DO NOT use on seed potatoes or seed pieces.			
Ensure the Decco Fludi SC solution remains in suspension by using agitation. [*For these uses, use in combination with Decco AZ (EPA Reg. No. 70506-351-2792).]			

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage

DO NOT store near heat or open flame. Store in original container only. Keep container closed when not in use. **DO NOT** store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of waste.

Pesticide Disposal

Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of nearest EPA Regional Office for guidance.

Container Handling

[Rigid Plastic Less Than or Equal to 5 Gallons. Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Rigid Plastic Greater Than 5 Gallons. Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of empty container in a sanitary landfill or by other procedures allowed by State and local authorities.]

[Rigid Plastic Bulk and Mini-Bulk Containers. Refillable container. Refill this container with pesticide only. **DO NOT** use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities.]

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

WARRANTY AND DISCLAIMER

DECCO US POST-HARVEST INC. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DECCO US POSTHARVEST INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL DECCO US POST-HARVEST INC. OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES

RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS; LABOR COST, OR OTHER EXPENSES INCURRED IN REPACKAGING, SORTING OR REPROCESSING.

DECCO US POST-HARVEST INC. and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed

by a duly authorized representative of DECCO US POST-HARVEST INC.

Manufactured for:

DECCO US POST-HARVEST INC.

1713 S. California Ave.

Monrovia, CA 91016-0120