



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

December 4, 2025

Marc Sheahin
Senior Regulatory Specialist, Agent
Decco US Post-Harvest Inc.
1713 S. California Ave.
Monrovia, CA 91016-0120

Subject: PRIA Label Amendment – Addition of stone fruit and tomato uses
Product Name: Decco PROPI EC
EPA Registration Number: 2792-90
Application Date: 06/16/2025
Case Number: 662007

Dear Marc Sheahin:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or

claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Raven Crosby at 202-566-0829 or at crosby.raven@epa.gov.

Enclosure

A handwritten signature in cursive script, reading "Carmer Swinger", is displayed on a light-colored rectangular background.

for Manjula Unnikrishnan
Product Manager 21 Fungicide
Branch Registration Division
Office of Pesticide Programs



[MASTER LABEL]

PROPICONAZOLE	GROUP	3	FUNGICIDE
---------------	-------	----------	-----------

Decco PROPI EC

Alternate Brand Name: Decco Pro Max

For Control of Certain Post-Harvest Diseases of Citrus, Stone Fruit and Tomato

Active Ingredient:

Propiconazole: (CAS No. 60207-90-1)..... 20.37%

Other Ingredients:79.63%

Total: 100.00%

Contains 1.84 lbs of Propiconazole per gallon.

ACCEPTED

12/04/2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 2792-90

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See additional precautionary statements and direction for use [on the next panel][inside the booklet].

EPA Reg. No. 2792-90

EPA Est. No. _____

Net Contents: _____

FIRST AID

If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
---------------------	--

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going to doctor for treatment. For general information on this pesticide product call the National Pesticide Telecommunications Network at 1-800-858-7378, or contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST. You may also visit <http://npic.orst.edu>. For medical emergencies, call the poison control center at 1-800-222-1222.

NOTE TO PHYSICIAN

There is no specific antidote. Induce emesis or lavage stomach. Give a saline laxative and supportive therapy.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Discard clothing and other absorbent materials that have been contaminated with this product's concentrate. Do not reuse them.

User Safety Requirements

Follow the manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

User should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to fish and shrimp. **DO NOT** apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment wash water. Refer to product labeling for use restrictions to protect endangered species.

DIRECTIONS FOR USE

It is violation of Federal Law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

PRODUCT INFORMATION

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUE.

This product is a broad-spectrum fungicide used to aid in control of several post-harvest diseases in citrus post-harvest treatment facilities.

USE INFORMATION

Integrated Pest Management

This product should be integrated into an overall disease and pest management strategy (IPM) whenever the use of fungicide is required. Cultural practices known to reduce disease development should be followed. Consult your local agricultural authorities for additional IPM strategies established for your area. This product may be used in State Agricultural Extension advisory (disease forecasting) programs which recommended application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

PROPICONAZOLE	GROUP	3	FUNGICIDE
----------------------	--------------	----------	------------------

For resistance management, Decco PROPI EC contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to Decco PROPI EC and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide/bactericide resistance, take one or more of the following steps:

- Rotate the use of Decco PROPI EC or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Decco US Post Harvest Inc. at (626) 358-1838. You can also contact your pesticide distributor or university extension specialist to report resistance.

CROP USE DIRECTIONS

Citrus Fruit Crop Group 10-10 (Including all cultivars, varieties, and/or hybrids of these)			
Calamondin; Citron; Citrus Hybrids; Grapefruit; Grapefruit, Japanese Summer; Kumquat; Lemon; Lime; Australian Desert, Lime; Australian Finger, Lime; Australian Round, Lime; Brown River Finger, Lime; Mount White, Lime; New Guinea, Wild, Lime; Russell River, Lime; Sweet, Lime; Tahiti Lime; Mediterranean Mandarin; Satsuma Mandarin; Orange, Sour; Orange, Sweet; Tachibana, Orange; Trifoliate Orange; Pummelo; Tangelo; Tangerine (Mandarin); Tangor; Uniq Fruit			
Application Method	Target Disease	Rate (fl oz)	Application Instructions

In-line Dip/Drench	Green mold (<i>Penicillium</i> spp.) Blue mold (<i>Penicillium</i> spp.)	14.5 – 29.0 fl oz /100 gallons (0.21 – 0.42 lb ai/100 gallons)	<ul style="list-style-type: none"> Mix 14.5 – 29.0 fl oz of this product in 100 gallons of an appropriate water, wax/oil emulsion, or aqueous dilution of wax/oil emulsion. Dip for a minimum of 30 seconds and allow fruit to drain.
In-line Aqueous or Fruit-Coating Spray Application	Sour rot (<i>Geotrichum citri-aurantii</i>)	14.5 – 29.0 fl oz /250,000 lbs of fruit (0.21 – 0.42 lb ai/250,000 lbs of fruit)	<ul style="list-style-type: none"> Ensure proper coverage of the fruit. Mix the fungicide solution in an appropriate water, or wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated. Use T-jet, CDA, or similar application system.
PRECAUTIONS			
<ul style="list-style-type: none"> Ensure the product remains in suspension by using agitation. 			
USE RESTRICTIONS			
<ul style="list-style-type: none"> Maximum Single Application Rate: 29.0 fl oz (0.42 lb ai of propiconazole) Maximum Annual Rate: 58 fl oz (0.84 lb ai of propiconazole) DO NOT make more than two post-harvest applications to the fruit. For maximum decay control, treat fruit once before storage and once after storage, just prior to marketing. Minimum Retreatment Interval (RTI) is 0 days. 			

Stone Fruit Crop Group 12-12 (Including all cultivars, varieties, and/or hybrids of these)			
Apricot; Apricot, Japanese; Capulin; Cherry, Black; Cherry, Nanking; Cherry, Sweet; Cherry, Tart; Jujube, Chinese; Nectarine; Peach; Plum; Plum, American; Plum, Beach; Plum, Canada; Plum, Cherry; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plum, Klamath; Plum, Prune; Plumcot, Sloe			
Application Method	Target Disease	Rate (fl oz)	Application Instructions
In-line Dip/Drench	Blue mold (<i>Penicillium</i> spp.) Brown rot (<i>Monilinia</i> spp.)	3.6 – 7.3 fl oz /100 gallons (0.05 – 0.10 lb ai/100 gallons)	<ul style="list-style-type: none"> Mix 3.6 – 7.3 fl oz of this product in 100 gallons of an appropriate water, wax/oil emulsion, or aqueous dilution of wax/oil emulsion. Dip for a minimum of 30 seconds and allow fruit to drain.

In-line Aqueous or Fruit-Coating Spray Application	Green mold (<i>Penicillium</i> spp.) Sour rot (<i>Geotrichum candidum</i>)	3.6 – 7.3 fl oz /250,000 lb of fruit (0.05 – 0.10 lb ai/250,000 lb of fruit)	<ul style="list-style-type: none"> Ensure proper coverage of the fruit. Mix the fungicide solution in an appropriate water, or wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated. Use T-jet, CDA, or similar application system.
PRECAUTIONS			
<ul style="list-style-type: none"> Ensure the product remains in suspension by using agitation. 			
USE RESTRICTIONS			
<ul style="list-style-type: none"> Maximum Single Application Rate: 7.3 fl oz (0.10 lb ai of propiconazole) Maximum Annual Rate: 7.3 fl oz (0.10 lb ai of propiconazole) DO NOT make more than one post-harvest application to the fruit. Decco PROPI EC is stable at temperatures of 60°C (or 140°F) that can be used to disinfest high-volume, recycling tanks. 			

Tomato (Including all cultivars and/or varieties of these)			
Application Method	Disease	Rate (fl oz)	Application Instructions
In-line Dip/Drench	Black mold (<i>Alternaria alternata</i>) Sour rot (<i>Geotrichum candidum</i>)	7.3 – 14.5 fl oz /100 gallons (0.10 – 0.21 lb ai/100 gallons)	<ul style="list-style-type: none">• Mix 7.3 – 14.5 fl oz of this product in 100 gallons of an appropriate water, wax/oil emulsion, or aqueous dilution of wax/oil emulsion.• Dip for a minimum of 30 seconds and allow fruit to drain.• Must be used in tank mixture with fludioxonil. May be used with a food grade adjuvant.
High-Volume (Dilute-Spray) Application		7.3 – 14.5 fl oz /50,000 lb of fruit (0.10 – 0.21 lb ai/50,000 lb of fruit)	<ul style="list-style-type: none">• Ensure proper coverage of the fruit.• Mix the fungicide solution in an appropriate water, or wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated.• Must be used in tank mixture with fludioxonil. May be used with a food grade adjuvant.• Use T-jet, CDA, or similar application system.
PRECAUTIONS			
<ul style="list-style-type: none">• Ensure the product remains in suspension by using agitation.			
USE RESTRICTIONS			

- Maximum Single Application Rate: 14.5 fl oz (0.21 lb ai of propiconazole)
- Maximum Annual Rate: 14.5 fl oz (0.21 lb ai of propiconazole)
- **DO NOT** make more than one post-harvest application to the fruit.
- Decco PROPI EC is stable at temperatures of 60°C (or 140°F) that can be used to disinfest high-volume, recycling tanks.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

DO NOT store near heat or open flame. Store in original containers only. Keep container closed when not in use.

DO NOT store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal waste.

Pesticide Disposal

Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling

Non-refillable plastic container. DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follow: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

WARRANTY AND DISCLAIMER

DECCO US POST-HARVEST INC. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. DECCO US POST-HARVEST INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT ALLOWED BY APPLICABLE LAW, IN NO CASE SHALL DECCO US POST-HARVEST INC. OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS; LABOR COST, OR OTHER EXPENSES INCURRED IN REPACKAGING, SORTING OR REPROCESSING.

DECCO US POST-HARVEST INC. and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement

in writing signed by a duly authorized representative of DECCO US POST-HARVEST INC.

Manufactured for:

DECCO US POST-HARVEST INC.

1713 S. California Ave.

Monrovia, CA 91016-0120