

<b>ACTIVE INGREDIENTS</b>	17.9%	
Diphenylamine		35.0%
Turkey-red Oil (Sulfonated Castor Oil)		7.9%
Benzene		35.0%
<b>INERT INGREDIENTS:</b>	22.1%	
Lamp Black		20.0%
Water		2.1%
<b>TOTAL</b>	<b>100.0%</b>	

**WARNING: FLAMMABLE! KEEP AWAY FROM FIRE, SPARKS, AND HEATED SURFACES. AVOID PROLONGED BREATHING OF VAPOR. AVOID PROLONGED CONTACT WITH HUMAN SKIN.**

## SCREW WORM SMEAR 62

(USDA Formula 62)

### WARNING:

**KEEP OUT OF THE  
REACH OF CHILDREN**

(See Left and Right Side Panels for  
Additional Warnings.)

USDA Reg. No. 2800-62

NET WEIGHT 1 LB. 14 OZS.



# HUMCO

LABORATORY

TEXARKANA, TEXAS 75501

**USE:**  
FOR TREATING SCREW WORM INFESTATIONS IN HORSES AND MULES.  
**WARNING: DO NOT USE ON CATTLE, GOATS, SHEEP, CATS OR DOGS REGARDLESS OF AGE.**  
**DIRECTIONS FOR USE:** Swab thoroughly with a one inch paint brush in the open wound and surrounded areas, push smear into open holes. Repeat treatment in one or two days and twice weekly until wound heals.  
**ANIMALS BEING TREATED WITH THIS PREPARATION SHOULD RECEIVE DAILY SUPERVISION.**  
**NOTICE: NOT EFFECTIVE IN SOME POPULATIONS OR STRAINS OF SCREW WORMS WHICH ARE RESISTANT TO THIS PRODUCT. KEEP JAR TIGHTLY SEALED. STIR CONTENTS BEFORE USING.**

ACCEPTED

JUN 24 1971

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POLYMER PRODUCTS  
L.A. UNDER NO. 2800-62  
ATTACHED COMMENTS