

FIRST AID:


In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg-whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NET CONTENTS:

8 OZS.

CONTROL

NO



DANGER: Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. See left panel for first aid.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard it.

For recommended use levels, consult technical bulletin. For Reformulating Use Only.

RII5-8B

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg-whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ROCCAL® II-50%

Benzalkonium Chloride U.S.P.

ACTIVE INGREDIENTS:
 Benzalkonium Chloride U.S.P. (C₁₂H₂₅N) 50%
 Ethyl Alcohol 50%

INERT INGREDIENTS:
 Water 49%

For use in industrial, institutional, and professional disinfection.

E.P.A. REG. NO. 283-16-AA



2235 Langdon Farm Rd. / Cincinnati, Ohio 45237

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. See left panel for first aid.

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard it.

For recommended use levels, consult technical bulletin. For Reformulating Use Only.

RII5-8P