

Aug 19-1971

2831-12

GENERAL INFORMATION

A liquid weed killer to be applied diluted at the rate of 10 to 1 with diesel or water as a spray on non-cropland acres where bare ground is desired. The degree of weed control and duration of effect will vary with the amount of chemical applied, soil type, rainfall and other conditions.

DIRECTIONS

Apply any time of year except when the ground is frozen. Best results are obtained after predominant species are well established in late spring or early summer. Before treating, calibrate equipment to insure uniform cover of the area to be treated. Measure out the proper amount and apply.

EQUIPMENT

Power sprayers, compressed air hand sprayers, knapsack sprayers or an ordinary sprinkler can be used to apply. After dilution slight agitation is recommended.

WARNING: Harmful if swallowed or absorbed through skin. Causes skin irritations. Do not breathe vapor or spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. In case of contact, wash skin with soap and water; for eyes, immediately flush with large amounts of water for at least 15 minutes and get medical attention. Remove and wash contaminated clothing before re-use.

U.S.D.A. REG. No. 2831-12.

NAPASCO

WEED-A-WAY

**NON-SELECTIVE
WEED AND BRUSH KILLER CONCENTRATE
FAST ACTING — AFFECTS ALL TYPES OF WEEDS**

ACTIVE INGREDIENTS

Petroleum Distillates, Aromatic Petroleum Hydrocarbon, Pentachlorophenol, and other Chlorophenols and Related Products. 40.09%

INERT INGREDIENTS

Tail Oil Fatty Acid Potassium Soap, Water, Nonylphenoxypolyethoxyethanol and Diacetone Alcohol. 59.91%

This Formulation Contains .36 Lb. Pentachlorophenol per gallon

WARNING: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or absorbed through skin. Causes skin irritation. See side panels for further caution.

GENERAL WEED CONTROL

To control most weeds for extended period of time on non-cropland areas such as parking lots, fence rows, loading ramps, plant sites, apply 3 to 5 gallons per 1,000 sq. feet (120 to 200 gallons per acre.)

To control annual weeds and grasses such as foxtail, ryegrass, wild oats, crabgrass, cheat grass, broomsedge, ragweed, lambsquarter and turkey mullein, apply 1 to 2 gallons per 1,000 sq. feet (40 to 80 gallons per acre.)

For other **HARD TO KILL** perennial weeds and grasses such as Johnson grass, soft grass, bouncing bet, broken fern and horsetail apply 3 to 5 gallons per 1,000 sq. feet (120 to 200 gallons per acre.)

To control perennial weeds and grasses such as smooth brome, Canada bluegrass, purple-top, dandelion, dog fennel, goldenrod, plantain and wild carrot apply 2 to 4 gallons per 1,000 sq. feet (80 to 160 gallons per acre.)

WARNING

Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not apply liquid directly to, or otherwise permit it to come into contact with susceptible plants such as vegetables, flowers, fruit trees, ornamentals, cotton, grapes, or other desirable plants and do not permit spray mists containing it to drift onto them. (Coarse sprays are less likely to drift). Vapors from this product may injure susceptible plants in the immediate vicinity. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Do not contaminate waters used for irrigation or domestic purposes. Do not re-use this container for other materials without thorough cleaning. Do not apply when weather conditions favor drift from areas treated.

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.

**DO NOT USE, POUR, SPILL OR STORE NEAR
HEAT OR OPEN FLAME**



NAPASCO CHEMICAL CO.
THIBODAUX, LOUISIANA

NAPASCO CHEMICAL CO.
NEW ORLEANS, LOUISIANA

NAPASCO CHEMICAL CO.
BATON ROUGE, LOUISIANA

NAPASCO CHEMICAL CO.
LAFAYETTE, LOUISIANA

