

#### DESCRIPTION

NAPASCO BRAND TB-15 STAPHEX-S is an activated germicidal cleaner. Staphex can be used as a disinfectant for cleaning and disinfecting recovery rooms, surgical rooms and all kinds of floors and walls. Staphex can be used in homes, hospitals, hotels and schools.

Also effective against a broad spectrum of gram-positive and gram-negative bacteria. Conforms to the specifications of A.O.A.C use dilution test against *S. Choleraesuis* at 1:128 dilution.

NAPASCO BRAND TB-15 STAPHEX-S cleaning system is formulated with heavy duty builders and synthetic detergents for easy cleaning of heavy soil buildup and stripping of oxidized and polymerized old wax residues. TB-15 STAPHEX-S is conductive, and will not adversely effect the conductivity of hospital operating room floors. This dual purpose product provides more cleaning and disinfecting economy than single purpose disinfectants and cleaners.

**DANGER: Keep Out of Reach of Children.** Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

(Continued in Right Panel)

**NAPASCO INTERNATIONAL**  
THIBODAU, LOUISIANA 70301

# NAPASCO

## TB-15

### Staphex Conductive

ACTIVE INGREDIENTS		
Potassium 4- and 6- Chloro-2- phenylphenate	.....	22.16%
Sodium Matasilicate	.....	10.00%
Diethanolamide of Coconut Oil Fatty Acid	.....	5.09%
Isopropyl Alcohol	.....	4.07%
Essential Oils	.....	2.50%
INERT INGREDIENTS	.....	0.50%
Water, Potassium Hydroxide, Diethylene Glycol Monoethyl Ether, Sodium Tripoly Phosphate, and Ethylene Glycol Monobutyl Ether.	.....	77.84%

Phenol Coefficient - 15 *S. Typhosa* — A.O.A.C. Method

**DANGER: KEEP OUT OF REACH OF CHILDREN**

See First Aid and Other Precautions In Side Panels

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

#### DIRECTIONS

- All synthetic detergent system.
- Broad spectrum phenolic germicide for effective disinfecting action.

1. For damp mopping, cleaning walls, woodwork, and painted surfaces, use half ounce per gallon of water.
2. For cleaning soiled floors and other heavily soiled areas, use 3 to 4 ounces per gallon of water.
3. For stripping old wax buildup, use 6 to 8 ounces per gallon of water. Adjust amount according to intensity of buildup.
4. For effective disinfecting use 1 to 2 ounces per gallon of water, let stand for 10 minutes.

Rinse empty container out with water and discard it.

EPA Reg. No. 2831-21  
EPA EST 2831 — LA-1

ACCEPTED

JAN 14 1975

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 2831-21