

NS. WARNING

onto desirable
res or streams.
not store near

If spilled on
ter and soap
lation of spray
ffs, streams or
f children and
tic animals. ef-
ful if swallowed.
treated forage.
ting and soil.

paty and soap
ngredients and
implied or ex-
ner seller shall
erty. Damage or
e or use of the
techniques. Her-
coats and use
S. 227-11-1

IONS



DAL-E-RAD 70 LIQUID

FOR DALLISGRASS CONTROL
IN BERMUDA TURF LAWNS

2753-22

**A
Selective
Dallis
Grass
Destroyer**

**A
Selective
Dallis
Grass
Destroyer**

ACTIVE INGREDIENT:
Monosodium Acid Methylenesulfonate 70.00
INERT INGREDIENTS 30.00
Total Active Ingredient 70.00
Total Net Weight 100.00
CAUTION: Keep this product out of the reach of children. Do not use near water.

VICHEM AND CHEMICAL COMPANY
1985

DIRECTIONS

For use on bermuda turf lawns. Apply to the lawn in the early morning or late afternoon. Do not apply if the lawn is wet or if rain is expected within 24 hours. Do not apply to areas where children or animals play.

RATE OF APPLICATION

Apply 1.0 to 1.5 lbs. of active ingredient per 1,000 sq. ft. of lawn. Do not apply to areas where children or animals play.

CAUTIONS

Keep this product out of the reach of children. Do not use near water. Do not apply to areas where children or animals play.

VICHEM

WEED-E-RAD LIQUID

FOR CONTROL OF JOHNSON GRASS
AND OTHER WEEDS IN COTTON

ACTIVE INGREDIENT:

Monosodium Acid Methanearsonate 51.19%

INERT INGREDIENTS: 48.81%

TOTAL 100.00%

Total Arsenic, as elemental,
all in water soluble form 23.70%

MSMA per gallon 6.66 lbs.

CAUTION: Keep Out of the Reach of Children
See side panel for precautionary measures.

NET CONTENTS GALLONS

Manufactured By

VINELAND CHEMICAL COMPANY

VINELAND, NEW JERSEY

*A
Selective
Weed
Destroyer
for
Cotton*

"WEED-E-RAD 120" (MSMA) Liquid Herbicide used Post-Emergence for weed control in cotton and non-crop areas.

"WEED-E-RAD 120" will control JOHNSON GRASS, GOOSE GRASS, COCKLEBUR, NUT GRASS and DALLIS GRASS.

FOR BAND TREATMENT ON COTTON GROWN IN 40 INCH ROWS

- (1) Mix at the rate of 2½ pints of "WEED-E-RAD 120" plus 2 pints of a suitable surfactant in 40 gallons of water. Add "WEED-E-RAD 120" to water in spray tank. Thoroughly mix by agitation, add surfactant and complete mixing by recirculating solution prior to application.
- (2) Apply at the rate of one (1) gallon of the mixture for each one (1) inch of band width. (13,068 ft. long).
Example: For 12 inch band, apply 12 gallons of the mixture per acre.
- (3) Apply as a directed basal spray to weeds when cotton is 3 inches high and up to first bloom. Do not apply after first bloom opens. Adjust nozzles on spray equipment to allow thorough wetting of weeds with no contact with cotton foliage. Use fan type nozzle, low volume-sprayer, 25 lbs. or less pressure.

Two applications may be made. Should regrowth occur, retreat at the above rates 1 to 3 weeks after first application



Oiling shoes and insecticide sprayers may be modified to apply "WEED-E-RAD 120". Set spray nozzles approximately 5 inches from ground.

Clean equipment after use by flushing with water. Run rinse water in area away from domestic water sources. Do not reuse container.

NOTE: WEED-E-RAD Powder (DSMA) may be used in place of WEED-E-RAD 120 Liquid (MSMA). Use 12-16 lbs. of DSMA-100% Hexahydrate to equal weed control attained with one (1) gallon of WEED-E-RAD 120.