

RAD-E-CATE 35

Surfactant Included

A Fast-acting, Non-Selective Grass and Weed Killer

ACTIVE INGREDIENTS:

*Sodium Dimethylarsinate	29.00%
Dimethylarsinic Acid	5.00%

INERT INGREDIENTS:	66.00%
--------------------	--------

Total	100.00%
-------	---------

*Sodium Cacodylate Arsenic as elemental, all in water soluble form	16.28%
Dimethylarsinic Acid equivalent per gallon	3.25 lbs.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

CAUTION: Harmful if swallowed. Avoid inhalation of spray mist. (Coarse sprays are less likely to drift.) Avoid contact with skin and eyes. Wash thoroughly with soap and water after using. Keep children and domestic animals off treated areas until water has dried soil. To protect fish and wildlife avoid contamination of surface water. Keep off clothing.

ADDITIONAL: If taken internally, induce vomiting and call physician at once.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

A Product Manufactured By
The Vineland Chemical Co.

ACCEPTED

AUG 14 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT CIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 2853-28 SUBJECT
TO ATTACHED COMMENTS.

VINELAND CHEMICAL COMPANY

VINELAND, NEW JERSEY

GENERAL INFORMATION: Rad-E-Cate 35 Herbicide is a solution of cacodylic acid and sodium cacodylate. It is a non-selective herbicide. Rad-E-Cate 35 will kill grasses and weeds on contact. Destruction of weeds and grasses around plants, shrubs, trees and bushes may be accomplished by directing spray to the undesired vegetation.

DIRECTIONS FOR USE: RAD-E-CATE 35 for lawn renovation. To destroy weedy sections of lawns and turf, apply Rad-E-Cate 35 at any time during the warm growing season. Sun and high temperature accelerates destruction. Foliage should be wet thoroughly to run-off using the spray solution at the following rates:

Rate	Amount of Water	Area Covered
1 quart	15-20 gallons	4,150 sq. ft.
1 gallon	65-80 gallons	16,600 sq. ft.

To prepare spray solution fill spray tank with the desired quantity of water. While agitating, add the required amount of Rad-E-Cate 35. Thoroughly mix prior to application. Rad-E-Cate 35 contains a suitable surfactant. NO OTHER WETTING AGENT NEED BE ADDED TO THE SPRAY SOLUTION. The spray should be carefully calibrated for best results and the spray solution applied at the proper rate in the proper amount of water to the properly measured area.

RAD-E-CATE 35 will top-kill perennials. Repeat application may be necessary. Apply second treatment 5 days after first application.

Rad-E-Cate 35 Herbicide may be used in non-crop areas, along rights-of-way, and in drainage ditches. Do not use in vegetable gardens. Do not reuse container.

Clean equipment after use by flushing with water. Run rinse water in area away from domestic water sources.

USE IN ACCORDANCE WITH DIRECTIONS, WARNING AND CAUTION

Reseeding after weed control should be preceded by heavy watering. Consult agriculture experimental station and extension service weed specialists for most desirable seeds and mixtures. Seeds may be worked into soil if necessary, by scarifying or raking. A light top dressing of screened peat moss will help retain moisture during germination. Delay reseeding a minimum of 5 days after last application.

BEFORE USING ANY HERBICIDE, STOP AND READ THE LABEL.

NET CONTENTS 30 GALLONS

EPA REG. NO. 2853-28AA

Printed in U.S.A.

2853-28

08/14/1972

1/1