

GENERAL INFORMATION

VIDELON is a stable aqueous solution containing a mixture of phenoxy salts equivalent to 48 ozs. of oil per gallon. It is non-flammable and non-corrosive to steel.

VIDELON may be used on blue grass, fescues, Bermudagrass and Kentucky blue grass. VIDELON should not be used on golf course greens.

Weeds controlled

Susceptible

- Dandelions (*Taraxacum* spp.)
- Clover (*Trifolium* spp.)
- Common Chickweed (*Stellaria media*)
- Mouse Ear Chickweed (*Ceratistium vulgatum*)
- Plantain (*Plantago* spp.)
- Pigweed (*Amaranthus* spp.)
- Ragweed (*Ambrosia* spp.)
- Lambsquarters (*Chenopodium album*)

Moderately Susceptible*

- Knotweed (*Polygonum* spp.)
- Black Medick (*Medicago sativa*)
- English Daisy (*Bellis perennis*)
- Dock (*Rumex* spp.)
- Purslane (*Portulaca oleracea*)
- Woolly St. John's Wort (*Hypericum acerosum*)
- Mallow (*Malva* spp.)

*For best results apply VIDELON to young, early stages of growth.

NON-WARRANTY

Seller makes no warranty of any kind, expressed or implied concerning the use of this product. User assumes all risk of use or failing which in accordance with the laws or not and accepts the product on these conditions.

ACCEPTED

SEP 8 1966

235333 Subj
REC'D. 10/10/66

For best results use 1 1/2 fl. ozs. of Vipar in 2 gallons of water per 1,000 sq. ft. for smaller areas, 4 pints in 30-40 gallons of water per acre for larger areas.

DIRECTIONS FOR USE

DIRECTIONS FOR USE

TIME OF APPLICATION: Apply when weeds are growing vigorously. Some discoloration may result if this product is applied in very hot weather (over 90°F). Do not apply in dry, hot conditions.

APPLICATION RATE 4 lbs per acre (1 1/2 fl.oz. per 1,000 sq ft)

MIXING In order to ensure even mixing fill half the spray tank with clean water. Add required amount of VIGORATE, mix thoroughly and finally add more water as required.

APPLICATION. When applying **VIMCAF**, always use equipment which applies the product uniformly. A boom type sprayer is recommended. To allow for best distribution of the chemical by the weeds, spray as follows: 6-8-2 days before or after spraying.

METHOD: To get best results spray after cutting or mowing turf for 2-3 days before or after spraying. Avoid cutting or mowing turf for 2-3 days before or after spraying.
Avoid watering for 24 hours after spraying.
FULL EFFECTS OF "VIGORAP" ARE NOT VISIBLE FOR ABOUT TWO-THREE WEEKS


PRECAUTIONS

PRECAUTIONS
Store containers, tightly closed, away from greenhouse, sends plants, fertilizers and water supplies. Protect contents from freezing. "TOPAR" is

Do not spray flower beds, borders or shrubbery. VISCAPAR is a plant killer. Avoid allowing spray to drift over flowering plants, vegetable gardens, etc. The first mowing after treatment should not be delayed as a mulch for flowers and vegetables.

~~be used as a mulch for flowers and vegetables.~~ After
Wash out the spray equipment before using "VIECPAR". After
use, first rinse equipment, including nozzles thoroughly, with water con-
taining a wetting agent or detergent fluid, refill with clean water
and let stand overnight.

Empty spray equipment the following day
Pails stored in building and water sources MUST NOT be contaminated by direct application of washings from spray equipment



Whiz-Kill
MCPP
HICPP (2,4-D)

**HERBICIDE for CONTROL
of BROAD-LEAF WEEDS in TURF, FAIRWAYS and LAWNS**

USDA REGISTRATION No. 728-86

ACTIVE INGREDIENTS:

Diethanolamine salt of 2-(2-methyl-4-chlorophenoxy)propionic acid* 30.1%

Diethanolamine salt of 2,4-dichlorophenoxyacetic acid* 14.9%

INERT INGREDIENTS: 55.0%

*Equivalent to 2 lbs. (21.4%) 2-(2-methyl-4-chlorophenoxy)propionic acid (MCPP) and 1 lb. (10.7%) 2,4 dichlorophenoxyacetic acid (2,4-D) per gallon

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes and clothing. Harmful if swallowed.

After using, wash hands and other parts of the body contaminated with concentrate, spray or washings.

NET CONTENTS:

VINELAND CHEMICAL COMPANY

VINELAND, NEW JERSEY