A Selective Weed Destroyer for

Cotton

ide may

nge yards

GRASS,

GREEN

wetting

plication.

foliage

tion will

domestic

d forage.

ldren and

ed areas.

breathing

at once.

rant that

out make mical Co.

y damage

product.

risks and

onditions.

CRE



ACCEPTED

6/5/67

UNDER THE FEDERAL INSECTICIDE

FUNGICIDE AND RODENTICIDE ACT
FOR LOONOFIC PUISON REGISTERED UNDER NO. T. JOUBLECT
ATTACHED COMMENTS.

 \boldsymbol{A}

Selective

Weed

Destroyer

for

Cotton

LIQUID MSMA + SURFACTANT

FOR CONTROL OF JOHNSON GRASS
AND OTHER WEEDS IN COTTON

ACTIVE INGREDIENT:

Monosodium Acid Methanearsonate

INERT INGREDIENTS:

TOTAL 100.00%

Total Arsenic, as elemental

all in water soluble form 16.34% MSMA per gallon 4.02 lbs.

CAUTION: Keep Out of the Reach of Children.
See side panel for additional cautions.

NET CONTENTS 1 GALLON

Distributed By

TRIANGLE CHEMICAL COMPANY MACON GEORGIA

USDA Reg. No. 2853-33

Printed in U.S.A

35.33%

64.67%

DIRECTIONS FOR USE IN COTTON

DIE-WEED WETTABLE (MSMA plus SURFACTANT) Liquid Herbicide used Post-Emergence for weed control in cotton and similar non-crop areas. WILL CONTROL JOHNSON GRASS, GOOSE GRASS, WATER GRASS, COCKLEBUR, NUT GRASS AND DALLIS GRASS.

DIE-WEED WETTABLE is ready for use, only water need be added. No other wetting agent is necessary. Use in accordance with directions.

FOR BAND TREATMENT OF COTTON GROWN IN 40 INCH ROWS

- (I) Mix at the rate of 1 pint in each 10 gallons of water. Example: Mix 5 pints of the material in 50 gallons of water. Add DIE-WEED WETTABLE to water, thoroughly mix by agitation or recirculating solution in spray tank prior to application.
- (2) Apply at the rate of one (1) gallon of the mixture for each one inch of band width. (13,068 ft. long).

 Example: For 12 inch band, apply 12 gallons of the mixture per acre.
- 43) APPLY AS A DIRECTED SPRAY TO WEEDS WHEN COTTON IS 3 INCHES HIGH AND UP TO FIRST BLOOM. DO NOT APPLY AFTER FIRST BLOOM OPENS

Nozzles on spray equipment should be adjusted to allow thorough coverage of weeds with no contact with cotton foliage. Use fan type nozzle, low volume sprayer. (25 lbs. or less pressure) Clean equipment after use by flushing with water. Bun rinse water in area away from comestic water sources.

Should regrowth occur, a second application will be necessary and may be applied at the above rates 1 to 3 weeks after the first treatment.

NO NOT REUSE CONTAINER

Before using any Herbicide stop and read the label.