

# LYSTADS

# SANITIZE

## DISINFECTANT AND SANITIZER

### INSTRUCTIONS FOR USE

#### EATING AND DRINKING ESTABLISHMENTS:

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with good detergent or compatible cleaner such as MARVEL.
3. Rinse with clean water.
4. Sanitize in a solution of 1 oz. to 4 gals. of water (200 ppm). Immerse all utensils for at least 2 min. or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drain-board to air dry.
6. The State of Wisconsin does not permit rinsing with water after sanitizing.

E.P.A. NO. 2881-20  
NET CONTENTS

### ACTIVE INGREDIENTS:

n-alkyl (C12-40%, C14-50%, C16-10%)  
dimethyl benzyl ammonium chlorides 10.00%  
Tetrasodium ethylenediamine tetraacetate 0.20%

### INERT INGREDIENTS:

89.80%  
100.00%

### DIRECTIONS FOR DILUTIONS

**For Sanitize 200 ppm solution:**  
Add & stir 1 oz. (2 tbs.) Sanitize 10% in 4 gals. water.  
**For Sanitize 400 ppm solution:**  
Add & stir 1 oz. (2 tbs.) Sanitize 10% in 2 gals. water.  
**For Sanitize 800 ppm solution:**  
Add & stir 1 oz. (2 tbs.) Sanitize 10% in 1 gal. water.

## DANGER

### Keep Out of Reach of Children

Severe eye irritant. May cause eye and severe skin irritation. Do not get in eyes or, on skin. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Avoid contamination of food.

**FIRST AID:** In case of eye contact, flush immediately with plenty of water for at least 15 minutes. Get prompt medical attention. In case of skin contact flush skin with plenty of water for at least 15 minutes. If swallowed, drink milk, egg white or large quantities of water. Call a physician.

ANOTHER QUALITY PRODUCT OF

LYSTADS CHEMICALS — GRAND FORKS, N. D. 58201

Milk Farms, Creameries and Food Processing Plants: First clean utensils and equipment with a good detergent, rinse well and then sanitize with Sanitize solution 200 ppm by immersion for 2 minutes. Use same dilution for flank and udder wash prior to milking. Before re-use of equipment coming in direct contact with food, a rinse with safe potable water is recommended.

Clean floors, walls, and storage areas with good detergent, rinse well, and then wipe or mop with Sanitize solution 400 ppm or spray with Sanitize solution 800 ppm.

Hospital and Household Uses: A concentration of at least 800 ppm should be recommended for previously cleaned non-porous surfaces in hospitals. For disinfecting applications in other localities, a concentration of 400 ppm should be recommended for previously cleaned non-porous surfaces and 800 ppm for previously porous surfaces.

Before re-use of equipment coming in direct contact with foods, a rinse with potable water is recommended.

In recommender use dilutions, Sanitize is effective in waters up to 500 ppm hardness when tested by the official procedures.

Rinse empty container thoroughly with water and discard it.

ACCEPTED

9-26-72

UNIVERSITY OF WISCONSIN  
LABORATORY FOR  
FOOD SAFETY  
AND  
HYGIENE  
MADISON, WISCONSIN