CAUTION KEEP OUT OF REACH OF CHILDREN


ilarmful if -wallowed. To not apply directly to tood. Do not apply while tood processing is underway. Thoroughly wash with potable vater treated chutes, processing surfaces, etc. that tood will touch prior to resuse. In case of impracticality of washing, do not allow tood to come in contact with treated surfaces for at least 18 hours. Remove bird cages and fish bowls during spray application. Do not pour, store or spill near heat or open flame. Keep container closed.

HARMFUL TO FISH AVOID SPRAYING OR CONTAMINATING OF PONDS THAT COULD CONTAIN FISH

This spray is especially designed for use in food sterage warehouses, dry milk plants, meat packing plants, flour mills, breweries, potato flaking plants and warehouses, grain bins and elevators, restaurants, grocery stores and other food handling or processing plants.

Do not reuse empty container. Destroy is by perforating or erushing. Bury or discard in a safe place away from water supply.

NET CONTENTS



DIRECTIONS FOR USE

FLEAS (INFESTING PREMISES): Remove and destroy animals' bedding and spray sleeping quarters freely, forcing the spray into all cracks in the floor and about the room. Repeat when re-infested. Do not remain in treated areas, Ventilate the area after treatment is completed.

and crevices in woodwork, underneath sinks, dark places behind pipes, and all places when roaches and waterbugs intest, contacting as many insects is possible. Repeat the operation as often as necessary. Do not remain in treated areas. Ventilate the area after treatment is completed.

CLOTHES MOTHS, CARPET BEETLES: Spray all articles to be protected, giving particular attention to folds and seams, Interior surfaces of storage containers such as closets, chests, and trunks should be sprayed regularly. The treatment should be reneated every 30 days unless the articles are stored in moth-tight containers. Do not remain in treated areas. Ventilate the area after treatment is completed.

GRANARY WEEVIL, RICE WEEVIL, CONFUSED FLOUR BEETLE, GRAIN MITES (ACCESSIBLE STAGES); Wet surfaces thoroughly and treat all infested surface areas, insect runways and hiding places at the rate of I gallon of this material to 1000 square feet of area, or to the point of run-off. Repear when necessary. Do not remain in treated areas. Ventilate the area after treatment is completed.

FOR FOGGING FLIES, GNATS& MOSQUITOES Close all doors and windows. Determine the number of cubic feet of space to be treated. Point nozzle upward and direct the aerosol mist to all parts of the area to be treated, hag particularly well in areas in which insects are most likely to congregate or rest. Fog at the rate of 1½ to 2 ounces per 1000 cubic feet of space. Keep the treated area closed if possible for 15 minutes. After treatment, air out thoroughly and sweep up and remove dead insects. Do not remain in treated areas, Veniclate the area after treatment is completed.