

# LYSTADS

# LISTOL

ACCESSION  
APR 1 1973  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, Lystol 10% is registered under EPA Reg. No. 2881-52

## DISINFECTANT & SANITIZER

### INSTRUCTIONS FOR USE

#### EATING AND DRINKING ESTABLISHMENTS:

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with good detergent or compatible cleaner such as MARVEL.
3. Rinse with clean water.
4. Sanitize in a solution of 1 oz. to 4 gals. of water (200 ppm). Immerse all utensils for at least 2 min. or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drain-board to air dry.
6. The State of Wisconsin does not permit rinsing with water after sanitizing.

EPA Reg. No. 2881-52

EPA Est. 5579-ND-1

NET CONTENTS

### ACTIVE INGREDIENTS:

n-alkyl (C12-40%, C14-50%, C16-10%)  
dimethyl benzyl ammonium chlorides 10.00%  
Tetrasodium ethylenediamine tetraacetate 0.20%

### INERT INGREDIENTS:

(Includes Ethanol less than 5.0%)  
100.00%

### DIRECTIONS FOR DILUTIONS

#### For Listol 200 ppm solution:

Add & stir 1 oz. (2 tbs.) Listol 10% in 4 gals. water.

#### For Listol 400 ppm solution:

Add & stir 1 oz. (2 tbs.) Listol 10% in 2 gals. water.

#### For Listol 800 ppm solution:

Add & stir 1 oz. (2 tbs.) Listol 10% in 1 gal. water.

## DANGER

### Keep Out of Reach of Children

Causes skin irritation. May cause eye damage. Do not get in eyes. Avoid contact with skin. Protect eyes when handling concentrate. For eyes, flush thoroughly and immediately get medical attention. If skin irritation persists, get medical attention. Harmful if swallowed. Avoid contamination of food.

Dilute with water as directed. Do not mix with soap or anionic detergents. Articles to be sanitized should first be washed and rinsed.

MANUFACTURED FOR

LYSTADS CHEMICALS - GRAND FORKS, N. D. 58201

### Milk Farms, Creameries and Food Processing

Plants: First clean utensils and equipment with a good detergent, rinse well and then sanitize with Listol solution 200 ppm by immersion for 2 minutes.

Use 200 ppm for flank and udder wash prior to milking.

Before re-use of equipment coming in direct contact with food, a rinse with safe and potable water is recommended.

Clean floors, walls, and storage areas with good detergent, rinse well, and then wipe or mop with Listol solution 400 ppm or spray with Listol solution 800 ppm.

**Hospital and Household Uses:** A Concentration of at least 800 ppm is recommended for previously cleaned non-porous surfaces such as walls, floors, counter tops, table tops, etc. in hospitals. In disinfection applications in non-hospital areas, a concentration of 400 ppm is recommended for previously cleaned non-porous areas such as floors, walls, table tops, counter tops, etc.

In recommended use dilutions, Listol is effective in waters up to 500 ppm hardness when tested by the official procedures.

Rinse empty container thoroughly with water and discard it.