

2881-56

For control of flying and crawling insects listed in poultry layer and broiler houses.

Flying insects include: House flies, Mosquitoes, Gnats and small flying moths. Crawling insects include: Cockroaches and Grain Mites.

DIRECTIONS FOR USE

This product is to be used in Lystads spraying system. No need to remove birds from houses before spraying. No need to enter poultry house while spraying is going on. To operate unit, turn on switch with air pressure regulated to 40 PSI. Allow unit to run for one minute when two spray head nozzles are used for each 8,000 cubic feet of room space. Each nozzle dispenses approximately 4.2 ounces of insecticide per minute.

NET CONTENTS

EPA Reg. No. 2881-56
EPA Est. 5579-ND-1

LYSTADS PYRISPRAY POULTRY HOUSE SPRAY

CAUTION

Read carefully caution statement on side panel.

KEEP OUT OF REACH OF CHILDREN

PYRISPRAY
1975
Insecticide,
Fungicide Act,
the pesticide
2881-56

ACTIVE INGREDIENTS: 100%
Pyrethrins .30%
Piperonyl butoxide, technical 1.50%
Petroleum distillate 98.20%
*Equivalent to 1.20% (butylcarbityl)(6-propylpiperonyl) ether and .30% related compounds.

CAUTION

Harmful if swallowed. Do not apply directly to poultry food or supplements. Do not apply directly to birds.

Toxic to fish. Do not contaminate any body of water by cleaning of equipment or disposal of wastes.

Do not re-use drum. Return to drum reconitioner or destroy by perforating or crushing and burying in a safe place.

Mix well in drum before sampling and each use. Keep container closed and store in a cool place. DO NOT USE, POUR OR SPILL NEAR HEAT OR OPEN FLAME.

Manufactured for
LYSTADS, INCORPORATED
Grand Forks, N. D. 58201

2881-56

For control of flying and crawling insects listed in poultry layer and broiler houses.

Flying insects include: House flies, Mosquitoes, Gnats and small flying moths. Crawling insects include: Cockroaches and Grain Mites.

DIRECTIONS FOR USE

This product is to be used in Lystads spraying system. No need to remove birds from houses before spraying. No need to enter poultry house while spraying is going on. To operate unit, turn on switch with air pressure regulated to 40 PSI. Allow unit to run for one minute when two spray head nozzles are used for each 8,000 cubic feet of room space. Each nozzle dispenses approximately 4.2 ounces of insecticide per minute.

NET CONTENTS

EPA Reg. No. 2881-58
EPA Est. 5579-ND-1



PYRISPRAY

POULTRY HOUSE SPRAY

CAUTION

Read carefully caution statement on side panel.

KEEP OUT OF REACH OF CHILDREN

P 1975
Insecticide,
Fungicide Act.
the pesticide
2881-56

ACTIVE INGREDIENTS: 100%

Pyrethrins	.30%
Piperonyl butoxide, technical	1.50%
Petroleum distillate	98.20%
*Equivalent to 1.20% (butylcarbityl)(6-propylpiperonyl) ether and .30% related compounds.	

CAUTION

Harmful if swallowed. Do not apply directly to poultry food or supplements. Do not apply directly to birds.

Toxic to fish. Do not contaminate any body of water by cleaning of equipment or disposal of wastes.

Do not re-use drum. Return to drum reconitioner or destroy by perforating or crushing and burying in a safe place.

Mix well in drum before sampling and each use. Keep container closed and store in a cool place. DO NOT USE, POUR OR SPILL NEAR HEAT OR OPEN FLAME.

Manufactured for
LYSTADS, INCORPORATED
Grand Forks, N. D. 58201