


LYSTADS

LISTOL II



DISINFECTANT SANITIZER DEODORIZER

USE DIRECTIONS

HOUSEHOLD

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. Apply solution with mop or cloth. Sanitize with one ounce LISTOL II to 4 gallons of water. Disinfect with 3 ounces LISTOL II to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

HOSPITALS

For floors, walls and other hard surfaces. In nursing homes, institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use LISTOL II at 3½ ounces per 5 gallons of water. Apply with cloth, mop or sponge. At this level LISTOL II will be effective against Pseudomonas aeruginosa PRD-10. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

FOOD PROCESSING EQUIPMENT

For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce LISTOL II per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, NO POTABLE WATER RINSE is required.

DAIRIES

To sanitize dairy equipment such as tanks, lines, pails and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce LISTOL II to 4 gallons of water (200 ppm). At this level, NO POTABLE WATER RINSE is required. Consult local health board for additional information.

RESTAURANT AND BAR RINSE

Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent, rinse thoroughly and immerse in a sanitizing solution containing 1 ounce LISTOL II to 4 gallons of water. NO TERMINAL POTABLE WATER RINSE REQUIRED.

This product when used on environmental, inanimate, hard surfaces at 3½ ounces per 5 gallons of water is effective against Influenza A2, Herpes Simplex, Adenovirus type 5 and Vaccinia Viruses.

ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides 5%
n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides 5%

INERT INGREDIENTS:

90%
100%

LISTOL II is a concentrate and should be diluted before using.

DANGER

Keep Out of Reach of Children

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Do not reuse empty container. Rinse thoroughly with water before discarding or returning drum to drum reconditioner.

COLD DISINFECTION

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1½ ounce LISTOL II per gallon of water for 10 minutes.

FOR ROOM FOGGING

DIRECTIONS:

To reduce the danger of cross-contamination from environmental surfaces, disinfect with LISTOL II as directed. As an adjunct prior to standard routine terminal cleaning and disinfecting of room surfaces, apply solution of LISTOL II as a fog as described below:

1. Remove all human, animal and plant life from room. 2. Open closet doors and drawers. 3. Set up 34" revolving platform in center of room. 4. Mount Jet FOG-MASTER TRI-JET MODEL 6208 or other fogging device delivering equivalent spray at 3 RPM. 5. Fill sprayer reservoir with 1½ ounce of LISTOL II to one gallon of water to produce 1150 ppm active ingredients. 6. Set sprayer mechanism to deliver one gallon of solution. 7. Fog for 15 minutes for an average 2100 cubic foot room. For different room sizes, vary spray time proportionately to ensure complete wetting of exposed surfaces. 8. Wait 2 hours before entering room after treatment. 9. Rinse fogging equipment thoroughly with clear water following use. Room surfaces should then be cleaned and disinfected.

WARRANTY

Our recommendation for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its mis-use as such, or in combination with other materials. Statements concerning the use of these products are not to be construed as recommending the infringement of any patent, and no liability for infringement arising out of any such use is assumed.

EPA REG. NO. 2881-61
EPA Est. 5579-ND-1

NET CONTENTS ONE GALLON

MANUFACTURED FOR
LYSTADS INCORPORATED - GRAND FORKS, NORTH DAKOTA 58201