

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS**

**CAUTION:** Keep away from children, domestic animals and pets. May be irritant or fatal if swallowed. In case of contact, may reduce the effectiveness of the product and cause bleeding.

**NOTE TO PHYSICIAN:** If ingested, administer Vitamin K intramuscularly or orally as indicated in the Physician's Manual. Repeat as necessary based on medical judgment and response.

**ENVIRONMENTAL HAZARDS**

Keep out of lakes, streams or ponds.


**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed. Storage and disposal procedures are printed on the label.

**PESTICIDE DISPOSAL:** Bait that cannot be used should not be stored. It may be disposed of in accordance with applicable local, state, or federal regulations.

**DO NOT REUSE EMPTY CONTAINER WRAP AND PUT IN TRASH COLLECTION**

**WARRANTY:** Lystads makes no warranty, expressed or implied, for use of this product other than as indicated on the label. Buyer assumes all risk of use in the handling of this material when the manufacturer's handling instructions are not followed.



**KILLS RATS AND MICE**

**CAUTION  
KEEP OUT OF REACH OF  
CHILDREN**

See other side for additional information and instructions.

NET WEIGHT

1.00 lb (454 g)

Lystads, Inc., Grand Forks, N.D. 58201

© 1988 Lystads, Inc. All rights reserved.

Printed in the U.S.A.

**DIRECTIONS FOR USE**

For control of Norway rats, Brown rats, and House mice in and around human habitations.

**USE RESTRICTIONS:**

For control of Norway rats, Brown rats, House mice in and around human habitations, commercial, agricultural buildings, and any man-made structure. Lystads Rat and Mouse Killer may also be used inside trailers, vehicles, ships, trains, aircraft and in and around related port or terminal buildings. Treated baits must be placed in locations not accessible to children, pets, domestic animals, wildlife or in tamper proof bait stations. Do not place baits in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS:**

Determine area where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove all food if possible.

**APPLICATION DIRECTIONS**

**RATS:** Apply 11-17 baits without baiting intervals of 15 to 20 days per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity are gone.

**HOUSE MICE:** Apply 15 to 19 baits without baiting intervals of 8 to 12 days per placement. Larger placements up to 25 baits may be needed at plants of very high mouse activity. Maintain uninterrupted supply of fresh bait until signs of mouse activity are gone. Replace contaminated or spoiled bait immediately. Collect and dispose of all unused bait and dead exposed animals according to Container Disposal paragraph to prevent reinfestation. Eliminate food, water and harborage, as much as possible, to prevent reinfestation. Repeat treatment if necessary. Always use bait stations and traps when bait is needed.

ACCEPTED

1988