

*2881-70

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply space sprays while food processing is underway. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions above in those establishments which do NOT operate under Federal meat, poultry, shell egg grading and egg products inspection programs. After spraying in meat packing plants, bakeries and other food processing plants, all equipment, benches, shelving, etc. where exposed food will be handled must be washed with an effective cleaning compound and rinsed with potable water. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SPACE APPLICATIONS: For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths.

Close windows and doors and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. In Official Meat and Poultry Establishments use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. In other than Official Meat and Poultry Establishments a combination of spot surface treatment and fogging is recommended.

IN BARN, POULTRY HOUSES and STORAGE ROOMS: To control house flies, stable flies and horn flies in Barns, Poultry Houses, Warehouses, Storage Bins and Storage Rooms dilute 1 part of material with 4 parts of oil. Apply as a fine mist over the livestock in stanchions or holding pens. Use a fogger or mist generator to direct spray above the animals and to the center of the area where animals are housed. Apply 1 ounce per 1000 cubic feet uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. If used in Dairy Barns, aerate the barn thoroughly before milking. Repeat daily in the morning and afternoon, or as required.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, mealworm, grain mites, and

Lystads
Inc.

**.5% FOGGING
SPRAY**

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional cautions.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician or Poison Control Center immediately. Do not induce vomiting unless directed to do so by a physician. Aspiration is a potential hazard.

IF IN EYES: Flush with plenty of water. Contact a physician if irritation persists.

IF ON SKIN: Wash affected areas with plenty of soap and water.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

**Apply with Equipment Designed and Adjusted to
Deliver a True Aerosol Mist**

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
*Piperonyl Butoxide, Technical	5.0%
INERT INGREDIENTS	94.5%

*Equivalent to 4.0% (butylcarbyl) (6-propylpiperonyl) ether and 1.0% related compounds.

EPA Reg. No. 2881

EPA Est. 2881-ND-1

Manufactured by Lystads, Incorporated
901 University Avenue, Grand Forks, ND 58201

100.0% ACCEPTED
WITH COMMENTS
EPA Letter 10/1/70

SEP 25 1985

2881-70

DIR

... Use a
Direct 1/4" spray
around containers,
shelves and drawers,
neats and points
floors, walls, and
feet of surface.

To control carpenter
and rug. Make
directly into cracks
as necessary.

To control bedbugs
seams. Take bed
mattings, and fl.

To control fleas on
beds, resting on
baseboards, mo
floor and floor
quarters following
proved insecticide.

To control clothes
thoroughly and
liberally to keep
the spray into cr
be treated slight
staining. Store c
thly.

OUTDOOR ADULT MOSQUITOES
treat the entire area
exceed 5 miles
may rest, low m
ly at a rate of 1
feet).

Using truck-mo
to treat a 300 s
minute.

Do not con
PESTICIDE
Store upright
temperature
material suc
with chemi
PESTICIDE
Pesticide, s
ding to labo
waste dispo
CONTAIN
Triple rinse
ing, or pun
proved Stai

Buyer assu
product ne
herewith.