

## CONCENTRATE MUST BE DILUTED BEFORE USE —FOR MANUAL APPLICATION ONLY

## **DIRECTIONS**

LIGHT DUTY CLEANING AND SANITIZATION: One ounce (2 tablespoonsful) of Totil added to four gallons of water will provide 200 ppm active quaternary germicide. Apply solution and allow to soak on surface for 5-10 initiates before rinsing. For residual bacteriostatic activity on the cleaned surface, do not rinse-remove excess cleaning solution with a mop, sponge, or wet vacuum. Rinse food preparation surfaces with potable water before reuse.

HEAVY DUTY CLEANING AND DISINFECTION: One ounce (2 tablespoonsful) of Total added to each gallon of water will provide 780 ppm active quaternary germicice. Follow instructions for Light Duty Cleaning and Sanitization.

Phenol Coefficients (AOAC 20 C): Staphyloccoccus aureus 50, Salmonella typhosa 38

See TOTIL I terature for farm and additional uses.

DANGER: KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or an clothing. Protect eyes and skin when handling Harmful if swallowed. Avoid contamination of food

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a process and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quartity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water Call a physician immediately


NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Do not use Total with soap or other detercents. Rinse empty container thoroughly with - water and discard it

E P.A. Reg. No. 2914-37



CALGON COMMERCIAL DIVISION
7501 PAGE AVENUE ST. LOUIS, MISSOURI 63166



EB 10-092



## STAPHYLOCIDAL/PSEUDAMONACIDAL/SALMONELLACIDAL CLEANS/DISINFECTS/DEODORIZES/SANITIZES

Recommended for cleaning, disinfecting, and deodorizing in hospitals, nursing homes, veterinary clinics, schools and other public institutions, food processing plants, beverage processing plants, restaurants, farms and other areas where the control of bacteria is essential. Effective on hard surfaces such as floors, walls, wood work and metal surfaces.

Recommended for high risk cross-infection areas such as operating rooms, sathology suites, isolation areas and physical therapy suites. Effective as a pre-autoclave sanitizer for surgical instruments, and as a sanitizer for physical therapy equipment. Safe on soft metals.

## Active Ingredients:

n-alkyl (40% $C_{12'}$ 50% $C_{14'}$ 10% $C_{18}$ ) dimethyl
benzyl ammonium chloride
Tetrasodium ethylenediamine tetraacetate
Inert Ingredients: *
*Includes water softening, deodorizing, solubilizing and stabilizing agents

DANGER: KEEP OUT OF REACH OF CHILDREN

(See side panel for first aid and additional precautions.)

**CONTENTS: 55 GALLONS** 

Andrew V. T. Carling