


2915-12



INDUSTRIAL INSECT SPRAY

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
*Technical piperonyl butoxide	4.0%
Petroleum hydrocarbon oil base	10.5%
*Equivalent to 3.2% butylcarbityl (6-propylpiperonyl) ether and 0.8% of other related compounds.	
INERT INGREDIENTS	85.0%



WARNING: CONTENTS UNDER PRESSURE
Keep Out of Reach of Children
Read side panel

INDUSTRIAL INSECT SPRAY is particularly suited for use where food and food products are packed, stored or served, or wherever insects are apt to invade.

DIRECTIONS

Remove cap from top of container. Hold container upright and point spray jet away from you. Press plastic button and treat as follows: Do not remain in treated area and ventilate the area after treatment is completed.

- 1. Flour Beetle, Weevils, Ants, Roaches.** Direct spray toward insects, into cracks, crevices, around machinery, containers of stored products, and other areas which might house insects. A treatment of 30 seconds for every 100 square feet of floor space is recommended. Close treated area tightly for 45 minutes and repeat as required.
- 2. Indian Meal (Chocolate) and Flour Moths.** Close doors and windows. Direct spray to all parts of the room, especially into dark enclosures and around stored product containers where adult moths may be lurking. A treatment of 6 to 8 seconds for every 1,000 cubic feet of space is recommended. Close area for 30 minutes. Repeat weekly or as other flying moths appear.
- 3. Flies, Mosquitoes, Wasps, Gnats.** Determine area to be treated. Shut all doors and windows. Direct aerosol mist to all parts of the room, especially windows and other sources of light which attract these insects. Keep aerosol in constant circular motion. Apply from 3 to 5 seconds for every 1,000 cubic feet.

WARNING: Contents under pressure Exposure to prolonged sunlight or other source of heat may cause bursting. Do not puncture or incinerate when full or empty.

Do not spray direct on foods or food utensils. Food processing plants should not be in operation while spraying. Cover or remove food products and processing surfaces during treatment. Any exposed working surfaces should be washed before plant is returned to production.

Do not use this product where birds and live fish or other aquatic or cold blooded life is kept.

Spraying closer than 36 inches. or spraying longer than

2915-12
 ACCOUNTED
 8/25/71