

# 5% MALATHION DUST

ACTIVE INGREDIENT:

Malathion\*

5.00%

INERT INGREDIENTS:

95.00%

Total — 100.00%

O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

## CAUTION

### KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Avoid prolonged breathing of dust.  
Avoid repeated or prolonged contact with skin. Wash thoroughly.  
Avoid contamination of food and feed stuff.  
This product is toxic to fish.  
Keep out of lakes, streams, and rivers.  
Do not apply near water.  
Do not use in or near homes, schools, or places where people live or work.  
Do not use in or near areas where people or animals graze.  
Malathion is toxic to bees and other beneficial insects. Do not apply to blooming crops or weeds where bees are active.  
Do not use in or near areas where people or animals graze.  
Do not use in or near areas where people or animals graze.

#### NOTE TO PHYSICIANS:

Malathion, upon repeated, prolonged careless use may cause cholinesterase inhibition. Atropine is antidotal.

#### DIRECTIONS FOR USE

**STRAWBERRIES** (Aphids, Spider Mites) — Use 40 lbs. CAROLINA BRAND 5% Malathion dust per acre. Do not apply less than three (3) days before harvest.

**BLUEBERRIES** (Cranberry Fruit Worm, Cherry Fruit Worm, Blueberry Maggot) — Use 25 lbs. CAROLINA BRAND 5% Malathion dust per acre. Do not apply less than three (3) days before harvest.

**BEANS** (Mexican Bean Beetle, Leafhopper) — Use 30 to 35 lbs. CAROLINA BRAND 5% Malathion dust per acre. Do not apply less than three (3) days before harvest.

**BROCCOLI, CABBAGE, KALE, TURNIPS, MUSTARD** (Aphids, Imported Cabbage Worm, Cabbage Looper) — Use 30 lbs. CAROLINA BRAND 5% Malathion dust per acre. Do not apply less than seven (7) days before harvest.

**TOBACCO** (Aphids) — Use 20 to 30 lbs. CAROLINA BRAND 5% Malathion dust per acre. Do not apply less than seven (7) days before harvest.

#### NOTICE: Conditions of Sale

**CARTER INSECTICIDE & CHEMICAL CO., INC.**

P. O. Box 209

Wallace, N. C. 28466

Phone AT 5-2144

E-2367

USDA REG. NO. 2917-31