

2% Parathion Dust

ACTIVE INGREDIENT:

Parathion* 2.0%

INERT INGREDIENTS:

..... 98.0%

TOTAL — 100.0%

*O, O-diethyl O-p-nitrophenyl thiophosphate

DANGER

Danger!



Poison

KEEP OUT OF REACH OF CHILDREN & ANIMALS

May Be Fatal If Swallowed, Inhaled, or Absorbed Through Skin! Rapidly Absorbed Through Skin! Do not get in eyes or on skin. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. ~~Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Burn or destroy, never re-use, containers, stay out of smoke or fumes.~~

Danger!



Poison

This product is toxic to fish and wildlife.

Keep out of lakes, streams, and ponds.

Apply this product only as specified on the label.

Do not apply when weather conditions favor drift or carry dust from areas treated.

Rinse empty metal containers and dispose of wastes by burying in non-crop lands away from water supplies. Metal containers should be disposed of by punching holes in them and burying with wastes. Glass containers should be disposed of by breaking & burying with wastes. Paper or cardboard containers should be disposed of by burning (Keep out of smoke).

DIRECTIONS FOR USE

TOBACCO: Green Peach Aphid and Flea Beetles in tobacco plant beds, make 1 to 2 applications using ½ lb. per 100 sq. yards of plant bed area. Be sure material is uniformly applied over plants. **HANDLERS OF TREATED SEED BED PLANTS SHOULD WEAR RUBBER GLOVES.**

Green Peach Aphid and Flea Beetles in tobacco fields, apply 7 to 12 lbs. 2% Parathion per acre. **APPLICATIONS SHOULD NOT BE MADE WITHIN 5 DAYS OF PRIMING OR 15 DAYS OF CUTTING. AVOID PLANT JUICES COMING IN CONTACT WITH THE SKIN OR OTHER PARTS OF THE BODY OF THOSE WHO ARE ENGAGED IN PRIMING OR CUTTING THE CROP**

COTTON: Aphids, make 1 application of 12 to 15 lbs. per acre 2% Parathion when honeydew first appears and leaves begin to curl downward. **DO NOT APPLY LESS THAN 30 DAYS BEFORE HARVEST.**

CABBAGE, COLLARDS, BROCCOLI: Aphids, apply 10 to 12 lbs. 2% Parathion dust per acre when infestation first appears. **DO NOT APPLY WITHIN 30 DAYS OF HARVEST OR AFTER HEADS OR OTHER EDIBLE PARTS HAVE BEGUN TO FORM.**

POISON

ATROPINE IS ANTIDOTAL

KEEP AN EMERGENCY SUPPLY ON HAND

FIRST AID TREATMENT

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED PARATHION POISONING

SYMPTOMS: See physician's note below.

INHALATION: Remove patient from contaminated atmosphere. Keep him under continuous observation. Don't wait for doctor; give atropine (2 tablets) at once if symptoms include blurred vision or pin-point pupils.

SKIN: Immediately remove contaminated clothing and scrub skin with soap and water for 15 minutes. Don't wait for doctor; give atropine (2 tablets) at once if symptoms include blurred vision or pin-point pupils.

INTERNAL: If swallowed, make patient vomit by giving warm salty or soapy water. Don't wait for doctor; give atropine (2 tablets) at once if symptoms include blurred vision or pin-point pupils.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PATIENT.

PHYSICIAN'S NOTE: Warning symptoms include weakness, headache, chest pain, blurred vision, pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and cramps.

TREATMENT: Give atropine, preferably by injection, grains, 1/100 two or three tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. Never give morphine. Clear chest by postural drainage. Give artificial respiration if patient becomes comatose. Oxygen may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood sample tests.

NOTICE: Conditions of Sale

This product is sold as warranted, and unless otherwise stated herein, the product sold hereunder is warranted to be of merchantable quality. The manufacturer makes no representation or warranty of any kind, express or implied, with respect to the product. Any claim for damages or other loss shall be deemed waived unless made in writing and received by the Manufacturer within 30 days after Buyer's receipt of the product.

— Manufactured By —

CARTER INSECTICIDE & CHEMICAL CO., INC.

P. O. Box 209

Wallace, N. C. 28466

Phone AT 5-2144

E-1962-A

USDA Reg. No. 2917-46

ACCEPTED
7/12/67