

MAY 27 1967

LINDANE DUST

ACTIVE INGREDIENTS:

Gamma Isomer of Benzene Hexachloride (From Lindane) 1.0%

INERT INGREDIENTS

99.0%

TOTAL — 100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

This material is poisonous if swallowed or absorbed through the skin. It is also very irritating to the eyes. Wash thoroughly with soap and water after handling. Do not breathe dust. Avoid contact with food.

This material is not to be used on food crops. Keep out of reach of children and animals. Apply this product only as directed.

Do not use this material on food crops. Keep out of reach of children and animals. Apply this product only as directed. Do not use this material on food crops. Keep out of reach of children and animals. Apply this product only as directed.

DIRECTIONS FOR USE

For control of melonworm, pickleworm, striped and spotted cucumber beetle on cantaloupe, cucumber and squash: Apply 25 to 30 lbs. Carolina Brand P-6 Lindane Dust per acre. REMOVE INSECTS FROM EDIBLE PARTS BY WASHING BEFORE USING FOR FOOD.

For control of cucumber beetle on ~~cucumbers~~. Use 25 to 30 lbs. Carolina Brand P-6 Lindane Dust per acre. DO NOT APPLY AFTER THE EDIBLE PARTS START TO FORM. For control of cucumber beetle on squash: Apply 25 to 30 lbs. Carolina Brand P-6 Lindane Dust per acre.

REGISTERED U.S. PATENT OFFICE

— Manufactured By —

CARTER INSECTICIDE & CHEMICAL CO., INC.

P. O. Box 209

Wallace, N. C. 28466

Phone AT 5-2144

E 2823 A

USDA REG. NO. 2917-47

MAY 25 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED IN NO. 2917-47

1% LINDANE DUST

ACTIVE INGREDIENTS:

Gamma Isomer of Benzene Hexachloride (From Lindane) 1.0%

INERT INGREDIENTS 99.0%

TOTAL — 100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes or on skin. Wash thoroughly with soap and water after handling this material. Do not contaminate feed or foodstuffs. Avoid breathing of dust.

This product is toxic to fish and wildlife.
Keep out of lakes, streams, and ponds.
Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Metal containers should be disposed of by punching holes in them and burying with wastes. Glass containers should be disposed of by breaking & burying with wastes. Paper or cardboard containers should be disposed of by burning. Keep out of smoke.

DIRECTIONS FOR USE

For control of mellonworm, pickleworm, striped and spotted cucumber beetle on cantaloupe, cucumber and squash: Apply 25 to 30 lbs. Carolina Brand 1% Lindane Dust per acre. REMOVE EXCESS RESIDUE FROM EDIBLE PARTS BY WASHING BEFORE USING FOR FOOD OR FEED.

For control of lace bugs on eggplant: Use 25 to 30 lbs. Carolina Brand 1% Lindane Dust per acre. DO NOT APPLY AFTER THE EDIBLE PARTS START TO FORM.
Do not apply within 1 day of harvest for cucumbers and squash.

NOTICE: Conditions of Sale

Manufacturer-Seller warrants that, unless otherwise specified herein, the products sold hereunder shall meet Manufacturer-Seller's standard specifications and shall be of merchantable quality. Except as otherwise expressly stated herein, Manufacturer-Seller makes no representation or warranty of any kind, express or implied, with respect to the product. All claims for alleged defective products, shortage, or other cause shall be deemed waived unless made in writing and received by Manufacturer-Seller within 20 days after Buyer's receipt of the product.

— Manufactured By —

CARTER INSECTICIDE & CHEMICAL CO., INC.

P. O. Box 209

Wallace, N. C. 28466

Phone AT 5-2144

E-2823-A

USDA REG. NO. 2917-47