


CUKE DUST



ACTIVE INGREDIENT:

Gamma Isomer of Benzene Hexachloride (from Lindane) 1%

Copper (Expressed as Metallic in the form of Tri Basic
Copper Sulphate) 5%

INERT INGREDIENTS: Not more than 94%

TOTAL—100%

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Avoid prolonged breathing of dust or spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

DIRECTIONS

CUCUMBERS: Downy Mildew, Striped and spotted cucumber beetles, melonworm and pickleworm, apply with a good duster at rate of 20 to 35 lbs. per acre when disease or insects first appear. Repeat applications twice a week until infestation is under control. **DO NOT APPLY LESS THAN ONE DAY BEFORE HARVEST.**

NOTICE: Conditions of Sale

Manufacturer Seller warrants that the product, as otherwise specified herein, the product sold hereunder shall meet Manufacturer Seller's standard of quality and shall be of merchantable quality, but Manufacturer Seller makes no warranty as to the results of the product's use, either alone or in combination with other substances. Seller's liability for the product is limited to the extent of the remedy and Manufacturer Seller's limit of liability for any loss or damage, including consequential loss or damage, shall be deemed waived unless made in writing and received by Manufacturer Seller within twenty days after Buyer's receipt of the product. Buyer's remedy and Manufacturer Seller's limit of liability for any loss or damage, including consequential loss or damage, shall be for the purchase price of the particular delivery, which is subject to which losses or damages are claimed plus any transportation charges paid by Buyer. Manufacturer Seller shall not be liable for, and Buyer assumes responsibility for, all personal injury and property damage resulting from the handling, possession or use of the product by Buyer.

CUKE DUST

ACTIVE INGREDIENT:

Gamma Isomer of Benzene Hexachloride (from Lindane)	1%
Copper (Expressed as Metallic in the form of Tri Basic Copper Sulphate)	5%

INERT INGREDIENTS: Not more than 94%

TOTAL—100%

DIRECTIONS

CUCUMBERS: Downy Mildew, Striped and spotted cucumber beetles, melonworm and pickle worm, apply with a good duster at rate of 20 to 35 lbs. per acre when disease or insects first appear. Repeat applications twice a week until infestation is under control. DO NOT APPLY LESS THAN ONE DAY BEFORE HARVEST.