

2917-67

Parathion (O,O-diethyl-O-p-nitrophenyl thiophosphate)

4.00%

96.00%

TOTAL — 100.00%

Danger!

KEEP OUT OF REACH OF CHILDREN AND ANIMALS *Danger!*

Danger!



Poison

Poison

[illegible]

DIRECTIONS:
TOPACCO For control of southern Potato Wire Worm, apply 250 pounds per acre of CALIFORNIA BRAND 1 PARATHION GRANULAR per acre after land has been tilled with a rotary plow. Bury into soil immediately. Treatment should be made about two weeks prior to planting. DO NOT APPLY WITH NAKED HANDS. READ WARNING ABOVE BEFORE APPLYING.

ATROPINE + ANTHOLAM

KEEP AN EMERGENCY SUPPLY ON HAND
FIRST AID TREATMENT

THE APPLICABLE LAW OF THE INALIENABLE RIGHTS OF THE PEOPLE

SYMPTOMS: see table, which is not boxed

INTERLUDE: In your patient room, take vital signs, then, if you are under center, stand observation. Don't wait for doctor. Call a nurse (2nd place) at once if you notice any problems. If you are in center, stand observation.

4. Rinse with warm water. Remove contents from container and let's rub with the soap and water for 15 minutes. Don't wait for doctor, you can cure it yourself at the 2 symptoms. I heard heard vision on bright light.

1. PNAH-16: 100% water, about 100°C, water warm, salty or soapy water. Do not wait for deicing, just measure and take the readings and be prepared to record readings.

STAYING CALM AND IN CONTROL BY MOUTH TO AN UNCONTROLLED PATIENT

[illegible][illegible]

NOTICE: Conditions of Sale

[illegible]

UNDER THE FEDERAL INSECTICIDE ACT
REGISTERED
2917-64

4% PARATHION GRANULAR

ACTIVE INGREDIENT:

Parathion (O,O-diethyl-O-p-nitrophenyl thiophosphate) 4.00%

INERT INGREDIENTS 96.00%

TOTAL — 100.00%

WARNING

Danger!



Poison

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

May Be Fatal If Swallowed, Inhaled, or Absorbed Through Skin! Rapidly Absorbed Through Skin! Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of type passed by the U. S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where they may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Burn or destroy, never re-use, containers, stay out of smoke or fumes.



Poison

DIRECTIONS

TOBACCO: For control of Southern Potato Wire Worm, broadcast uniformly 50 lbs. of CAROLINA BRAND 4% PARATHION GRANULAR per acre, after land has been turned with bottom plow. Disc into soil immediately. Treatment should be made about two weeks prior to transplanting. DO NOT APPLY WITH NAKED HANDS. READ WARNING ABOVE BEFORE APPLYING.

POISON

**ATROPINE IS ANTIDOTAL
KEEP AN EMERGENCY SUPPLY ON HAND
FIRST AID TREATMENT**

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED PARATHION POISONING

SYMPTOMS: See physician's note below.
INHALATION: Remove patient from contaminated atmosphere. Keep him under continuous observation. Don't wait for doctor; give atropine (2 tablets) at once if symptoms include blurred vision or pin-point pupils.
SKIN: Immediately remove contaminated clothing and scrub skin with soap and water for 15 minutes. Don't wait for doctor; give atropine (2 tablets) at once if symptoms include blurred vision or pin-point pupils.
INTERNAL: If swallowed, make patient vomit by giving warm salty or soapy water. Don't wait for doctor; give atropine (2 tablets) at once if symptoms include blurred vision or pin-point pupils.
NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PATIENT.
PHYSICIAN'S NOTE: Warning symptoms include weakness, headache, chest pain, blurred vision, pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and cramps.
TREATMENT: Give atropine, preferably by injection, grains, 1/100 two or three tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. Never give morphine. Clear chest by postural drainage. Give artificial respiration if patient becomes comatose. Oxygen may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood sample tests.

NOTICE: Conditions of Sale

Manufacturer/Seller warrants that, unless otherwise specified herein, the product sold hereunder shall meet Manufacturer/Seller's standard specifications and shall be of merchantable quality but Manufacturer/Seller makes no warranty as to the fitness of the product for a particular purpose or the results to be obtained from its use by Buyer either alone or in combination with other substances. Subject to the preceding sentence and except as otherwise expressly stated herein, Manufacturer/Seller makes no representation or warranty of any kind express or implied, with respect to the product. All claims for alleged defective product, shortage or other cause shall be deemed waived unless made in writing and received by Manufacturer/Seller within twenty days after Buyer's receipt of the product. Buyer's exclusive remedy and Manufacturer/Seller's limit of liability for any and all losses or damages resulting from a defective product or from any other cause shall be for the purchase price of the particular delivery with respect to which losses or damages are claimed plus any transportation charges paid by Buyer. Manufacturer/Seller shall not be liable for, and Buyer assumes responsibility for, all personal injury and property damage resulting from the handling, possession or use of the product by Buyer.