

7 1/2% SEVIN[®] DUST

2917-68

3-13-61

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)

7.50%

INERT INGREDIENTS

92.50%

TOTAL — 100.00%

Trademark of Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate.

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

HARMFUL TO HUMANS IF SWALLOWED. Do not inhale dust. Do not get internally. Skin Contact With Any Insecticide May Be Harmful. Wash thoroughly with soap and water after use. Do not eat, drink, or smoke while using this product. Do not use in or near buildings.

Note for Physician: Carbaryl is a carbamate. It may cause drowsiness, dizziness, and other symptoms. Avoid contamination of foodstuffs. Avoid contact with eyes, nose, mouth, and skin. If in contact with eyes, flush with water.

DIRECTIONS

TOBACCO: Budworms, Flea Beetles, Hand Dust Application: When hand dust applications are made on individual plants, apply 6 to 8 pounds per acre. Heavier applications may cause plant injury. Rotary dusters are recommended. Shaker cans and cloth bags usually result in excessive applications.

TOBACCO: Hornworms, Budworms, Japanese Beetles, Grasshoppers, Flea Beetles, Broadcast Application: Apply 15 to 25 pounds per acre depending on size of the plants. Use the highest rate only on large plants. Full coverage is essential. Begin treatment when worms are small and repeat as needed. Heavier applications may cause plant injury. **NOTE: (1) If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. (2) Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.**

COTTON: For early-season thrips, flea beetle, and cotton fleahopper. Also for cotton leafworm and striped blister beetle. Apply 15 to 20 pounds of CARTER CAROLINA BRAND 7 1/2% Dust per acre on appearance of insects.

For mid- and late-season boll weevil, bollworm, fall armyworm, cotton leaf perforator, leaf rollers, leafhoppers and turnip leaf plant bug. Apply 15 to 20 pounds of CARTER CAROLINA BRAND 7 1/2% Dust per acre, depending on the size of the cotton and insect population level. Applications for boll weevil should be made every five to seven days as long as control is necessary.

Aphid populations will be suppressed by repeated applications of this insecticide.

SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide.

Consult your local county agent or agricultural authorities for more specific recommendations for your locality.

ALLOW 7 DAYS BETWEEN LAST APPLICATION AND GRAZING TREATED FIELDS OR HARVESTING CROP RESIDUE FOR USE AS FEED OR BEDDING FOR DAIRY OR MEAT ANIMALS.

NOTICE: Coded by sale

This product is sold by the manufacturer in accordance with the provisions of the Federal Insecticide, Fungicide, and Rodenticide Act. The manufacturer warrants that the product is of the quality and quantity represented on the label. The manufacturer does not warrant the product for any other purpose. The manufacturer is not responsible for any damage to property or person caused by the use of this product. The manufacturer is not responsible for any damage to property or person caused by the use of this product. The manufacturer is not responsible for any damage to property or person caused by the use of this product.

7 1/2% SEVIN DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)

INERT INGREDIENTS

2917-68
3-13-67
AC 100
2917-68
3-13-67
FOR
BY
TO

7.50%

92.50%

TOTAL — 100.00%

*Trademark of Union Carbide Corp for 1-Naphthyl N-Methylcarbamate.

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. **Skin Contact With Any Insecticide May Be Harmful.** Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved clothing. Change to clean clothing daily.

Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptacles.

DIRECTIONS

TOBACCO: Budworms, Flea Beetles, Hand Dust Application: When hand dust applications are made on individual plants, apply 6 to 8 pounds per acre. Heavier applications may cause plant injury. Rotary dusters are recommended. Shaker cans and cloth bags usually result in excessive applications.

TOBACCO: Hornworms, Budworms, Japanese Beetles, Grasshoppers, Flea Beetles, Broadcast Application: Apply 15 to 25 pounds per acre depending on size of the plants. Use the highest rate only on large plants. Full coverage is essential. Begin treatment when worms are small and repeat as needed. Heavier applications may cause plant injury. **NOTE: (1) If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. (2) Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.**

COTTON: For early-season thrips, flea beetle, and cotton fleahopper. Also for cotton leafworm and striped blister beetle: Apply 15 to 20 pounds of CARTER CAROLINA BRAND 7 1/2% Dust per acre on appearance of insects.

For mid- and late-season boll weevil, bollworm, fall armyworm, cotton leaf perforator, leaf rollers, leafhoppers and tarnished plant bug: Apply 15 to 20 pounds of CARTER CAROLINA BRAND 7 1/2% Dust per acre, depending on the size of the cotton and insect population level. Applications for boll weevil should be made every five to seven days as long as control is necessary.

Aphid populations will be suppressed by repeated applications of this insecticide.

SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide.

Consult your local county agent or agricultural authorities for more specific recommendations for your locality.

ALLOW 7 DAYS BETWEEN LAST APPLICATION AND GRAZING TREATED FIELDS OR HARVESTING CROP RESIDUE FOR USE AS FEED OR BEDDING FOR DAIRY OR MEAT ANIMALS.

NOTICE: Conditions of Sale

Manufacturer/Seller warrants that, unless otherwise specified herein, the product sold hereunder shall meet Manufacturer/Seller's standard specifications and shall be of merchantable quality, but Manufacturer/Seller makes no warranty as to the fitness of the product for a particular purpose or the results to be obtained from its use by Buyer either alone or in combination with other substances. Subject to the preceding sentence and except as otherwise expressly stated herein, Manufacturer/Seller makes no representation or warranty of any kind express or implied, with respect to the product. All claims for alleged defective product, shortage or other cause shall be deemed waived unless made in writing and received by Manufacturer/Seller within twenty days after Buyer's receipt of the product. Buyer's exclusive remedy and Manufacturer/Seller's limit of liability for any and all losses or damages resulting from a defective product or from any other cause shall be for the purchase price of the particular delivery with respect to which losses or damages are claimed plus any transportation charges paid by Buyer. Manufacturer/Seller shall not be liable for, and Buyer assumes responsibility for, all personal injury and property damage resulting from the handling, possession or use of the product by Buyer.

E-1037-A

USDA REG. NO. 2917-68