



DUSTING SULFUR

ACCEPTED	
CERTIFICATE OF ANALYSIS	
No. 2935-48-AA	
DATE: 10/15/54	
BY: [Signature]	
LABORATORY: [Signature]	
TESTED AT: [Signature]	
TOTAL 100%	

DIRECTIONS FOR USE: This material may be applied with ground equipment or aircraft depending upon local conditions.

RECOMMENDATIONS FOR USE:

ALMONDS, CHERRIES, NECTARINES, PEACHES, PLUMS and PRUNES: For the control of BROWN ROT, CHERRY LEAF SPOT, PRUNE RUST and POWDERY MILDEW use at the rate of 40 to 50 lbs. per acre in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

ALFALFA: For the control of POWDERY MILDEW, TWO SPOTTED MITE, PACIFIC MITE, and STRAWBERRY MITE (Atlantic Mite) use 30 to 50 lbs. per acre. Apply when disease first appears and repeat at 14 day intervals or as necessary.

ASPARAGUS: For control of RUST use 15 to 35 lbs. per acre. Dust after cutting stops, irrigate, cultivate, then apply the sulfur. Repeat at 7-10 day intervals through season.

APPLES and PEARS: For control of SCAB and POWDERY MILDEW use 40 to 50 lbs. per acre. Apply in pre-bloom, bloom and post-bloom periods. On pears do not apply after the fruit begins to turn down.

BEANS and PEAS: For the control of POWDERY MILDEW use 15 to 30 lbs. per acre. Apply when disease first appears and repeat as necessary.

CARROTS and ONIONS: For the control of PETROBIA MITE use 30 to 35 lbs. per acre.

CITRUS: For the control of CITRUS THRIPS, CITRUS FLAT MITE, YUMA SPIDER MITE, SIX SPOTTED MITE and CITRUS RUST MITE use 100 to 125 lbs. per acre. Apply when pests first appear and repeat as necessary.

COTTON: For the control of STRAWBERRY MITE (Atlantic Mite) use 30 to 40 lbs. per acre when mites first appear and repeat as necessary.

FIGS: For Control of FIG RUST MITE in spring and early summer. Dust at 40 lbs. per acre when damage to foliage (russeting) appears. A second application may be necessary.

GRAPES: For control of POWDERY MILDEW use 8 to 20 lbs. per acre. Make first application when shoots are 6 in. to 8 in. long. Make a second application when shoots are 12 in. to 15 in. long and third about 14 days later. Repeat applications at 10 to 14 day intervals to protect new growth. Avoid over-applications in hot weather.

Concord and other labrusca type grapes may be injured by sulfur.

EPA Reg. No. 2935-48-AA

NOTE: The statements made on this label are based on the best available data, but because of conditions of use which are beyond our control, WILBUR ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation of any kind concerning this material or the use thereof, except in conformity with the statements on this label. Neither WILBUR ELLIS COMPANY, nor the seller, shall be held responsible in any manner for any persons, firms or corporations, damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and agrees to hold the material on these conditions.

ACTIVE INGREDIENT:
Sulfur

INERT INGREDIENTS

98%
2%
TOTAL 100%

EPA Reg. No. 2935-48-AA

CAUTION: KEEP OUT OF REACH OF CHILDREN!

Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

CAUTION: Applications of sulfur should not be made just prior to or during periods of high summer temperatures. Do not make applications within 2 weeks of an oil spray. Do not use or allow this material to drift onto sulfur sensitive Melons, Squash, Cucumbers, Apricots or other plants which may be injured by sulfur.

RECOMMENDATIONS FOR USE: (Continued)

MELONS: Use 20 to 25 lbs. per acre for control of POWDERY MILDEW. Powdery Mildew may be controlled by dusting but the sulfur often injures the vines, particularly if the temperature reaches 95° F. Treat when disease first appears. Use 30-50 lbs. per acre to control Spider Mites. DO NOT USE ON SULFUR SENSITIVE VARIETIES.

OLIVES: For control of OLIVE MITE. Apply 50 to 70 lbs. per acre. Do not use sulfur in hot weather as damage may result to crop and foliage.

ONIONS: For control of PETROBIA MITES and POWDERY MILDEW, use 10 to 40 lbs. per acre. This product may be applied by air or ground equipment. Apply when disease first appears and repeat at 14 day intervals or as necessary.

STRAWBERRIES, RASPBERRIES and BOYSENBERRIES: For the control of POWDERY MILDEW use 10 to 20 lbs. per acre. Apply when disease first appears and repeat as necessary.

SUGAR BEETS: For the control of POWDERY MILDEW, TWO SPOTTED MITE, PACIFIC MITE, and STRAWBERRY MITE (Atlantic Mite) use 35 to 50 lbs. per acre. Apply when disease first appears and repeat at 14 day intervals or as necessary.

TOMATOES: For the control of TOMATO RUSSET MITE use 30 to 40 lbs. per acre.



WILBUR-ELLIS COMPANY



FRESNO · LOS ANGELES · GLENDALE · PORTLAND · SEATTLE