

red-top

MALATHION 25 SPRAY POWDER

COMPOSITION:

ACTIVE INGREDIENT:

Malathion* 25%

INERT INGREDIENTS 75%

*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate

USDA Reg. No. 2935-84

KEEP OUT OF REACH OF CHILDREN!

CAUTION: Harmful if Swallowed. Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.

DIRECTION FOR USE: Recommendations on this label are based on a volume of 500 gallons of water per acre, unless otherwise stated. For a lower volume of water per acre multiply the recommended dosage by 5 to obtain maximum amount of Malathion allowed per acre. Consult local agricultural authorities for additional information, as the timing, number and rate of applications needed will vary with local conditions.

Do not use this product for any purpose other than that recommended on this label.

Do not allow this product to drift from the area being treated.

NOTICE: The statements made on the label, or by any of our agents, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except that it conforms to the chemical description on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or for any other person from the handling, storage or use of the material, whether or not in accordance with directions. The buyer assumes all risk of liability resulting from such handling, storage or use and accepts and uses this material on these conditions.

WILBUR - ELLIS COMPANY

FRESNO - LOS ANGELES - PHOENIX - PORTLAND - SEATTLE - SALT LAKE

Boise - Browley - Chowchilla - Chula Vista - Livingston - Medford

San Joaquin - Stockton - Wandland - Yakima - Yuma

E-565-84 B

ACCEPTED WITH COMMENTS

RECOMMENDATIONS FOR USE:

APPLES, APRICOTS, AVOCADOS, CHERRIES, FILBERTS, NECTARINES, PEACHES, PEARS, PECANS, PLUMS, PRUNES & QUINCES: Use 3 to 4 lbs. per 100 gal. of water to control APHIDS, THRIPS, SPIDER MITES, MEALYBUGS, CODLING MOTH, BIRD MOTH, EYE SPOTTED MOTH, PLUM CURCULIO, CHERRY FRUIT FLY, PEAR PSYLLA, SCALE CRAWLERS, LEAFROLLERS & ORANGE TORTRIX. For WILLAMETTE MITE and TWO-SPOTTED MITES on Apples make at least 2 applications 10 to 12 days apart in summer months. For SCALE on Peaches apply at completion of hatching, about last of June. For CHERRY FRUIT FLY repeat applications every 10 days. Injury may occur to McIntosh and Courtland varieties of Apples in summer sprays, and to certain varieties of sweet Cherries particularly in the Northwest, and fruit spotting may occur on nectarines. Do not apply within 1 day of harvest on Pears; within 3 days on Apples, Cherries, Plums, Prunes and Quinces; within 7 days on Apricots, Avocados, Nectarines and Peaches. No limitation on Filberts and Pecans.

ASPARAGUS, BEANS, BEETS, CARROTS DANDELIONS, EGGPLANT, ENDIVE, GARLIC, LEEKS, PARSLEY, PARSNIPS, PEAS, PEPPERS, SALSIFY, SHALLOTS, SPINACH, SWISS CHARD, & WATERCRESS: Use 4 to 5 lbs. per acre to control APHIDS, SPIDER MITES, THRIPS, LEAFHOPPERS, ASPARAGUS BEETLE, PEA WEEVIL & MEXICAN BEAN BEETLE. Do not apply within 1 day of harvest on Asparagus or Beans; within 3 days on Eggplants, Garlic, Leeks, Peas, Peppers and Shallots, within 7 days on Beets, Carrots, Dandelions, Endive, Parsley, Parsnips, Salsify, Swiss Chard, Spinach and Watercress.

BERRIES: CANEBERRIES (Blackberries, Boysenberries, Dewberries, Loganberries and Raspberries), CRANBERRIES and STRAWBERRIES: Use 5 to 6 lbs. per acre to control APHIDS, THRIPS, SPIDER MITES, WHITEFLIES, LEAFHOPPERS, ROSE SCALE, LYGUS BUGS, SPITTLEBUGS, LEAFROLLERS, BLACK HEADED FIREWORMS and STRAWBERRY ROOT WEEVIL. For STRAWBERRY ROOT WEEVIL apply to soil surface before planting and work into top 6 to 8 inches. Do not apply within 3 days of harvest on Cranberries or Strawberries and within 1 day on Caneberries.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, KALE, MUSTARD GREENS, & TURNIPS: Use 10 lbs. per acre to control APHIDS, IMPORTED CABBAGEWORM and CABBAGE LOOPER. Do not apply within 7 days of harvest on Brussels Sprouts, Cabbage, Kale, Mustard Greens and within 3 days on Broccoli or Turnips.

CELERY: Use 6 lbs. per acre to control APHIDS and SPIDER MITES. Do not apply within 7 days after harvest.

CITRUS: (Grapefruit, Lemons, Limes, Oranges, Tangerines, Tangelos and Kumquats): Use 8 lbs. per 100 gallons of water to control APHIDS, THRIPS, and SCALE CRAWLERS. Do not apply when trees are in bloom. Do not apply within 7 days of harvest.

GRAPES: Use 6 lbs. per acre to control GRAPE LEAFHOPPERS, MEALYBUGS and SPIDER MITES. Injury may occur to Bibier, Italia, Cardinal and Almeria grape varieties when Malathion is applied after the clusters appear. Do not apply within 3 days of harvest.

LETTUCE: Use 8 lbs. per acre to control APHIDS, CABBAGE LOOPER, LEAFHOPPERS and SPIDER MITES. Do not apply within 7 days of harvest on Heart Lettuce and 14 days on Leaf Lettuce.

PEPPERMINT and STEARMIN: Use 4 lbs. per acre to control APHIDS, SPIDER MITES, FLEABETTLER ADULTS and LEAFHOPPERS. Do not apply within 7 days of harvest.

POTATOES and TOMATOES: Use 8 lbs. per acre to control APHIDS, LEAFHOPPERS, BAISE, CHINCH BUG and SPIDER MITES. Do not apply within 1 day of harvest on Tomatoes and no time limitation on Potatoes.