

P1021

2935-92

182

RED-TOP

SPRAY SULFUR

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing for 24 hours after application. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should specify protective clothing, areas or fields that cannot be entered without this clothing, period of time area is to be vacated, and statement of practical treatment in case of exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "Caution: Area treated with Sulfur on (date of application). Do not enter without appropriate protective clothing for 24 hours (insert here action to take in case of accidental exposure)".

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food, or feed by storage or disposal. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

3. **Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

4. **General:** Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing, or pouring.

Some burning may occur when used during warm weather.

Sulfur will cause severe fruit and leaf injury to sulfur sensitive crops. Use only as directed on label.

Do not apply when shade temperatures exceed or are likely to exceed 90° F. Sulfur may burn foliage and fruit during periods of high temperature and under certain climatic conditions. Do not use sulfur within 30 days of an oil spray (60 days on citrus). If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulfur.

Pour recommended amount of this material through screen on surface of water in nearly filled spray tank, or pre-mix in a bucket half filled with water and pour mixture into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low-volume gear type spray equipment. Do not combine with emulsifiable liquids in the same tank.

This product is formulated for application by ground or air equipment.

Consult with State Agricultural Experiment Station or State Agricultural Extension Service personnel for additional information consistent with label cautions and directions.

For control of mites, use higher rates shown under conditions of high pest pressure. For control of fungi, use higher rates shown when weather conditions indicate a potential for increased pest pressure.

RECOMMENDATIONS FOR USE:

APPLES: SCAB, POWDERY MILDEW — Apply 4.5 lbs per 100 gallons from pink or pre-blossom spray through second cover spray. Add 1 gallon lime sulfur in pink and calyx sprays. Use on sulfur tolerant varieties only. Macintosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulfur under certain climatic conditions.

BEANS: For the control of SPIDER MITE and POWDERY MILDEW, use 4 to 5 lbs per 100 gallons of water. Sulfur may injure certain varieties of beans. Begin application when disease first appears. Repeat as necessary.

CHERRIES: BLOSSOM BLIGHT, BROWN ROT, LEAF SPOT, POWDERY MILDEW — Apply 4.5 lbs per 100 gallons. Begin in pink stage. Repeat at bloom, petal fall and in cover stages at 7 to 14 day intervals and as a post harvest foliar treatment for LEAF SPOT.

COTTON: For control of STRAWBERRY (Allant) MITES, use 3 to 10 pounds per acre. For air applications, use 5 to 10 gallons of water per acre. For ground applications use 20 to 40 gallons per acre.

5753

Wilbur-Ellis

P. O. Box 16458, Fresno, CA 93755

ACTIVE INGREDIENT

Sulfur

97.0%

INERT INGREDIENTS

3.0%

100.0%

EPA Reg. No. 2935-92 AA

EPA Est. No. 2935 CA 1

KEEP OUT OF REACH OF CHILDREN!

CAUTION:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust, vapor, or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Induce vomiting by sticking finger down throat. Never give anything by mouth to an unconscious person. Get medical attention.

If In Eyes: Wash eyes with plenty of clear water for at least 15 minutes. Get medical attention.

Harmful If Swallowed: Avoid breathing dust. May cause irritation of nose, throat, or skin. Avoid contact with eyes, skin and clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas.

PHYSICAL HAZARD

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks, or flame. Do not smoke while applying this product.

RECOMMENDATIONS FOR USE (Continued)

CITRUS: SIX SPOTTED and CITRUS MITES — Apply 4 lbs per 100 gallons. CITRUS FLAT and YUMA SPIDER MITES — Apply 4 lbs per 100 gallons (At least 60 lbs per acre).

GRAPES: POWDERY MILDEW — Apply 2 lbs per 100 gallons plus a wetting agent. Apply when shoots are 6 inches long, repeat before blossoms open and continue as necessary. Concord and other labrusca type grapes may be injured by sulfur.

PEACHES & NECTARINES: BROWN ROT, COHNEUM BLIGHT, RUST, PEACH SILVER MITE, POWDERY MILDEW — Apply 5-6 lbs per 100 gallons from popcorn stage up to harvest. For BROWN ROT control in summer sprays apply up to 10 lbs per 100 gallons starting about 4-5 weeks before harvest and continuing at weekly intervals.

PEARS: SCAB, POWDERY MILDEW — Apply 4 lbs plus 1 gallon lime sulfur per 100 gallons in the pink or pre-blossom spray. On Bartlett pears, apply 4-6 lbs at petal fall through second cover sprays. Sensitive varieties (e.g., Anjou, Comice) under certain climatic conditions may be injured by sulfur.

PLUMS & PRUNES: BROWN ROT, PRUNE RUST — Apply 5-6 lbs per 100 gallons in summer sprays. Make first application about 5 weeks before harvest and thereafter at weekly intervals if wet weather occurs.

STRAWBERRIES: POWDERY MILDEW — Apply 2.5 lbs per 100 gallons. Begin at first sign of disease and repeat as necessary. Discontinue applications during hot weather. Sulfur may injure strawberries under certain climatic conditions.

SUGAR BEETS: POWDERY MILDEW — Apply 8 to 10 lbs per acre in 40 to 50 gallons of water. Apply when disease first appears. Repeat applications at 3 to 4 week intervals or as necessary.

TOMATOES: TOMATO RUSSET MITE — Apply 6 to 10 lbs per acre in sufficient water for thorough coverage. May be applied with ground equipment or aircraft, depending on local conditions.

ORNAMENTALS

ROSES: BLACK SPOT, POWDERY MILDEW, RUST — Apply 3 tablespoons per gallon (5 lbs per 100 gallons) and spray for full coverage wetting both sides of leaves. Begin applications when new growth appears in the spring and repeat every 7 days, if necessary.

In Case of Emergency, Call CHEMTREC: (800) 424-9300

NOTICE: The statements made on this label are believed to be true and accurate, but because of variations of use which are beyond our control, WILBUR ELLIS COMPANY does not make, nor does it authorize any agent or representative to make any warranty, guarantee or representation as to the performance of this material, or as to the results to be obtained therefrom, or as to the safety of its use. WILBUR ELLIS COMPANY and its agents shall be responsible to the user for any personal injury or property damage or loss resulting from the use of this product. The user or the other person using handling, storage or use of this material shall be held liable for any and all such damage or loss. WILBUR ELLIS COMPANY and its agents shall not be held liable for any such damage or loss resulting from the use of this material.

ACCEPTED

JAN 6 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the use registered under EPA Reg. No. 2935-92

BEST AVAILABLE COPY

