

USE PRECAUTIONS

HAZARDS TO HUMANS — DANGER: Poisonous If Swallowed, Inhaled or Absorbed Through Skin. Rapidly Absorbed Through Skin. Repeated Inhalation or Skin Contact May, Without Symptoms, Progressively Increase Susceptibility to Methyl Parathion Poisoning.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors or spray mist. Wear clean, natural rubber gloves, clean, waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into eyes, immediately flush with water for at least 10 minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water, wash clothing before reuse.

ANTIDOTE: ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION POISONING. Consult your physician about obtaining an adequate supply of 1-100 grain atropine tablets for emergency use. 2 PAM is also an antidote for Methyl Parathion and may be administered in conjunction with Atropine. **CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

ENVIRONMENTAL HAZARDS — This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your agricultural extension service.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT — Treated areas should not be re-entered until drifting pesticide and volatile residues have dissipated.

STORAGE — NOT FOR USE OR STORAGE IN OR AROUND THE HOME. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DISPOSAL — Do not reuse empty drum. Return to distributor for proper disposal. Do not use solution and perforate and bury with wastes away from crops and water. Do not dump or run off. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of caustic soda and rinse with water.

EPA Reg No 2935-142

EPA Reg No 2935-142

RESTRICTED USE PESTICIDE:

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

red-top METHYL PARATHION 4 SPRAY

ACTIVE INGREDIENTS:

O, O-Dimethyl O-p-nitrophenyl thiophosphate*
Xylene Range Aromatic Solvent

INERT INGREDIENTS

By Weight
45%
48%
7%
100%

1 gallon contains 4 lbs. of *Methyl Parathion.

KEEP OUT OF REACH OF CHILDREN

DANGER  POISON

FIRST AID TREATMENT

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 table-spoon of salt to a glass of water). Never give anything by mouth to an unconscious person. Have victim lie down and keep quiet.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the Methyl Parathion, remove all contaminated clothing and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING
See Side Panels for Antidote and Additional Precautions

WILBUR-ELLIS COMPANY
FRESNO • LOS ANGELES • GLENDALE, CALIF. • PORTLAND • SEATTLE

DIRECTIONS FOR USE: Apply to plant surfaces with minimum of 5 gallons.

RECOMMENDATIONS FOR

ALFALFA: Use 1 pint per acre (up to 3rd instar), LEAFHOPPERS and SPINACH. Do not apply within 15 days of harvest.

ARTICHOKES, CARROTS, CABBAGE, CAULIFLOWERS, CUCUMBERS, ETC. Use 1 pint per acre to control APHIDS. Do not apply within 15 days of harvest.

BEANS, CABBAGE, CAULIFLOWERS, CUCUMBERS, ETC. Use 1 pint per acre to control APHIDS. Do not apply within 15 days of harvest.

BEETS: Use 1 pint per acre to control LEAFHOPPERS and SPINACH. Do not apply within 15 days of harvest.

COTTON: Use 3 to 6 pints per acre to control LEAFHOPPERS, THRIPS, TOBACCO BUDWORM, and other pests. Do not apply within 15 days of harvest.

LETTUCE AND SPINACH: Use 1 pint per acre to control CABBAGE LOOPER, and other pests. Do not apply within 15 days of harvest.

POTATOES: Use 1 1/2 pints per acre to control BEETLES, LEAFHOPPERS, and other pests. Do not apply within 15 days of harvest.

SMALL GRAINS (Barley, Oats, etc.): Use 1 pint per acre to control ARMYWORMS and other pests. Do not apply within 15 days of harvest.

NOTICE: The statements made on this label are based on our control, and are not intended to make any warranty or representation, except in connection with the use thereof, except in connection with the use thereof, except in connection with the use thereof, except in connection with the use thereof.

BES

ICIDE:
aters or persons under
covered by the Certified

p
4 SPRAY
By Weight
45%
48%
7%
100%

Parathion.

HILDREN
ISON

omiting immediately. This may
ing warm salt water (1 table-
mouth to an unconscious person.

mediately remove patient from
ed clothing and wash skin
eyes, wash immediately with

SUSPECTED POISONING
ional Precautions

PANY **W.D.**
ORTLAND - SEATTLE

DIRECTIONS FOR USE (Continued):

DIRECTIONS FOR USE: Apply specified rate per acre in sufficient water to thoroughly cover plant surfaces with ground driven equipment or aircraft. When applied by air use a minimum of 5 gallons of water per acre by ground a minimum of 10 gallons per acre

RECOMMENDATIONS FOR USE:

ALFALFA: Use 1 pint per acre to control SPOTTED ALFALFA APHIDS, APHIDS, ARMYWORMS (up to 3rd instar), LEAFHOPPERS and SPIDER MITES. Do not apply within 20 days of harvest or pasturing. In California, do not apply more than 1/2 pint per acre

ARTICHOKES, CARROTS, CELERY and PEPPERS: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest. Do not use treated carrot tops for food or feed

BEANS, CABBAGE, CAULIFLOWER, BROCCOLI and BRUSSELS SPROUTS: Use 1 1/2 to 3 pints per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES and SPIDER MITES. Do not apply within 21 days of harvest. Do not apply within 15 days of harvest on dry beans.

BEETS: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest (21 days if tops are to be used for food)

COTTON: Use 3 to 6 pints per acre to control APHIDS, BOLL WEEVILS, COTTON LEAFWORMS, LEAFHOPPERS, THRIPS, COTTON LEAF PERFORATORS, ARMYWORMS (up to 3rd instar), TOBACCO BUDWORMS, BOLLWORMS, SPIDER MITES, and PINK BOLLWORMS. Do not apply within 15 days of hand picking. Workers entering cotton fields within 24 hours after treatment should wear protective clothing


LETTUCE AND SPINACH: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), CABBAGE LOOPER, IMPORTED CABBAGE WORMS, FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 21 days of harvest

POTATOES: Use 1 1/2 pints per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 5 days of harvest

SMALL GRAINS (Barley, Oats, Rye and Wheat): Use 1 pint per acre to control APHIDS (Green Bug), ARMYWORMS (up to 3rd instar) and LEAFHOPPERS. Do not apply after the heads start to form

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

BEST DOCUMENT AVAILABLE



METHYL PARATHION MIXTURE, LIQUID