

Reg # 2935-142
PM 12
173014 - 1/5-
17/03
300

JUL 11 1986

Ms. Terry Miller
Wilbur-Ellis Company
Agricultural Services Corporate Office
191 West Shaw Avenue, Suite 107
Fresno, CA 93704-2876

Dear Ms. Miller:

Subject: Revision of Label Ingredient Statement
Methyl Parathion 4 Spray
EPA Registration No. 2935-142
Your Application Dated April 11, 1986

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Xylene Range Aromatic Solvent does not need to be listed as an inert ingredient in your label ingredient statement.

Sincerely yours,

Lawrence J. Schnaubelt ^{LJS}
Acting Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

92963:Schnaubelt:173014:E-1 to KENCO:6/5/86:6/11/86:TAR:LF

SYMBOL ▶								
SURNAME ▶								
DATE ▶								

P. 2/5

USE PRECAUTIONS

HAZARDS TO HUMANS — DANGER: Poisonous If Swallowed, Inhaled or Absorbed Through Skin. Rapidly Absorbed Through Skin. Repeated Inhalation or Skin Contact May, Without Symptoms, Progressively Increase Susceptibility to Methyl Parathion Poisoning.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors or spray mist. Wear clean, natural rubber gloves; clean, waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into eyes, immediately flush with water for at least 10 minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before reuse.

FIRST AID TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the Methyl Parathion, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING

ANTIDOTE: ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION POISONING.

Consult your physician about obtaining an adequate supply of 1/100 grain atropine tablets for emergency use. "2-PAM" is also an antidote for Methyl Parathion and may be administered in conjunction with Atropine. **CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

ENVIRONMENTAL HAZARDS — This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your agricultural extension service.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep off of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MP4-3

RESTRICTED USE

For retail sale to and use only by Certified their direct supervision and only for those Applicator's certification.

Wilbur-Ell

METHYL PARATHION

ACTIVE INGREDIENTS:

O, O-Dimethyl O-p-nitrophenyl thiophosphate*.....

INERT INGREDIENTS

Xylene Range Aromatic Solvent ... 48%

1 gallon contains 4 lbs. of *

EPA Reg. No. 2935-142-ZA

KEEP OUT OF REACH

**DANGER
PELIGRO**



PRECAUTION AL USUARIO: Si usted lee hasta que le etiqueta haya sido expli

See Side Panels for Antidote

Wilbur-E

P. O. Box 16458, Fresno

RED

BEST AVAILABLE COPY

1977
APR 27 1977

USE PESTICIDE:

Certified Applicators or persons under or those uses covered by the Certified

ur-Ellis

THION 4 SPRAY

By Weight
45%
55%
100%
lbs. of *Methyl Parathion.
EPA Est. 33691 AZ-1

EACH OF CHILDREN



POISON

ed lee ingles, use este producto
sido explicudo ampliamente.

Antidote and Precautions

ur-Ellis

58, Fresno, CA 93755

DIRECTIONS FOR USE (Continued):

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas for 48 hours unless appropriate protective clothing is worn. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should specify protective clothing (see left), areas or fields that cannot be entered without this clothing, period of time area is to be vacated, and statement of practical treatment (see left) in case of exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "Danger. Area treated with Methyl Parathion on (date of application). Do not enter without appropriate protective clothing for 48 hours. [Insert here actions to take in case of accidental exposure - see left]."

STORAGE AND DISPOSAL

1. Prohibitions — Do not contaminate water, food or feed by storage or disposal.
2. Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.
3. Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
4. Storage: Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing, or pouring. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

Apply specified r
driven equi
per acre - 1
of 5 gallon

RECOMMENDATIONS

ALFALFA: Use 1
(up to 3rd in
or pasturing

ARTICHOKES, CA
WORMS (up
within 15 d

BEANS, CABBAGE
acre to cont
Do not appl
dry beans.

BEETS: Use 1 pin
LEAFHOPPERS
are to be us

COTTON: Use 3 t
LEAFHOPPERS
TOBACCO BU
within 15 d
treatment sh

LETTUCE AND SPIN
CABBAGE LO
MITES. Do n

POTATOES: Use 1
BEETLES, LEA

SMALL GRAINS (B
Bug), ARMY
start to form

NOTICE: The
tions of use w
authorize any
pilled, concern
Neither WILB
personal injury
storage or use o
resulting from

PA
88

415

In Case of Emergency, Call CI

DIRECTIONS FOR USE (Continued)

Apply specified rate per acre in sufficient water to thoroughly cover plant surfaces with ground driven equipment or aircraft. When applied by air use a minimum of 3 gallons of water per acre - by ground a minimum of 10 gallons per acre. [In California use a minimum of 5 gallons of water per acre by air.]

RECOMMENDATIONS FOR USE:

ALFALFA: Use 1 pint per acre to control SPOTTED ALFALFA APHIDS, APHIDS, ARMYWORMS (up to 3rd instar), LEAFHOPPERS and SPIDER MITES. Do not apply within 20 days of harvest or pasturing. In California, do not apply more than 1/4 pint per acre.

ARTICHOKES, CARROTS, CELERY and PEPPERS: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest. Do not use treated carrot tops for food or feed.

BEANS, CABBAGE, CAULIFLOWER, BROCCOLI and BRUSSELS SPROUTS: Use 1 1/2 to 3 pints per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES and SPIDER MITES. Do not apply within 21 days of harvest. Do not apply within 15 days of harvest on dry beans.

BEETS: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 15 days of harvest (21 days if tops are to be used for food).

COTTON: Use 3 to 6 pints per acre to control APHIDS, BOLL WEEVILS, COTTON LEAFWORMS, LEAFHOPPERS, THRIPS, COTTON LEAF PERFORATORS, ARMYWORMS (up to 3rd instar), TOBACCO BUDWORMS, BOLLWORMS, SPIDER MITES, and PINK BOLLWORMS. Do not apply within 15 days of hand picking. Workers entering cotton fields within 24 hours after treatment should wear protective clothing.

LETTUCE AND SPINACH: Use 1 pint per acre to control APHIDS, ARMYWORMS (up to 3rd instar), CABBAGE LOOPER, IMPORTED CABBAGE WORMS, FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 21 days of harvest.

POTATOES: Use 1 1/2 pints per acre to control APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES, LEAFHOPPERS and SPIDER MITES. Do not apply within 5 days of harvest.

SMALL GRAINS (Barley, Oats, Rye and Wheat): Use 1 pint per acre to control APHIDS (Green Bug), ARMYWORMS (up to 3rd instar) and LEAFHOPPERS. Do not apply after the heads start to form.

F-1185

NOTICE: The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guarantee or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on this label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

h drift expose workers
ected persons.
ishing is worn. Because
ious crops treated with
her information.
be in a treated area
ngs should specify pro-
l without this clothing,
al treatment (see left)
be given in a language
n if there is reason to
nnings must include the
ion on (date of appli-
48 hours. [Insert here

ge or disposal.
er disposal of excess
If these wastes can-
your State Pesticide
nitive at the nearest

ycling or recondition-
procedures approved

ore in closed storage
Decontaminate areas
ali and detergent and



METHYL PARATHION

NA 2

BEST AVAILABLE COPY

to EPA

JUL

Under the
Pungicide,
as annu-
registered
29

1.5/5

BEST AVAILABLE COPY

In Case of Emergency, Call CHEMTREC 800/424-9300

INSTRUCTIONS FOR USE (Continued)

Water to thoroughly cover plant surfaces with ground
m applied by air use a minimum of 3 gallons of water
of 10 gallons per acre. (In California use a minimum
y air.)

not SPOTTED ALFALFA APHIDS, APHIDS, ARMYWORMS
and SPIDER MITES. Do not apply within 20 days of harvest
t apply more than 3/4 pint per acre.

LEAFHOPPERS: Use 1 pint per acre to control APHIDS, ARMY-
WORMS, LEAFHOPPERS and SPIDER MITES. Do not apply
t use treated carrot tops for food or feed.

COCOA and BRUSSELS SPROUTS: Use 1 1/2 to 3 pints per
acre (up to 3rd instar), FLEABEETLES and SPIDER MITES.
harvest. Do not apply within 15 days of harvest on

APHIDS, ARMYWORMS (up to 3rd instar), FLEABEETLES,
Do not apply within 15 days of harvest (21 days if tops

control APHIDS, BOLL WEEVILS, COTTON LEAFWORMS,
LEAF PERFORATORS, ARMYWORMS (up to 3rd instar),
MS, SPIDER MITES, and PINK BOLLWORMS. Do not apply
Workers entering cotton fields within 24 hours after
clothing.

acre to control APHIDS, ARMYWORMS (up to 3rd instar),
BOLL WORMS, FLEABEETLES, LEAFHOPPERS and SPIDER
ys of harvest.

control APHIDS, ARMYWORMS (up to 3rd instar), FLEA-
MITES. Do not apply within 5 days of harvest.

(Wheat): Use 1 pint per acre to control APHIDS (Green
ear) and LEAFHOPPERS. Do not apply after the heads

F-1185

are believed to be true and accurate, but because of condi-
WILBUR-ELLIS COMPANY does not make, nor does it
any warranty, guaranty or representation, express or im-
plied, except in conformity with the statements on the label.
The seller shall be held responsible in any manner for any
resulting to the buyer or to the other person from handling,
once with directions, the buyer assumes all risk and liability
for use and accepts and uses this material on these conditions.



6

METHYL PARATHION MIXTURE, LIQUID

NA 2783

REGISTERED
with CHEMTREC
in EPA Letter Dated:

JUL 11 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2935-142