

DIRECTIONS FOR USE: Apply specified rate per acre in sufficient water to thoroughly cover plant surfaces with ground driven equipment or air craft. When applied by air use a minimum of 5 gallons of water per acre - by ground a minimum of 10 gallons per acre.

RECOMMENDATIONS FOR USE:

ALFALFA & CLOVER: Use 1 pt. per acre to control APHIDS, GRASSHOPPERS, LEAFHOPPERS, CUTWORMS (Climbing), LYGUS BUGS, ALFALFA WEEVIL LARVAE and SPIDER MITES. Do not apply within 1 day of harvest.

APPLES, PEACHES, PEARS & PLUMS: Use 1/2 pt. per 100 gallons of water to control APHIDS, GRASSHOPPERS, LYGUS BUGS, RED-BANDED LEAF ROLLER and SPIDER MITES. Do not apply within 1 day of harvest. Do not use more than 3.2 lbs. actual Phosdrin per acre on Apples and Pears or more than 2.56 lbs. actual per acre on Peaches and Plums.

BEANS, BEETS (Including Beet Tops), CARROTS, CELERY, EGGPLANTS, LETTUCE, ONIONS (Including Green Onions), PEAS (Including Forage), PEPPERS & SPINACH. Use 1 pt. per acre to control APHIDS, GRASSHOPPERS, LEAFHOPPERS, MEXICAN BEAN BEETLES, CABBAGE LOOPERS, CUTWORMS (Climbing), DIPTEROUS LEAFMINER (Adult), FALSE CHINCH BUGS, IMPORTED CABBAGE WORM, SALT MARSH CATERPILLAR, LYGUS BUGS, CORN EARWORM, THRIPS and SPIDER MITES. Do not apply within 1 day of harvest on Beans, Carrots, Eggplants, Onions, Peas or Peppers; within 2 days on Lettuce, within 3 days on Beets or Celery and within 4 days on Spinach.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, KALE, MUSTARD GREENS & TURNIPS (Including Turnip Tops.): Use 1 pt per acre to control APHIDS, CABBAGE LOOPER, IMPORTED CABBAGE WORM, GRASSHOPPERS, LEAFHOPPERS, SALT-MARSH CATERPILLAR, CUTWORMS (Climbing), DIPTEROUS LEAFMINER (Adults), LYGUS BUGS, FALSE CHINCH BUGS and SPIDER MITES. Do not apply within 1 day of harvest on Broccoli or Cabbage, and within 3 days on Brussels Sprouts, Cauliflower, Collards, Kale, Mustard Greens or Turnips.

CORN (Field, Sweet, Pop, Forage & Grain): Use 1 pt. per acre to control APHIDS. Do not apply within 1 day of harvest.

GRAPES: Use 2 pts. per acre to control APHIDS, LEAF FOLDER, LEAFHOPPERS, RED-BANDED LEAF ROLLER, LYGUS BUGS and SPIDER MITES. Do not apply within 5 days of picking.

LETTUCE: Use 1 pint per acre to control CABBAGE LOOPER and CORN EARWORM. Do not use within 4 days of harvest. Do not use more than 1 pint per acre.

PEPPERS: For control of APHIDS, apply 1/4 to 1/2 pint per acre. For control of GRASSHOPPERS, LEAFHOPPERS, and MITES, use 1/2 to 1 pint per acre. Do not treat within 2 days of harvest. For control of ORGANOPHOSPHATE-RESISTANT GREEN PEACH APHIDS (California only), use 1 to 2 pints per acre. Do not treat within 4 days of harvest. If this crop is grown for processing, do not use the higher rates recommended for control of Organophosphate-Resistant Green Peach Aphids without consulting your processor field representative.

POTATOES: For control of APHIDS, GRASSHOPPERS, LEAFHOPPERS, and SPIDER MITES, use 1/2 pint per acre. Do not apply within 1 day of harvest. For control of GREEN PEACH APHID (California only), use 1 quart per acre. Do not apply less than 30 gallons of water per acre for ground application and 5 gallons of water for aerial application. Do not allow workers to re-enter fields within two days of application. Do not make applications within four days of harvest. Do not apply more than 1 quart per acre.

TOMATOES: For control of APHIDS, GRASSHOPPERS, LEAFHOPPERS, and SPIDER MITES, use 1 pint per acre. Do not apply within 1 day of harvest.

red-top PHOSDRIN® 4 SPRAY

ACTIVE INGREDIENTS:

Alpha Isomer of 2-Carbomethoxy-1-methylvinyl Dimethyl Phosphate 30.0%*

Related Compounds 20.0%*

INERT INGREDIENTS 50.0%

*Equivalent to 50.0% W. Phosdrin Insecticide. 100.0%

EPA Reg. No. 2935-167-AA



POISON

DANGER: KEEP OUT OF REACH OF CHILDREN!

Poisonous If Swallowed, Inhaled or Absorbed Through Skin. Rapidly Absorbed Through Skin or Eyes. Repeated Inhalation or Skin Contact May, Without Symptoms, Progressively Increase Susceptibility to Phosdrin Insecticide Poisoning.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors. Wear heavy natural rubber gloves and waterproof protective clothing and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Phosdrin insecticide protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into eyes, immediately flush with water for at least ten minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before re-use. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with strong lye solution, then destroy containers to prevent re-use.

"This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label."

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. (See additional caution and antidotes on side panel.)

WILBUR-ELLIS COMPANY
FRESNO • LOS ANGELES • GLENDALE, ARIZ. • PORTLAND • SEATTLE

RECOMMENDATIONS

RASPBERRIES: Use 1 day of harvest

SORGHUMS (Grain): not apply with

STRAWBERRIES: Use ROLLER, SALT-1 day of harvest

SQUASH (Summer melons): Use 1

LOOPER, DIPTEROUS HOPPERS. Do not

ANTIDOTE AND FIRST AID

Insecticide Poisoning: Give grain atropine

Suspected Poisoning: Internal. If the

done by introduction of salt to a glass of water

By Mouth To a Glass of Water. External. If the

the vicinity of the body and running water for at least 15 minutes

Start Artificial Respiration. Immediately Take to Hospital

PRECAUTIONS IN USE

humans or bees. Do not enter treated fields or pastures with animals. Do not

To prevent re-use, wash equipment by washing the material on these conditions

by washing the material on these conditions

by washing the material on these conditions

by washing the material on these conditions

by washing the material on these conditions

by washing the material on these conditions

by washing the material on these conditions

by washing the material on these conditions

by washing the material on these conditions

by washing the material on these conditions

by washing the material on these conditions

by washing the material on these conditions

RECOMMENDATIONS FOR USE:

RASPBERRIES: Use 1 pt. per acre to control APHIDS and LEAFHOPPERS not apply within 3 days of harvest.

SORGHUMS (Grain & Forage): Use 1 pt. per acre to control APHIDS and CORN EARWORMS. Do not apply within 3 days of harvest.

STRAWBERRIES: Use 2 pts. per acre to control APHIDS, GRASSHOPPERS, STRAWBERRY LEAF ROLLER, SALT-MARSH CATERPILLAR, LYGUS BUGS and SPIDER MITES. Do not apply within 1 day of harvest.

SQUASH (Summer) & MELONS (Including Cantaloupes, Honey-Dew, Muskmelons and Watermelons): Use 1/2 pt. per acre to control APHIDS. Use 1 pt. per acre to control CABBAGE LOOPER, DIPTEROUS LEAFMINER (Adults), LEAFHOPPERS, LYGUS BUGS, MITES and GRASSHOPPERS. Do not apply within 1 day of harvest.

ANTIDOTE AND FIRST AID TREATMENT: Atrophine Is The Emergency Antidote For Phosdrin Insecticide Poisoning. Consult your physician about obtaining an adequate supply 1/100 grain atropine tablets for emergency use. Call Physician Immediately In All Cases of Suspected Poisoning.

Internal. If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). Have victim lie down and keep quite. **Never Give Anything By Mouth To An Unconscious Person.**

External. If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

Start Artificial Respiration Immediately If Patient Is Not Breathing. Transport The Patient Immediately To The Nearest Physician.


PRECAUTIONS IN USING: Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Unprotected humans and animals should be kept out of treated fields on the day of treatment. Keep out of the reach of children and domestic animals. Do not contaminate food or feed products. Keep away from heat and open flame. To prevent re-use, empty container is to be decontaminated and destroyed. Decontaminate by washing thoroughly with lye solution, then rinsing with water.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

NOTICE: The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

F-675 EPA Est. 2935-CA-1



**ORGANIC PHOSPHATE COMPOUND MIXTURE,
LIQUID, NOIBN**

DIRECTIONS FOR USE: Apply specified rate per acre in sufficient water to thoroughly cover plant surfaces with ground driven equipment or air craft. When applied by air use a minimum of 5 gallons of water per acre - by ground a minimum of 10 gallons per acre.

RECOMMENDATIONS FOR USE:

ALFALFA & CLOVER: Use 1 pt. per acre to control APHIDS, GRASSHOPPERS, LEAFHOPPERS, CUT-WORMS (Climbing), LYGUS BUGS, ALFALFA WEEVIL LARVAE and SPIDER MITES. Do not apply within 1 day of harvest.

APPLES, PEACHES, PEARS & PLUMS: Use 1/2 pt. per 100 gallons of water to control APHIDS, GRASSHOPPERS, LYGUS BUGS, RED-BANDED LEAF ROLLER and SPIDER MITES. Do not apply within 1 day of harvest. Do not use more than 3.2 lbs. actual Phosdrin per acre on Apples and Pears or more than 2.56 lbs. actual per acre on Peaches and Plums.

BEANS, BEETS (Including Beet Tops), CARROTS, CELERY, EGGPLANTS, LETTUCE, ONIONS (Including Green Onions), PEAS (Including Forage), PEPPERS & SPINACH. Use 1 pt. per acre to control APHIDS, GRASSHOPPERS, LEAFHOPPERS, MEXICAN BEAN BEETLES, CABBAGE LOOPERS, CUTWORMS (Climbing), DIPTEROUS LEAFMINER (Adult), FALSE CHINCH BUGS, IMPORTED CABBAGE WORM, SALT MARSH CATERPILLAR, LYGUS BUGS, CORN EARWORM, THRIPS and SPIDER MITES. Do not apply within 1 day of harvest on Beans, Carrots, Eggplants, Onions, Peas or Peppers; within 2 days on Lettuce, within 3 days on Beets or Celery and within 4 days on Spinach.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, KALE, MUSTARD GREENS & TURNIPS (Including Turnip Tops.): Use 1 pt per acre to control APHIDS, CABBAGE LOOPER, IMPORTED CABBAGE WORM, GRASSHOPPERS, LEAFHOPPERS, SALT-MARSH CATERPILLAR, CUTWORMS (Climbing), DIPTEROUS LEAFMINER (Adults), LYGUS BUGS, FALSE CHINCH BUGS and SPIDER MITES. Do not apply within 1 day of harvest on Broccoli or Cabbage, and within 3 days on Brussels Sprouts, Cauliflower, Collards, Kale, Mustard Greens or Turnips.

CORN (Field, Sweet, Pop, Forage & Grain): Use 1 pt. per acre to control APHIDS. Do not apply within 1 day of harvest.

GRAPES: Use 2 pts. per acre to control APHIDS, LEAF FOLDER, LEAFHOPPERS, RED-BANDED LEAF ROLLER, LYGUS BUGS and SPIDER MITES. Do not apply within 5 days of picking.

LETTUCE: Use 1 pint per acre to control CABBAGE LOOPER and CORN EARWORM. Do not use within 4 days of harvest. Do not use more than 1 pint per acre.

PEPPERS: For control of APHIDS, apply 1/4 to 1/2 pint per acre. For control of GRASSHOPPERS, LEAFHOPPERS, and MITES, use 1/2 to 1 pint per acre. Do not treat within 2 days of harvest. For control of ORGANOPHOSPHATE-RESISTANT GREEN PEACH APHIDS (California only), use 1 to 2 pints per acre. Do not treat within 4 days of harvest. If this crop is grown for processing, do not use the higher rates recommended for control of Organophosphate-Resistant Green Peach Aphids without consulting your processor field representative.

POTATOES: For control of APHIDS, GRASSHOPPERS, LEAFHOPPERS, and SPIDER MITES, use 1/2 pint per acre. Do not apply within 1 day of harvest. For control of GREEN PEACH APHID (California only), use 1 quart per acre. Do not apply less than 30 gallons of water per acre for ground application and 5 gallons of water for aerial application. Do not allow workers to re-enter fields within two days of application. Do not make applications within four days of harvest. Do not apply more than 1 quart per acre.

TOMATOES: For control of APHIDS, GRASSHOPPERS, LEAFHOPPERS, and SPIDER MITES, use 1 pint per acre. Do not apply within 1 day of harvest.

red-top PHOSDRIN® 4 SPRAY

ACTIVE INGREDIENTS:

Alpha Isomer of 2-Carbomethoxy-1-methylvinyl Dimethyl Phosphate 30.0%*

Related Compounds 20.0%*

INERT INGREDIENTS 50.0%

*Equivalent to 50.0% W. Phosdrin Insecticide. 100.0%

EPA Reg. No. 2935-167-AA



DANGER: KEEP OUT OF REACH OF CHILDREN!

Poisonous If Swallowed, Inhaled or Absorbed Through Skin. Rapidly Absorbed Through Skin or Eyes. Repeated Inhalation or Skin Contact May, Without Symptoms, Progressively Increase Susceptibility to Phosdrin Insecticide Poisoning.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe vapors. Wear heavy natural rubber gloves and waterproof protective clothing and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Phosdrin insecticide protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into eyes, immediately flush with water for at least ten minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before re-use. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with strong lye solution, then destroy containers to prevent re-use.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

(See additional caution and antidotes on side panel.)

WILBUR-ELLIS COMPANY
 FRESNO • LOS ANGELES • TUCUMAN, ARIZ. • PORTLAND • SEATTLE

RECOMMENDATIONS FOR USE:

RASPBERRIES: Use 1 pt. per acre to control APHIDS and LEAFHOPPERS. Do not apply within 3 days of harvest.

SORGHUMS (Grain & Forage): Use 1 pt. per acre to control APHIDS and CORN EARWORMS. Do not apply within 3 days of harvest.

STRAWBERRIES: Use 2 pts. per acre to control APHIDS, GRASSHOPPERS, STRAWBERRY LEAF ROLLER, SALT-MARSH CATERPILLAR, LYGUS BUGS and SPIDER MITES. Do not apply within 1 day of harvest.

SQUASH (Summer) & MELONS (Including Cantaloupes, Honey-Dew, Muskmelons and Watermelons): Use 1/2 pt. per acre to control APHIDS. Use 1 pt. per acre to control CABBAGE LOOPER, DIPTEROUS LEAFMINER (Adults), LEAFHOPPERS, LYGUS BUGS, MITES and GRASSHOPPERS. Do not apply within 1 day of harvest.

ANTIDOTE AND FIRST AID TREATMENT: Atrophine Is The Emergency Antidote For Phosdrin Insecticide Poisoning. Consult your physician about obtaining an adequate supply 1/100 grain atropine tablets for emergency use. Call Physician Immediately In All Cases of Suspected Poisoning.

Internal. If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). Have victim lie down and keep quite. **Never Give Anything By Mouth To An Unconscious Person.**

External. If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

Start Artificial Respiration Immediately If Patient Is Not Breathing. Transport The Patient Immediately To The Nearest Physician.


PRECAUTIONS IN USING: Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Unprotected humans and animals should be kept out of treated fields on the day of treatment. Keep out of the reach of children and domestic animals. Do not contaminate food or feed products. Keep away from heat and open flame. To prevent re-use, empty container is to be decontaminated and destroyed. Decontaminate by washing thoroughly with lye solution, then rinsing with water.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

NOTICE: The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

F-675 EPA Est. 2935-CA 1



**ORGANIC PHOSPHATE COMPOUND MIXTURE,
LIQUID, NOIBN**