

SEVIN® 5 DUST

ACCEPTED
APR 8 - 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 2935-193 SUBJECT
TO ATTACHED COMMENTS.

RECOMMENDATIONS FOR USE:

ALFALFA and CLOVERS: Use 30 lbs. per acre to control THRIPS, LEAFHOPPERS, THREE-CORNERED ALFALFA HOPPER, JAPANESE BEETLE, ALFALFA CATERPILLAR, ARMYWORMS, CORN EARWORMS and STINK BUGS.

BEANS: Use 40 lbs. per acre to control MEXICAN BEAN BEETLE, LEAFHOPPERS, BEAN LEAF BEETLE, CUCUMBER BEETLES, WESTERN BEAN CUTWORMS, FLEA BEETLES, CORN EARWORM, ARMYWORMS, VELVET BEAN CATERPILLAR, LYGUS and STINK BUGS.

CORN: Use 40 lbs. per acre to control CORN EARWORM, EUROPEAN CORN BORER, FALL ARMYWORMS, SAP BEETLES and JAPANESE BEETLES. Do not apply within 7 days of harvest for forage.

COTTON: Use 40 to 50 lbs. per acre to control THRIPS, FLEA BEETLE, FLEAHOPPERS, LEAFWORM, BOLLWORM, FALL ARMYWORM, LEAF PERFORATOR, LEAF ROLLERS, LEAFHOPPERS, LYGUS BUGS, SALT MARSH CATERPILLAR, STINK BUGS and GRASSHOPPERS.

CUCUMBERS and SUMMER SQUASH: For control of FLEA BEETLES, CUCUMBER BEETLES, SQUASH BUGS, MELONWORM and PICKLEWORM apply 20 lbs. per acre.

GRAPES: Use 40 to 50 lbs. per acre to control GRAPE LEAF FOLDER, GRAPE LEAFHOPPER, GRAPE BERRY MOTH and JAPANESE BEETLE. No time limitation.

EGGPLANTS, PEPPERS and TOMATOES: Use 40 lbs. per acre to control TOMATO FRUITWORM, FALL ARMYWORM, FLEA BEETLES, LEAFHOPPERS, TOMATO HORNWORM, EUROPEAN CORN BORER, STINK BUGS and LYGUS BUGS. No time limitation.

POTATOES: Use 30-40 lbs. per acre to control LEAFHOPPERS, FLEA BEETLES and COLORADO POTATO BEETLE. Apply to foliage only.

STRAWBERRIES: Use 40 lbs. per acre to control MEADOW SPITTLEBUG, and LEAF ROLLER. Do not apply within 1 day of harvest.

FILBERTS: For control of APHIDS, LEAFROLLERS and FILBERTWORM apply 60-80 lbs. per acre. For LEAFWORM apply when eggs are hatching. For FILBERTWORM apply on first appearance of adult moths and repeat 3 to 4 weeks later.

COMPOSITION:

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS 95%

USDA Reg. No. 2935-193 100%

KEEP OUT OF REACH OF CHILDREN!

CAUTION: Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing dust. Skin contact with any insecticide may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. To protect fish and wildlife, do not contaminate any body of water with this material.

Note For Physicians: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED ON THIS LABEL.

DO NOT ALLOW THIS PRODUCT TO DRIFT FROM THE AREA BEING TREATED.

● Sevin is a trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

RECOMMENDATIONS FOR USE: (Continued)

FRUIT TREES (Apples, Cherries, Peaches, Pears, Plums and Prunes): For control of APHIDS, PEAR PSYLLA, PEAR SLUG, WESTERN TUSSOCK MOTH, CODLING MOTH, CHERRY FRUIT FLY, PLUM CURCULIO, LEAFROLLERS, ORANGE TORTRIX, PEACH TWIG BORER, CAT-FACING INSECTS and PERIODICAL CICADA apply 40-60 lbs. per acre by aircraft. To avoid possible fruit foliage injury on McIntosh and York apples do not apply before second cover. Do not apply within 1 day of harvest.

F-966-193-A