

USE CAUTION

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil.

J.W. 6 1974
DESTROY THIS CONTAINER WHEN EMPTY — NEVER RE-USE

DIRECTIONS FOR USE

2935-208

1 [**BEANS: (Root and Stem Rot - Rhizoctonia solani): FURROW:** Application rates are to 14,500 linear feet of row for bush type beans and to 8,430 linear feet of row for pole beans. Apply 2-3 qts. in 8 to 10 gallons of water. Spray planting furrow and covering soil at planting time. Avoid applying directly to seed; otherwise delayed emergence may occur. IN CALIFORNIA AND ARIZONA apply 3-4 pints in 8 to 10 gals. of water. Apply in-furrow and incorporate with covering soil at planting. Use lower rate of lighter soils. Avoid application to bare seed.

2 [*BEANS*
SNAP AND DRY BEANS: (White Mold-Sclerotinia sclerotiorum): Use 1 gal. mixed in 8 to 10 gals. of water and apply to 14,500 linear feet of row for bush type beans or to 8,430 linear feet of row for pole beans. Spray in an 8" wide band, centered on the row, immediately after or at time of seeding. Avoid applying directly to seed; otherwise delayed emergence may occur. (Where dinitro type pre-emergence herbicides are used add 1 gal. to the final herbicide spray mix and apply simultaneously.) If disease is severe, application may be repeated at 2 to 3 week intervals, using 1 or 2 nozzles per row and directing spray at base of plants. Do not apply after first bloom. Soil should remain undisturbed after treatment. Do not feed treated vines to livestock.

3 [**COTTON (Damping Off-Rhizoctonia Solani):** Use 1/2 to 1 gallon per acre in at least 15 gallons of water (per 13,000 linear feet of row based on 40" spacings). Spray on the seed and surrounding soil at time of planting. Do not allow the feeding or grazing of cotton foliage by livestock.

red-top

Terrador® 2 Spray

ACTIVE INGREDIENT:		
Pentachloronitrobenzene	24.0%
INERT INGREDIENTS:	76.0%
	TOTAL	100.0%

Contains 2 lbs. of Pentachloronitrobenzene per gallon

EPA Reg. No. 2935-208

CAUTION: KEEP OUT OF REACH OF CHILDREN!

Harmful if swallowed. Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face thoroughly after using.

Do not store near heat or open flame. Use with adequate ventilation. Keep container closed. Avoid contamination of feed and foodstuffs.

CAUTION COMBUSTIBLE LIQUID

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Terrador® Trademark of Olin Corporation

W E WILBUR-ELLIS COMPANY W E
 FRESNO · LOS ANGELES · GLENDALE · PORTLAND · SEATTLE

4 [**GARLI**
to 21,
2 noz:
and fl

5 [**POTAI**
with 1
disc a
disc
formin

POTAI
linear
Spray
seed
nozzle
above
fan
shoula

DO N
LABEL

NOTIC
which
present
use the
seller
buyer
materia

red-top

Terrador[®] 2 Spray

ACTIVE INGREDIENT:

Pentachloronitrobenzene 24.0%

INERT INGREDIENTS:

..... 76.0%

TOTAL 100.0%

Contains 2 lbs. of Pentachloronitrobenzene per gallon

EPA Reg. No. 2935-208

CAUTION: KEEP OUT OF REACH OF CHILDREN!

Harmful if swallowed. Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face thoroughly after using.

Do not store near heat or open flame. Use with adequate ventilation. Keep container closed. Avoid contamination of feed and foodstuffs.

CAUTION COMBUSTIBLE LIQUID

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Terrador[®] Trademark of Olin Corporation

WE

WILBUR-ELLIS COMPANY

WE

FRESNO · LOS ANGELES · GLENDALE · PORTLAND · SEATTLE

- 4 | **GARLIC (White Rot - Sclerotium cepivorum):** Use 10 gallons mixed in 100 gallons of water and to 21,800 linear feet of row at planting time. Attach a suitable rig to the machine planter using 2 nozzles per planting furrow. Direct the front nozzle to spray the bottom of the open furrow and the "seed" as it is dropped. Direct the rear nozzle to spray the covering soil.
- 5 | **POTATOES (Rhizoctonia solani) - BROADCAST APPLICATION** - Apply 9 to 12½ gallon. mixed with 100 to 150 gallons of water to one acre. Apply to the soil surface prior to planting and disc and cross disc to a depth of 4 to 6 inches. A drag harrow is usually attached behind the discs on the last discing. It is very important to thoroughly mix the Terrador into the tuber-forming zone of the soil. It is desirable to make treatment when the soil is slightly moist.

POTATOES (Rhizoctonia solani) - ROW (In furrow) APPLICATION: Use 5 gallons per 12,400 linear feet of row mixed in sufficient water to total approximately 20 gallons of finished spray. Spray several points in the planting row to treat all of the soil that surrounds and covers the seed piece and potato forming zone. Three nozzles attached to the planter work best; one fan nozzle ahead of opening disc (approximately 5 gals. of spray per acre); one cone nozzle placed above and slightly ahead of closing discs (approximately 10 gallons of spray per acre); one fan nozzle to rear of closing discs (approximately 5 gals. of spray per acre). Spray pattern should treat 12 to 14 inch band centered on the row.

DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED ON THIS LABEL. DO NOT ALLOW THIS PRODUCT TO DRIFT FROM THE AREA BEING TREATED.

NOTICE: The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

RECEIVED
JAN 21 1974

F-1173