

RECEIVED
 1955
 JUN 21 1955
 U.S. DEPARTMENT OF AGRICULTURE
 AGRICULTURAL RESEARCH SERVICE
 ARS 111-10-10

ATTENTION - READ OUT OF REAR OF CONTAINER

Wash hands thoroughly with soap and water after use. Avoid contamination of food and feedstuffs. Do not take any part of the body or crop to be used for seed or forage. Do not use for removal of body sores or insect infestations on humans.

Keep out of lakes, streams, ponds, fish markets, and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from the treated area. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Use containers and equipment and disposal of wastes by burying in non-polluting area with concrete. Containers should be disposed of by burning with removal of burning (keep out of smokes).

Directions for Use (Continued)
 Apply 1.5 to 2.0 lbs. per acre to control TWO-SPOTTED and PACIFIC MITE. Do not apply within 7 days of harvest.

Apply 1.5 to 2.0 lbs. per acre to control EUROPEAN RED, CLOVER, TWO-SPOTTED and PACIFIC MITE. Do not repeat applications within 7 days. Do not apply within 7 days of harvest. Do not use more than 2 lbs. per acre.

Apply 1.5 to 2.0 lbs. per acre to control TWO-SPOTTED, BEAN-LEAF and PACIFIC MITE. Do not apply within 7 days of harvest.

Apply 1.5 to 2.0 lbs. per acre to control EUROPEAN RED MITE, TWO-SPOTTED and PACIFIC MITE. Apply at 1.5 lbs. per acre to control BEAN-LEAF MITE. Do not apply within 7 days of harvest. Do not use more than 2 lbs. per acre.

Apply 1.5 to 2.0 lbs. per acre to control TWO-SPOTTED and PACIFIC MITE. Do not apply within 7 days of harvest.

Apply 1.5 to 2.0 lbs. per acre to control EUROPEAN RED, CLOVER, TWO-SPOTTED and PACIFIC MITE. Do not repeat applications within 7 days. Do not apply within 7 days of harvest. Do not use more than 2 lbs. per acre.

Apply 1.5 to 2.0 lbs. per acre to control TWO-SPOTTED, BEAN-LEAF and PACIFIC MITE. Do not apply within 7 days of harvest.

Apply 1.5 to 2.0 lbs. per acre to control EUROPEAN RED MITE, TWO-SPOTTED and PACIFIC MITE. Apply at 1.5 lbs. per acre to control BEAN-LEAF MITE. Do not apply within 7 days of harvest. Do not use more than 2 lbs. per acre.

Apply 1.5 to 2.0 lbs. per acre to control TWO-SPOTTED and PACIFIC MITE. Do not apply within 7 days of harvest.

Apply 1.5 to 2.0 lbs. per acre to control EUROPEAN RED, CLOVER, TWO-SPOTTED and PACIFIC MITE. Do not repeat applications within 7 days. Do not apply within 7 days of harvest. Do not use more than 2 lbs. per acre.

Apply 1.5 to 2.0 lbs. per acre to control TWO-SPOTTED, BEAN-LEAF and PACIFIC MITE. Do not apply within 7 days of harvest.

Apply 1.5 to 2.0 lbs. per acre to control EUROPEAN RED MITE, TWO-SPOTTED and PACIFIC MITE. Apply at 1.5 lbs. per acre to control BEAN-LEAF MITE. Do not apply within 7 days of harvest. Do not use more than 2 lbs. per acre.