

**RED-TOP BRAND 2,4, 5- TP 256**

A plant hormone-type product effective in controlling the harvest drop of apples of all varieties tested including McIntosh, Delicious, Golden Delicious, Starking, Starman, Turley, Winesap, and Rome Beauty.

**DIRECTIONS FOR DROP CONTROL OF APPLES -** RED-TOP 2,4, 5-TP 256 becomes effective within 7 to 12 days after application and maintains a high degree of drop control for 5 to 7 weeks. Apply 4 days before harvest drop normally begins (7 to 20 days before harvest.) Regional growing conditions may require a different time of application.

**GROUND FIG SPRAY -** Mix 3.8 fl. oz. of RED-TOP 2,4, 5-TP 256 with 100 gallons of water and stir vigorously. This spray solution contains 20 ppm of Silvex (2, 4, 5-TP). Wet foliage thoroughly to obtain good coverage. Consult agricultural authorities for local recommendations.

**TO IMPROVE SET & YIELD OF ANJOU PEARS -** RED-TOP 2, 4, 5-TP 256 applied to Anjou Pear trees immediately after harvest time is very effective in increasing fruit set the following year. 1 fl. oz. of this product thoroughly mixed with 20 gallons of water will make a 7.5 ppm spray solution. Do not apply a stronger spray. Apply spray at bulk volume during the week after harvest.

**RED-TOP**

**2, 4, 5 - TP 256  
A HORMONE TYPE SPRAY**

For control of preharvest drop of apples

To Improve Set & Yield of Anjou Pears, for control of preharvest drop of Italian Type Prunes

**ACTIVE INGREDIENT:**

Silvex (2-(2, 4, 5-Trichlorophenoxy) propionic acid) as the triethanolamine salt\* . . . . . 11.45%

**INERT INGREDIENTS . . . . . 88.55%**

\*Equivalent to 7.37% of 2 (2, 4, 5-Trichlorophenoxy) propionic acid or 256 grams per gallon

USDA Reg. No. 2935

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
(18 point) (12 point)

**CAUTION:** Avoid contact with skin, eyes, and clothing. Wash hands after use. Harmful if swallowed. Avoid spray drift to nearby susceptible plants as this product may injure cotton, beans, peas, grapes, and ornamentals. Coarse sprays are less likely to drift. Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Avoid contamination of irrigation ditches or water used for domestic purposes.

**FOR CONTROL OF PREHARVEST DROP OF ITALIAN TYPE PRUNES**  
Spray trees until dripping wet with Red-Top 2, 4, 5-TP 256 diluted with water to 20 ppm (3.8 fl. oz. concentrate to 100 gallons water). Application should be made not less than 60 days before harvest and not earlier than 800 to 1100 degree days after full bloom when above 43 degrees F. For correct time of application, consult local agricultural advisor, since these directions are based on use in the Pacific Northwest.

**DIRECTIONS FOR USE -** Keep agitator running during filling, mixing and spraying operations. Apply as a spray wetting foliage uniformly and completely. Do not delay harvest as fruit condition and keeping qualities will be impaired if left on trees beyond normal maturity.

