

# SEVIN® 7.5 SULFUR 50 DUST

This product may be applied by ground or aircraft equipment.

## RECOMMENDATIONS FOR USE:

**ALFALFA and CLOVERS:** Use 20 lbs. per acre to control THRIPS, LEAFHOPPERS, THREE-CORNERED ALFALFA HOPPER, JAPANESE BEETLE, ALFALFA CATERPILLAR, ARMYWORMS, CORN EARWORMS and STINK BUGS.

**ASPARAGUS:** For the control of ASPARAGUS BEETLE, use 25 pounds per acre during cutting season. Do not repeat applications within 3 weeks. Do not apply within 1 day of harvest.

**BEANS:** Use 20 to 25 lbs. per acre to control MEXICAN BEAN BEETLE, LEAFHOPPERS, BEAN LEAF BEETLE, CUCUMBER BEETLES, WESTERN BEAN CUTWORMS, FLEA BEETLES, CORN EARWORM, ARMYWORMS, BEAN BEETLE, BEAN CATERPILLAR, LYGUS and STINK BUGS.

**CORN:** Use 20 to 25 lbs. per acre to control CORN EARWORM, CORN BORER, FALL ARMYWORMS, SAP BEETLES and JAPANESE BEETLES. Do not apply within 7 days of harvest for forage.

**COTTON:** Use 25 to 30 lbs. per acre to control THRIPS, FLEA BEETLE, FLEA-HOPPERS, LEAFWORM, BOLLWORM, FALL ARMYWORM, LEAF PERFORATOR, LEAF ROLLERS, LEAFHOPPERS, LYGUS BUGS, SALT MARSH CATERPILLAR, STINK BUGS and GRASSHOPPERS.

**GRAPES:** Use 25 to 30 lbs. per acre to control GRAPE LEAF FOLDER, GRAPE LEAFHOPPER, GRAPE BERRY MOTH and JAPANESE BEETLE. No time limitation.

**EGGPLANTS, PEPPERS and TOMATOES:** Use 30 to 40 lbs. per acre to control TOMATO FRUITWORM, FALL ARMYWORM, FLEA BEETLES, LEAFHOPPERS, TOMATO HORNWORM, EUROPEAN CORN BORER, STINK BUGS and LYGUS BUGS. No time limitation.

**POTATOES:** Use 20 to 25 lbs. per acre to control LEAFHOPPERS, FLEA BEETLES and COLORADO POTATO BEETLE. Apply to foliage only.

**STRAWBERRIES:** Use 20 to 25 lbs. per acre to control MEADOW SPITTLEBUG, and LEAF ROLLER. Do not apply within 1 day of harvest.

**DO NOT USE THIS PRODUCT FOR ANY PURPOSE OTHER THAN THAT RECOMMENDED ON THIS LABEL.**

**DO NOT ALLOW THIS PRODUCT TO DRIFT FROM THE AREA BEING TREATED.**

© Sevin is a trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate.

EPA Est. 2935-CA-1 F-875

## ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) .....	7.5%
Sulfur .....	50.0%
<b>INERT INGREDIENTS</b> .....	<b>42.5%</b>
	<b>100.0%</b>

EPA Reg. No. 2935-233-AA

**CAUTION:**  
 AUG 25 1975  
 Federal Insecticide, Fungicide, and Rodenticide Act  
 2935-233

**KEEP OUT OF REACH OF CHILDREN!**

**HARMFUL IF INHALED OR SWALLOWED.** Avoid Breathing of Dust. Do Not Take Internally. Avoid Contact with Skin and Eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. Do Not Use 2-PAM, opiates, or cholinesterase inhibiting drugs.

**AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES.** Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**CAUTION:** Applications of sulfur should not be made just prior to or during periods of high summer temperatures. Do not use or allow this material to drift onto sulfur sensitive Melons, Squash, Cucumbers, Apricots or other plants which may be injured by sulfur.

Carbaryl can be injurious to tender foliage. Do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

Do not reuse empty container. Containers should be disposed by burying with wastes or burning (keep out of smoke).