

2935-246

01/10/1973

1/2

BEN-SUL 60 DUST

ACTIVE INGREDIENT:

Sulfur - - - - - 60%

INERT INGREDIENTS: - - - - - 40%


Total 100%

EPA REG. NO. 2935-246
(12 point)CAUTION: KEEP OUT OF REACH OF CHILDREN!
(18 point) (14 point)


CAUTION: Some fruits and other plants are susceptible to injury from (18 pt.) sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage when the temperature is high (90° or above). Do not make sulfur applications at such times. Do not use this product for any purpose other than that recommended on this label. Never use sulfur before, with, or immediately following foliage oil sprays.

Do not reuse container. Destroy when empty.

NOTICE: The statements made on this label, are believed to be true and accurate, but because of conditions of use which are beyond our control, WILBUR-ELLIS COMPANY does not make, nor does it authorize any agent or representative to make, any warranty, guaranty or representation, express or implied, concerning this material or the use thereof, except in conformity with the statements on the label. Neither WILBUR-ELLIS COMPANY nor the seller shall be held responsible in any manner for any personal injury or property damage or loss resulting to the buyer or to the other person from handling, storage or use of this material, not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.



BEST DOCUMENT AVAILABLE



2/2 19 5

Left Panel

DIRECTIONS FOR USE: This material may be applied with ground equipment or aircraft.

Where recommended for prevention of Powdery Mildew, apply when disease first appears and repeat at 7 to 10 day intervals throughout the growing season. Shorten the interval during humid, rainy weather.

Where recommended for prevention of Brown Rot, apply at pink bud or popcorn stage during bloom and at petal fall. Also repeat at 7 to 10 day intervals late in the season as conditions warrant.

RECOMMENDATIONS FOR USE:

ALFALFA: For control of Atlantic Mite, Spider Mite, and Clover Mite use 25 lbs. per acre.


HOPS: For control of Red Spider Mites use 25 lbs. per acre. For prevention of Brown Rot on plums and prunes use 40 to 50 lbs. per acre. Apply when fruit starts to ripen.

BEANS: For the control of Pacific Mite (Red Spider) use 25 lbs. per acre. For control of Powdery Mildew use 40 to 50 lbs. per acre.

CORN: For the control of Spider Mites use 25 lbs. per acre.

GRAPES: For the control of Powdery Mildew use 8 to 20 lbs. per acre. Make first application when shoots are 6 inches to 8 inches long. Make a second application when shoots are 12 inches to 15 inches long and the third about 14 days later. Repeat applications at 10 to 14 day intervals to protect new growth. Avoid applications during hot weather as injury may occur. Concord and other labrusca type grapes may be injured by sulfur.

PEACHES: For control of Powdery Mildew and prevention of Brown Rot use 40 to 50 lbs. per acre. For prevention of Brown Rot apply when fruit starts to ripen for the control of Peach Silver Mite.



BEST DOCUMENT AVAILABLE

