



BEN-SUL 60 DUST

DIRECTIONS FOR USE: This material may be applied with ground equipment or aircraft.

Where recommended for prevention of Powdery Mildew, apply when disease first appears and repeat at 7 to 10 day intervals throughout the growing season. Shorten the interval during humid, rainy weather.

Where recommended for prevention of Brown Rot, apply at pink bud or popcorn stage during bloom and at petal fall. Also repeat at 7 to 10 day intervals late in the season as conditions warrant.

RECOMMENDATIONS FOR USE:

ALFALFA: For control of Atlantic Mite, Spider Mite, and Clover Mite use 25 lbs. per acre.

BEANS: For the control of Pacific Mite (Red Spider) use 25 lbs. per acre. For control of Powdery Mildew use 40 to 50 lbs. per acre.

CORN: For the control of Spider Mites use 25 lbs. per acre.

GRAPES: For the control of Powdery Mildew use 8 to 20 lbs. per acre. Make first application when shoots are 6 inches to 8 inches long. Make a second application when shoots are 12 inches to 15 inches long and the third about 14 days later. Repeat applications at 10 to 14 day intervals to protect new growth. Avoid applications during hot weather as injury may occur. Concord and other labrusca type grapes may be injured by sulfur.

F-473

ACTIVE INGREDIENT:

Sulfur 60%

INERT INGREDIENTS: 40%

TOTAL 100%

EPA REG. NO. 2935-246

CAUTION: KEEP OUT OF REACH OF CHILDREN!

CAUTION: Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage when the temperature is high (90° or above). Do not make sulfur applications at such times. Do not use this product for any purpose other than that recommended on this label. Never use sulfur before, with, or immediately following foliage oil sprays.

Do not reuse container. Destroy when empty.

RECOMMENDATIONS FOR USE: (Continued)

HOPS: For control of Red Spider Mites use 25 lbs. per acre. For prevention of Brown Rot on plums and prunes use 40 to 50 lbs. per acre. Apply when fruit starts to ripen.

PEACHES: For control of Powdery Mildew and prevention of Brown Rot use 40 to 50 lbs. per acre. For prevention of Brown Rot apply when fruit starts to ripen. For the control of Peach Silver Mite, use 40 to 50 lbs. per acre.

BEST DOCUMENT AVAILABLE